arXiv:1207.7233v1 [math.AG] 31 Jul 2012

FOURIER-MUKAI AND AUTODUALITY FOR COMPACTIFIED JACOBIANS. I

MARGARIDA MELO, ANTONIO RAPAGNETTA, FILIPPO VIVIANI

(WITH AN APPENDIX BY ANA CRISTINA LÓPEZ MARTÍN)

Abstract. To every singular reduced projective curve X one can associate, following E. Esteves, many fine compactified Jacobians, depending on the choice of a polarization on X, each of which yields a modular compactification of a disjoint union of the generalized Jacobian of X. We prove that, for a reduced curve with locally planar singularities, the Fourier-Mukai transform with kernel the Poincaré sheaf from the derived category of the generalized Jacobian of X to the derived category of any fine compactified Jacobian of X is fully faithful, generalizing a previous result of D. Arinkin in the case of integral curves. As a consequence, we prove that there is a canonical isomorphism (called autoduality) between the generalized Jacobian of X and the connected component of the identity of the Picard scheme of any fine compactified Jacobian of X and that algebraic equivalence and numerical equivalence coincide on any fine compactified Jacobian, generalizing previous results of Arinkin, Esteves, Gagné, Kleiman, Rocha.

The paper contains two Appendices: in the first one, we explain how our work can be interpreted in view of the Langlands duality for the Hitchin fibration as proposed by Donagi-Pantev; the second one (due to Ana Cristina López Martín) contains a fully faithful criterion for Fourier-Mukai transforms that we use in the proof of our main theorem.

Contents

1. Introduction 1
2. Compactified Jacobians of reduced curves 8
3. Deformation theory 13
4. The universal fine compactified Jacobians 18
5. The Picard scheme of the universal fine compactified Jacobians 23
6. Abel maps 24
7. Examples: Locally planar curves of arithmetic genus 1 31
8. The Poincaré bundle 34
9. Cohomology of restricted Poincaré bundles 39
10. Proof of Theorem C for nodal curves 41
11. Proof of Theorem A, Corollary B and Theorem C 43
12. Proof of Theorem D 48
13. Appendix A: Hitchin fibration vs compactified Jacobians of spectral curves 53
14. Appendix B: Fully faithfulness criteria (by Ana Cristina López Martín) 55
References 59

1. Introduction

Let C be a smooth irreducible projective curve over an algebraically closed field k and let $J(C)$ be its Jacobian variety. Then $J(C)$ is an abelian variety that carries lots of information about the curve itself. Among abelian varieties, Jacobians have the important property of being "autodual", i.e., they are canonically isomorphic to their dual abelian varieties. This is equivalent to the existence of a Poincaré line bundle $\mathcal{P}$ on $J(C) \times J(C)$ which is universal as a family of algebraically trivial line bundles on $J(C)$. In the breakthrough work [Muk81], S. Mukai proved that the Fourier-Mukai transform with kernel $\mathcal{P}$ is an auto-equivalence of the bounded derived category of $J(C)$ [1].

Key words and phrases. Compactified Jacobians, Fourier-Mukai transform, autoduality, Poincaré bundle, Abel map.

[1] More generally, for an arbitrary abelian variety A with dual abelian variety $A^\vee$, Mukai proved that the Fourier-Mukai transform associated to the Poincaré line bundle on $A \times A^\vee$ gives an equivalence between the bounded derived category of A and that of $A^\vee$.

The aim of this paper and its sequel [MRV2] is to extend these results to fine compactified Jacobians (as defined by E. Esteves in [Est01]) of *reduced projective curves with locally planar singularities.* The case of integral (i.e. reduced and irreducible) projective curves with locally planar singularities was dealt with by D. Arinkin in [Ari11] and [Ari], generalizing previous partial results of Esteves-Gagné-Kleiman [EGK02] and Esteves-Kleimann [EK05] for integral projective curves with double point singularities. Moreover, the autoduality result has been recently extended by Esteves-Rocha [ER] to tree-like curves, i.e. curves with locally planar singularities such that the unique singular points lying in more than one irreducible component are separating nodes (e.g. nodal curves of compact type).

The main motivation for this work comes from the (conjectural) Langlands duality for the Hitchin fibration associated to the group GL_r (as introduced by Donagi-Pantev in [DP12]), which in particular predicts a Fourier-Mukai autoequivalence of the bounded derived category of fine compactified Jacobians of spectral curves. In an appendix of this paper, we recall the description of the fibers of the Hitchin fibration in terms of fine compactified Jacobians of spectral curves (see Fact 13.3) andt the (fiberwise) Langlands duality for the Hitchin fibration (see Conjectures 13.5 and 13.6). In the sequel to this work, we will prove Theorem E below, which will give a positive answer to Conjecture 13.6 for reduced spectral curves (i.e. on the so-called regular locus of the Hitchin fibration), extending the work of Arinkin for integral spectral curves (i.e. on the so-called elliptic locus of the Hitchin fibration).

Before stating our results, we need to briefly recall how fine compactified Jacobians of singular curves are defined.

1.1. **Fine compactified Jacobians of singular curves.** Let X be a *reduced* (projective) curve. The ***generalized Jacobian*** $J(X)$ of X is the connected component of the Picard scheme of X containing the identity. It is not difficult to see that $J(X)$ is a smooth irreducible algebraic group of dimension equal to the arithmetic genus $p_a(X)$ of X, parametrizing line bundles on X that have multidegree zero, i.e. degree zero on each irreducible component of X. However, for a singular curve X, the generalized Jacobian $J(X)$ is rarely complete. The problem of compactifying it is very natural and it has attracted the attention of many mathematicians, starting from the pioneering work of Mayer-Mumford and of Igusa in the 50's, till the more recent works of Oda-Seshadri, Altmann-Kleiman, Caporaso, Pandharipande, Simpson, Esteves, etc.. (we refer to the introduction of [Est01] for an account of the different approaches).

Here we will consider Esteves's ***fine compactified Jacobians***, as constructed by E. Esteves in [Est01]. We briefly review the definition of Esteves's fine compactified Jacobians of a reduced curve X (following the notation introduced in [MV12]); we refer the reader to §2.2 for more details. A *polarization* on X is a tuple of rational numbers $\underline{q} = \{\underline{q}_{C_i}\}$, one for each irreducible component C_i of X, such that $|\underline{q}| := \sum_i \underline{q}_{C_i} \in \mathbb{Z}$. A torsion-free rank-1 sheaf I on X of degree $\deg I := \chi(I) - \chi(\mathcal{O}_X)$ equal to $|\underline{q}|$ is called *$\underline{q}$-semistable* (resp. *$\underline{q}$-stable*) if for every proper subcurve $Y \subset X$, we have that

$$\deg_Y(I) \geq \sum_{C_i \subseteq Y} \underline{q}_{C_i} - \frac{\#(Y \cap \overline{X \setminus Y})}{2} \text{ (resp. } >),$$

where $\#(Y \cap \overline{X \setminus Y})$ is the scheme-theoretic intersection of Y with its complementary subcurve $\overline{X \setminus Y}$ and $\deg_Y(I)$ is the degree of the biggest torsion-free quotient of the restriction $I_{|Y}$ of I to the subcurve Y. A polarization $\underline{q}$ is called *general* if there are no strictly $\underline{q}$-semistable sheaves, i.e. if every $\underline{q}$-semistable sheaf is also $\underline{q}$-stable (see Definition 2.6 for a numerical characterization of general polarizations). A fine compactified Jacobian of X is the fine moduli space $\overline{J}_X(\underline{q})$ of torsion-free rank-1 sheaves of degree $|\underline{q}|$ on X that are $\underline{q}$-semistable (or equivalently $\underline{q}$-stable) with respect to a general polarization $\underline{q}$ on X. Indeed, it is known that $\overline{J}_X(\underline{q})$ is a projective scheme over k (see Fact 2.11) on which the generalized Jacobian $J(X)$ of X acts naturally by tensor product. It is easy to see that if the curve X is integral, then all the fine compactified Jacobians of X are isomorphic among them (because all of them are equivalent by translation in the sense of Definition 2.13). However, this fails for reduced non-irreducible curves: the authors have found some explicit examples of nodal curves which admit non-isomorphic fine compactified Jacobians (details will be published somewhere else).

If the curve X has locally planar singularities, then we can prove that any fine compactified Jacobian $\overline{J}_X(\underline{q})$ of X has the following remarkable properties:

- $\overline{J}_X(\underline{q})$ is a reduced scheme with locally complete intersection singularities (see Corollary 2.12(i));

- The smooth locus of $\overline{J}_X(\underline{q})$ coincides with the open subset $J_X(\underline{q}) \subseteq \overline{J}_X(\underline{q})$ parametrizing line bundles; in particular $J_X(\underline{q})$ is dense in $\overline{J}_X(\underline{q})$ and $\overline{J}_X(\underline{q})$ is of pure dimension equal to $p_a(X)$ (see Corollary 2.12(ii) and (iii));
- $\overline{J}_X(\underline{q})$ is connected (see Corollary 4.4);
- $\overline{J}_X(\underline{q})$ has trivial dualizing sheaf (see Corollary 4.5);
- $J_X(\underline{q})$ is the disjoint union of a number of copies of $J(X)$ equal to the complexity $c(X)$ of the curve X (in the sense of Definition 4.10); in particular, $\overline{J}_X(\underline{q})$ has $c(X)$ irreducible components, independently of the chosen polarization $\underline{q}$ (see Corollary 4.12).

In the proof of all the above properties, we use in an essential way the fact that the curve has locally planar singularities and indeed we expect that many of the above properties are false without this assumptions (see also Remark 2.4).

After these preliminaries, we can now state our main results.

1.2. **Main results.** Since any fine compactified Jacobian $\overline{J}_X(\underline{q})$ of X is a fine moduli space for certain sheaves (as the name suggests), there exists a universal sheaf $\mathcal{I}$ on $X \times \overline{J}_X(\underline{q})$. Using this universal sheaf and the formalism of the determinant of cohomology, it is possible to define a Poincaré line bundle $\mathcal{P}$ on $\overline{J}_X(\underline{q}) \times J(X)$; we refer the reader to §8 for details.

Our first result concerns the Fourier-Mukai transform with kernel $\mathcal{P}$. This result can be seen as a first partial generalization of the above mentioned result of Mukai [Muk81] in the case of Jacobians. In Theorem E below (whose proof will appear in [MRV2]), we will give a second and more satisfactory generalization.

Theorem A. *Let X be a reduced curve with locally planar singularities over an algebraically closed field k. Let $J(X)$ be the generalized Jacobian of X and let $\overline{J}_X(\underline{q})$ be a fine compactified Jacobian of X. Denote by $D^b(J(X))$ and $D^b(\overline{J}_X(\underline{q}))$ the bounded derived categories of $J(X)$ and $\overline{J}_X(\underline{q})$, respectively. Let $\mathcal{P}$ be a Poincaré line bundle on $\overline{J}_X(\underline{q}) \times J(X)$. Then the Fourier-Mukai transform (or integral transform) with kernel $\mathcal{P}$*

$$\begin{array}{rcl} \Phi^{\mathcal{P}} : D^b(J(X)) & \longrightarrow & D^b(\overline{J}_X(\underline{q})) \\ \mathcal{E}^\bullet & \longmapsto & \mathbf{R}p_{1*}(p_2^*(\mathcal{E}^\bullet) \otimes \mathcal{P}) \end{array}$$

is fully-faithful, where with p_i we denote the projection of $\overline{J}_X(\underline{q}) \times J(X)$ on the i-th factor.

As a corollary of Theorem A, we can compute the cohomology of the line bundles $\mathcal{P}_M := \mathcal{P}_{|\overline{J} \times \{M\}}$ on $\overline{J}_X(\underline{q})$, as M varies in $J(X)$, generalizing the classical result for abelian varieties (see [Mum70, Sec. 13]).

Corollary B. *Same assumptions as in Theorem A. For any $M \in J(X)$, let $\mathcal{P}_M := \mathcal{P}_{|\overline{J}_X(\underline{q}) \times \{M\}} \in \mathrm{Pic}(\overline{J}_X(\underline{q}))$. Then we have that*

$$H^i(\overline{J}_X(\underline{q}), \mathcal{P}_M) = \begin{cases} 0 & \text{if } M \neq [\mathcal{O}_X], \\ \bigwedge^i H^1(X, \mathcal{O}_X) & \text{if } M = [\mathcal{O}_X]. \end{cases}$$

As we mentioned in the introduction, Jacobians of smooth curves are autodual. In other words, given a smooth projective curve C, its Jacobian $J(C)$ is canonically isomorphic to the dual abelian variety which, by definition, is equal to $\mathrm{Pic}^o(J(C))$, i.e. the connected component of the Picard scheme of $J(C)$ containing the origin. Our next result is a generalization of this autoduality result to fine compactified Jacobians.

Theorem C. *Same assumptions as in Theorem A. The morphism*

$$\begin{aligned} \beta_{\underline{q}} : J(X) &\longrightarrow \mathrm{Pic}^o(\overline{J}_X(\underline{q})) \\ M &\mapsto \mathcal{P}_M := \mathcal{P}_{|\overline{J}_X(\underline{q}) \times \{M\}} \end{aligned}$$

is an isomorphism of algebraic groups.

Finally, it is well known that a line bundle on an abelian variety A is algebraically equivalent to zero if and only if it is numerically equivalent to zero (see [Mum70, Cor. 2, p. 178]). In other words, the connected component $\mathrm{Pic}^o(A)$ of the Picard scheme $\mathrm{Pic}(A)$ of A containing the identity (which also parametrizes line bundles algebraically equivalent to zero) coincides with the open and closed subset $\mathrm{Pic}^\tau(A) \subseteq \mathrm{Pic}(A)$ parametrizing line bundles numerically equivalent to zero. We prove that the same holds true for fine compactified Jacobians.

Theorem D. *Same assumptions as in Theorem A. Then we have that*

$$\mathrm{Pic}^o(\overline{J}) = \mathrm{Pic}^\tau(\overline{J}).$$

Note that the above Theorem D is new even for irreducible curves in positive characteristic: the proof of Theorem D for irreducible curves by D. Arinkin (see [Ari11, Prop. 12]) uses in a crucial way that $\mathrm{char}(k) = 0$.

In our forthcoming paper [MRV2], we will use the results of this paper to prove the following theorem, which can be seen as a strengthening of Theorem A and a further generalization of Mukai's result in [Muk81] to the case of singular curves.

Theorem E ([MRV2])**.** *Same assumptions as in Theorem A. There exists a (naturally defined) Cohen-Macauly sheaf* $\overline{\mathcal{P}}$ *on* $\overline{J}_X(\underline{q}) \times \overline{J}_X(\underline{q})$ *such that the Fourier-Mukai transform (or integral transform) with kernel* $\overline{\mathcal{P}}$

$$\begin{aligned} \Phi^{\overline{\mathcal{P}}} : D^b(\overline{J}_X(\underline{q})) &\longrightarrow D^b(\overline{J}_X(\underline{q})) \\ \mathcal{E}^\bullet &\longmapsto \mathbf{R}p_{1*}(p_2^*(\mathcal{E}^\bullet) \overset{\mathbf{L}}{\otimes} \overline{\mathcal{P}}) \end{aligned}$$

is an equivalence.

Note that Theorem E provides a positive answer to Conjecture 13.6 for reduced spectral curves, at least in the case where the degree d and the rank r of the Higgs bundles are coprime (see Remark 13.4).

1.3. **Sketch of the proofs.** Let us now give a brief outline of the proofs of the main results, trying to highlight the main ingredients that we use.

Theorem A follows (using a fully-faithfulness criterion for Fourier-Mukai transforms due to A. C. López Martín; see Appendix B) from the formula

$$Rp_{2*}\mathcal{P} = \mathbf{k}(0)[-g] \tag{1.1}$$

where $\mathbf{k}(0)$ denotes the skyscraper sheaf supported at the origin $0 = [\mathcal{O}_X] \in J(X)$, $g = p_a(X)$ is the arithmetic genus of X and $p_2 : \overline{J}_X(\underline{q}) \times J(X) \to J(X)$ is the projection onto the second factor. Indeed, formula (1.1) is a generalization of a well-known result of Mumford (see [Mum70, Sec. III.13]) for abelian varieties which was indeed the crucial step for the celebrated original result of Mukai [Muk81].

In order to prove (1.1), the key idea, which we learned from D. Arinkin in [Ari11] and [Ari][2], is to prove a similar formula for the effective semiuniversal deformation family of X (see §3.3 for more details):

$$\begin{tikzcd} X \arrow[r, hook] \arrow[d] \arrow[dr, phantom, "\square"] & \mathcal{X} \arrow[d, "\pi"] \\ \mathrm{Spec}\, \mathrm{k} \arrow[r, hook] & \mathrm{Spec}\, \mathrm{R_X}. \end{tikzcd}$$

The generalized Jacobian $J(X)$ and the fine compactified Jacobian $\overline{J}_X(\underline{q})$ deform over $\mathrm{Spec}\, \mathrm{R_X}$ to, respectively, the universal generalized Jacobian $v : J(\mathcal{X}) \to \mathrm{Spec}\, \mathrm{R_X}$ (see Fact 4.6) and the universal fine compactified Jacobian $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{Spec}\, \mathrm{R_X}$ with respect to the polarization $\underline{q}$ (see Theorem 4.2). Therefore we get the following commutative diagram

$$\begin{tikzcd} & \overline{J}_{\mathcal{X}}(\underline{q}) \times_{\mathrm{Spec}\, \mathrm{R_X}} J(\mathcal{X}) \arrow[dl, "\widetilde{u}"'] \arrow[dr, "\widetilde{v}"] & \\ J(\mathcal{X}) \arrow[dr, "v"] \arrow[rr, phantom, "\square"] & & \overline{J}_{\mathcal{X}}(\underline{q}) \arrow[dl, "u"'] \\ & \mathrm{Spec}\, \mathrm{R_X} \arrow[ul, bend left, "\zeta"] & \end{tikzcd} \tag{1.2}$$

where ζ is the zero section of v. Moreover, the Poincaré line bundle $\mathcal{P}$ on $\overline{J}_X(\underline{q}) \times J(X)$ deforms to the universal Poincaré line bundle $\mathcal{P}^{\mathrm{un}}$ on the fiber product $\overline{J}_{\mathcal{X}}(\underline{q}) \times_{\mathrm{Spec}\, \mathrm{R_X}} J(\mathcal{X})$ (see §10).

---

[2] In loc. cit., D. Arinkin considers the stack of all integral curves with locally planar singularities, which is of finite type. Here, we replace this stack with the semiuniversal deformation space of X since the stack of all reduced curves with locally planar singularities is not of finite type.

Equation (1.1) will follow, by restricting to the central fiber of v, from the following universal version of it (which we prove in Theorem 11.1):

$$R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) = \zeta_*(\mathcal{O}_{\mathrm{Spec}\,\mathrm{R_X}})[-g]. \tag{1.3}$$

A key intermediate step in proving (1.3) consists in showing that

(*) $\quad R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) = R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})[-g]$ is a Cohen-Macaulay sheaf such that $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) = \mathrm{Im}(\zeta)$.

The proof of (*) has two main ingredients. The first ingredient is the study of the cohomology of the line bundles $\mathcal{P}_M \in \mathrm{Pic}(\overline{J}_X(\underline{q}))$, for $M \in J(X)$; see Section 9. Here, we use in an essential way the Abel map $A_L : X \longrightarrow \overline{\mathbb{J}}_X$, for $L \in \mathrm{Pic}(X)$, with value in the scheme $\overline{\mathbb{J}}_X$ parametrizing all simple torsion-free rank-1 sheaves on X; see section 6. The second ingredient is the equigeneric stratification of $\mathrm{Spec}\,\mathrm{R_X}$, i.e. the stratification of $\mathrm{Spec}\,\mathrm{R_X}$ according to the arithmetic genus of the normalization of the fibers of the universal family $\mathcal{X} \to \mathrm{Spec}\,\mathrm{R_X}$. As shown by Teissier [Tei80] and Diaz-Harris [DH88], if X has locally planar singularities (as in our case) then each equigeneric stratum has the expected codimension and all its generic points correspond to nodal curves (see Fact 3.3). These properties allow us to prove (*) over the generic points of each equigeneric stratum, using in an essential way Theorem C for nodal curves; see Section 10.

The proof of **Theorem C** follows the same idea of using the semiuniversal deformation family $\mathcal{X} \to \mathrm{Spec}\,\mathrm{R_X}$ of X. The map $\beta_{\underline{q}}$ of Theorem C deforms over $\mathrm{Spec}\,\mathrm{R_X}$ to a homomorphism

$$\beta_{\underline{q}}^{\mathrm{un}} : J(\mathcal{X}) \to \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \tag{1.4}$$

between two group schemes which are smooth, separated and of finite type over $\mathrm{Spec}\,\mathrm{R_X}$ (see Fact 4.6 and Theorem 5.1(iv)). We note here that the representability of $\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$ and its smoothness over $\mathrm{Spec}\,\mathrm{R_X}$ (proved in Theorem 5.1(iv)) use in a crucial way the fact that

$$h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X). \tag{**}$$

Formula (**) can be proved for nodal curves using results of Alexeev-Nakamura [AN99] and Alexeev [Ale04] on some special degenerations of abelian varieties called stable quasiabelian varieties (this is exactly what allows us to prove Theorem C for nodal curves in Section 10 before proving Theorem A). For an arbitrary curve X with locally planar singularities, formula (**) follows from Corollary B, hence from the Fourier-Mukai type result of Theorem A. A direct proof of (**) would allow to give a Fourier-Mukai's free proof of Theorem C (and also of Theorem D as we will see below).

In Theorem 10.2, we prove that the map $\beta_{\underline{q}}^{\mathrm{un}}$ is an isomorphism (assuming that (**) holds true), which therefore implies Theorem C restricting to the closed point of $\mathrm{Spec}\,\mathrm{R_X}$. The proof of Theorem 10.2 uses the fact (due to Esteves-Gagné-Kleiman [EGK02]) that $\beta_{\underline{q}}^{\mathrm{un}}$ is an isomorphism over the open subset $U \subseteq \mathrm{Spec}\,\mathrm{R_X}$ (whose complement has codimension at least two by Lemma 3.5) of curves having at most one node, which combined with Van der Waerden's theorem on the purity of the ramification locus and Zariski's main theorem, gives that $\beta_{\underline{q}}^{\mathrm{un}}$ is an open embedding, hence an isomorphism.

Finally, let us give a sketch of the proof of **Theorem D**, which we be given in Section 12.

In Theorem 12.1, we will first prove Theorem D in the special case where the fine compactified Jacobian $\overline{J}_X(\underline{q})$ admits an Abel map, i.e. if there exists $L \in \mathrm{Pic}(X)$ such that $\mathrm{ImA_L} \subseteq \overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$ (see Definition 6.8). Indeed, this hypothesis is quite restrictive for a fine compactified Jacobian since in general only of them will admit an Abel map (see e.g. Proposition 7.4). Once again, the strategy will be to work on the semiuniversal deformation family $\mathcal{X} \to \mathrm{Spec}\,\mathrm{R_X}$. Indeed, we can deform the line bundle $L \in \mathrm{Pic}(X)$ that gives the Abel map $A_L : X \to \overline{J}_X(\underline{q})$ to a line bundle $\mathcal{L}$ on $\mathcal{X}$ in order to obtain a universal Abel map $A_{\mathcal{L}} : \mathcal{X} \to \overline{J}_{\mathcal{X}}(\underline{q})$. By taking the pull-back via $A_{\mathcal{L}}$, we obtain the following commutative diagram of group schemes (all of which are smooth, separated and of finite type over $\mathrm{Spec}\,\mathrm{R_X}$, by Fact 4.6 and Theorem 5.1):

(1.5)

$$\begin{tikzcd}
\mathrm{Pic}^{\tau}(\overline{J}_{\mathcal{X}}(\underline{q})) \arrow[dr, two heads, "A_{\mathcal{L}}^{*,\tau}"] & \\
& J(\mathcal{X}) \\
\mathrm{Pic}^{o}(\overline{J}_{\mathcal{X}}(\underline{q})) \arrow[uu, hook, "i"] \arrow[ur, "\cong", "A_{\mathcal{L}}^{*,o}"'] &
\end{tikzcd}$$

where i is the natural open embedding and $A_{\mathcal{L}}^{*,o}$ is an isomorphism since it is the right inverse of $\beta_{\underline{q}}^{\mathrm{un}}$ (by Proposition 8.5), which is an isomorphism by Theorem 10.2. The morphism $A_{\mathcal{L}}^{*,\tau}$ is an isomorphism over the open subset $U \subseteq \operatorname{Spec} \mathrm{R_X}$ of curves having at most one node (as it follows from [EGK02]); using that $\operatorname{Spec} \mathrm{R_X} \setminus \mathrm{U}$ has codimension at least two, together with Van der Waerden's theorem on the purity of the ramification locus and Zariski's main theorem, we conclude that $A_{\mathcal{L}}^{*,\tau}$ is an open embedding, hence an isomorphism. Therefore i must be an equality and Theorem D in this special case follows by restricting to the closed point of $\operatorname{Spec} \mathrm{R_X}$.

In order to prove Theorem D in the general case, i.e. if $\overline{J}_X(\underline{q})$ does not admit an Abel map, we consider another fine compactified Jacobian $\overline{J}_X(\underline{q}')$ of X that does admit an Abel map (such a fine compactified Jacobian $\overline{J}_X(\underline{q}')$ exists by Corollary 6.9(ii)) and we are able to deduce Theorem D for $\overline{J}_X(\underline{q})$ knowing that it does hold true for $\overline{J}_X(\underline{q}')$ (by Theorem 12.1). The key ingredient is to compare their universal fine compactified Jacobians $\overline{J}_{\mathcal{X}}(\underline{q})$ and $\overline{J}_{\mathcal{X}}(\underline{q}')$ by showing that their are isomorphic over the open subset $U \subseteq \operatorname{Spec} \mathrm{R_X}$ of curves having at most one node (see Lemma 12.2). We refer to Section 12 for more details.

1.4. **Outline of the paper.** The paper is organized as follows.

Section 2 is devoted to collect several facts on fine compactified Jacobians of reduced curves: in §2.1, we consider the scheme $\overline{\mathbb{J}}_X$ parametrizing all simple torsion-free rank-1 sheaves on a curve X (see Fact 2.2) and we investigate its properties under the assumption that X has locally planar singularities (see Theorem 2.3); in §2.2, we introduce fine compactified Jacobians of X (see Fact 2.11) and study them under the assumption that X has locally planar singularities (see Corollary 2.12).

Section 3 is devoted to recall or prove some basic facts on deformation theory: we start with the deformation theory of a curve X (see §3.1) and then we continue with the deformation theory of a pair (X, I) consisting of a curve X together with a torsion-free, rank-1 sheaf I on X (see §3.2). Finally, in §3.3, we study the semiuniversal deformation spaces for a curve X and for a pair (X, I) as above.

In Section 4, we introduce the universal fine compactified Jacobians relative to the semiuniversal deformation of a curve X (see Theorem 4.2) and study it under the assumption that X has locally planar singularities (see Theorem 4.3). In §4.1, we use a result of J. Kass in order to prove that the pull-back of any universal fine compactified Jacobian under a 1-parameter regular smoothing of the curve is a compactification of the Néron model of the Jacobian of the general fiber (see Fact 4.9); a formula for the number of irreducible components of a fine compactified Jacobian follows (see Corollary 4.12).

Section 5 is devoted to study the representability of the relative Picard scheme of the universal fine compactified Jacobians and of its subfunctors parametrizing line bundles that are fiberwise algebraically or numerically equivalent to the trivial line bundle (see Theorem 5.1).

In Section 6, we introduce Abel maps: first for curves that do not have separating points (see §6.1) and then for curves all of whose separating points are nodes, e.g. Gorenstein curves (see §6.2).

In Section 7, we illustrate the general theory developed so far with the study of fine compactified Jacobians of Kodaira curves, i.e. curves of arithmetic genus one and with locally planar singularities.

In Section 8, we define the Poincaré line bundle and study its behavior with respect to the Abel maps (see Proposition 8.5).

In Section 9, we study the cohomology of the restricted Poincaré line bundles on a fine compactified Jacobian, obtaining some partial results that later on will turn out to be special cases of Corollary B.

Section 10 contains a proof of Theorem C for nodal curves while Section 11 contains the proof of Theorem A, Corollary B and the general case of Theorem C. Finally, Theorem D is proved in Section 12.

In the Appendix 13, we first discuss the Hitchin fibration and the description of its fibers in terms of compactified Jacobians of spectral curves (see Fact 13.3). Then we state the conjectural Langlands duality for the Hitchin fibration (see Conjecture 13.5) and its implication for the autoduality of the compactified Jacobians of spectral curves (see Conjecture 13.6).

Appendix 14, by A. C. López Martín, contains fully-faithfulness criteria for integral functors from a smooth connected quasi-projective scheme to a Gorenstein connected projective scheme, generalizing the classical Bondal-Orlov criterion.

The following notations will be used throughout the paper.

Notations.

1.1. k will denote an algebraically closed field (of arbitrary characteristic), unless otherwise stated. All **schemes** are k-schemes, and all morphisms are implicitly assumed to respect the k-structure.

1.2. A **curve** is a *reduced* projective scheme over k of pure dimension 1. Unless otherwise specified, a curve is meant to be connected.

Given a curve X, we denote by X_{sm} the smooth locus of X, by X_{sing} its singular locus and by $\nu : X^\nu \to X$ the normalization morphism. We denote by $\gamma(X)$, or simply by γ where there is no danger of confusion, the number of irreducible components of X.

We denote by $p_a(X)$ the *arithmetic genus* of X, i.e. $p_a(X) := 1-\chi(\mathcal{O}_X) = 1-h^0(X,\mathcal{O}_X)+h^1(X,\mathcal{O}_X)$. We denote by $g^\nu(X)$ the *geometric genus* of X, i.e. the sum of the genera of the connected components of the normalization X^ν.

1.3. A **subcurve** Z of a curve X is a closed k-scheme $Z \subseteq X$ that is reduced and of pure dimension 1. We say that a subcurve $Z \subseteq X$ is proper if $Z \neq \emptyset, X$.

Given two subcurves Z and W of X without common irreducible components, we denote by $Z \cap W$ the 0-dimensional subscheme of X that is obtained as the scheme-theoretic intersection of Z and W and we denote by $|Z \cap W|$ its length.

Given a subcurve $Z \subseteq X$, we denote by $Z^c := \overline{X \setminus Z}$ the **complementary subcurve** of Z and we set $\delta_Z = \delta_{Z^c} := |Z \cap Z^c|$.

1.4. A curve X is called **Gorenstein** if its dualizing sheaf ω_X is a line bundle.

1.5. A curve X has **locally complete intersection (l.c.i.) singularities at** $p \in X$ if the completion $\widehat{\mathcal{O}}_{X,p}$ of the local ring of X at p can be written as

$$\widehat{\mathcal{O}}_{X,p} = k[[x_1,\ldots,x_r]]/(f_1,\ldots,f_{r-1}),$$

for some $r \geq 2$ and some $f_i \in k[[x_1,\ldots,x_r]]$. A curve X has locally complete intersection (l.c.i.) singularities if X is l.c.i. at every $p \in X$. Clearly, a curve with l.c.i. singularities is Gorenstein.

1.6. A curve X has **locally planar singularities at** $p \in X$ if the completion $\widehat{\mathcal{O}}_{X,p}$ of the local ring of X at p has embedded dimension two, or equivalently if it can be written as

$$\widehat{\mathcal{O}}_{X,p} = k[[x,y]]/(f),$$

for a reduced series $f = f(x,y) \in k[[x,y]]$. A curve X has locally planar singularities if X has locally planar singularities at every $p \in X$. Clearly, a curve with locally planar singularities has l.c.i. singularities, hence it is Gorenstein.

1.7. A **separating point** of a curve X is a geometric point $n \in X$ for which there exists a subcurve $Z \subset X$ such that $\delta_Z = 1$ and $Z \cap Z^c = \{n\}$. If X is Gorenstein, then a separating point n of X is a node of X, i.e. $\widehat{\mathcal{O}}_{X,n} = k[[x,y]]/(xy)$ (see Fact 6.4). However this is false in general without the Gorenstein assumption (see Example 6.5).

1.8. Given a scheme S proper over a field k (not necessarily algebraically closed), we denote by $\text{Pic}(S)$ its **Picard scheme**, which exists by a result of Murre (see [FGA05, Chap. 9, Cor. 4.1.18] and the references therein). The **connected component of the identity** of $\text{Pic}(S)$, denoted by $\text{Pic}^o(S)$, parametrizes line bundles on S which are algebraically equivalent to the trivial line bundle (see [FGA05, Chap. 9, Sec. 5] for details). The **torsion component of the identity** of $\text{Pic}(S)$, denoted by $\text{Pic}^\tau(S)$, parametrizes line bundles on S which are numerically equivalent to the trivial line bundle or, equivalently, such that some powers of them lie in $\text{Pic}^o(S)$ (see [FGA05, Chap. 9, Sec. 6] for details). The scheme $\text{Pic}^\tau(S)$ is an open and closed group subscheme of $\text{Pic}(S)$ which is of finite type over k (see [FGA05, Prop. 9.6.12]).

On the other hand, given an arbitrary scheme S, we denote by $\mathcal{P}ic(S)$ the **Picard group** of S, i.e. the abstract group consisting of all isomorphism classes of line bundles on S with the operation of tensor product.

1.9. Given a curve X, the **generalized Jacobian** of X, denoted by $J(X)$ or by $\text{Pic}^{\underline{0}}(X)$, is the algebraic group whose k-valued points are the group of line bundles on X of multidegree $\underline{0}$ (i.e. having degree 0 on each irreducible component of X) together with the multiplication given by the tensor product. The generalized Jacobian of X is a connected commutative smooth algebraic group of dimension equal to $h^1(X,\mathcal{O}_X)$ which coincides with $\text{Pic}^o(X)$ and with $\text{Pic}^\tau(X)$.

1.10. Given a scheme X, we will denote by $D(X)$ the **derived category** of complexes of $\mathcal{O}_X$-modules with quasi-coherent cohomology sheaves and by $D^b(X) \subset D(X)$ the **bounded derived category** consisting of complexes with only finitely many non-zero cohomology sheaves.

1.11. Given a scheme X and a closed point $x \in X$, we will denote by $\mathbf{k}(x)$ the **skyscraper sheaf** supported at x.

2. Compactified Jacobians of reduced curves

The aim of this section is to collect several facts about compactified Jacobians of connected reduced curves, with special emphasis on connected reduced curves with locally planar singularities. Many of these facts are well-known to the experts but for many of them we could not find satisfactory references in the existing literature, at least at the level of generality we need, e.g. for reducible curves. Throughout this section, we fix a connected reduced curve X.

2.1. **Simple rank-1 torsion-free sheaves.**

We start by defining the sheaves on the connected curve X we will be working with.

Definition 2.1. A coherent sheaf I on a connected curve X is said to be:

(i) *rank*-1 if I has generic rank 1 at every irreducible component of X;
(ii) *torsion-free* if $\mathrm{Supp(I)} = \mathrm{X}$ and every non-zero subsheaf $J \subseteq I$ is such that $\dim \mathrm{Supp(J)} = 1$;
(iii) *simple* if $\mathrm{End}_k(I) = k$.

Note that any line bundle on X is a simple rank-1 torsion-free sheaf.

Consider the functor

$$\overline{\mathbb{J}}_X^* : \{\text{Schemes}/k\} \to \{\text{Sets}\} \tag{2.1}$$

which associates to a k-scheme T the set of isomorphism classes of T-flat, coherent sheaves on $X \times_k T$ whose fibers over T are simple rank-1 torsion-free sheaves. The functor $\overline{\mathbb{J}}_X^*$ contains the open subfunctor

$$\mathbb{J}_X^* : \{\text{Schemes}/k\} \to \{\text{Sets}\} \tag{2.2}$$

which associates to a k-scheme T to the set of isomorphism classes of line bundles on $X \times_k T$.

Fact 2.2 (Murre-Oort, Altman-Kleiman, Esteves)**.** *Let X be a connected reduced curve. Then*

(i) The étale sheafification of $\mathbb{J}_X^*$ is represented by a k-scheme $\mathrm{Pic}(X) = \mathbb{J}_X$, locally of finite type over k. Moreover, $\mathbb{J}_X$ is smooth.
(ii) The étale sheafification of $\overline{\mathbb{J}}_X^*$ is represented by a k-scheme $\overline{\mathbb{J}}_X$, locally of finite type over k. Moreover, $\overline{\mathbb{J}}_X$ is universally closed and $\mathbb{J}_X$ is an open subset of $\overline{\mathbb{J}}_X$.
(iii) There exists a sheaf $\mathcal{I}$ on $X \times \overline{\mathbb{J}}_X$ such for every $\mathcal{F} \in \overline{\mathbb{J}}_X^*(T)$ there exists a unique map $\alpha_{\mathcal{F}} : T \to \overline{\mathbb{J}}_X$ with the property that $\mathcal{F} = (\mathrm{id_X} \times \alpha_{\mathcal{F}})^*(\mathcal{I}) \otimes \pi_2^*(\mathrm{N})$ for some $N \in \mathrm{Pic}(T)$, where $\pi_2 : X \times T \to T$ is the projection onto the second factor. The sheaf $\mathcal{I}$ is uniquely determined up to tensor product with the pullback of an invertible sheaf on $\overline{\mathbb{J}}_X$ and it is called a universal sheaf.

Proof. Part (i): the representability of the étale sheafification of $\mathbb{J}_X^*$ follows from a result of Murre-Oort (see [BLR90, Sec. 8.2, Thm. 3] and the references therein). The smoothness of $\mathbb{J}_X$ follows from [BLR90, Prop. 2].

Part (ii): the representability of the étale sheafification of $\overline{\mathbb{J}}_X^*$ by an algebraic space $\overline{\mathbb{J}}_X$ locally of finite type over k follows from a general result of Altmann-Kleiman ([AK80, Thm. 7.4]). The fact that $\overline{\mathbb{J}}_X$ is a scheme follows from a general result of Esteves ([Est01, Thm. B]), using the fact that each irreducible component of X has a k-point (recall that k is assumed to be algebraically closed). Since $\mathbb{J}_X^*$ is an open subfunctor of $\overline{\mathbb{J}}_X^*$ then $\mathbb{J}_X$ is an open subscheme of $\overline{\mathbb{J}}_X$. Finally, the fact that $\overline{\mathbb{J}}_X$ is universally closed follows from [Est01, Thm. 32].

Part (iii) follows from another general result of Altman-Kleiman ([AK79, Thm. 3.4]) using again the fact that each irreducible component of X has a k-point.

$\square$

Given a coherent sheaf I on X, its degree $\deg(I)$ is defined by $\deg(I) := \chi(I) - \chi(\mathcal{O}_X)$, where $\chi(I)$ (resp. $\chi(\mathcal{O}_X)$) denotes the Euler-Poincaré characteristic of I (resp. of the trivial sheaf $\mathcal{O}_X$). Since the degree of a sheaf is constant under deformations, we get a decomposition

$$\begin{cases} \overline{\mathbb{J}}_X = \coprod\limits_{d \in \mathbb{Z}} \overline{\mathbb{J}}_X^d, \\ \mathbb{J}_X = \coprod\limits_{d \in \mathbb{Z}} \mathbb{J}_X^d = \coprod\limits_{\underline{d} \in \mathbb{Z}^{\gamma(X)}} \mathbb{J}_X^{\underline{d}}, \end{cases} \tag{2.3}$$

where $\overline{\mathbb{J}}_X^d$ (resp. $\mathbb{J}_X^d$, resp. $\mathbb{J}_X^{\underline{d}}$) denotes the open and closed subscheme of $\overline{\mathbb{J}}_X$ (resp. $\mathbb{J}_X$, resp. $\mathbb{J}_X^{\underline{d}}$) parametrizing simple rank-1 torsion-free sheaves I (resp. line bundles L) of degree $\deg I = d$ (resp. $\deg L = d$, resp. $\underline{\deg} L = \underline{d}$).

If X has locally planar singularities, then $\overline{\mathbb{J}}_X$ has the following properties.

Theorem 2.3. *Let X be a connected reduced curve with locally planar singularities. Then*

- *(i) $\overline{\mathbb{J}}_X$ is a reduced scheme with locally complete intersection singularities.*
- *(ii) $\mathbb{J}_X$ is dense in $\overline{\mathbb{J}}_X$.*
- *(iii) $\mathbb{J}_X$ is the smooth locus of $\overline{\mathbb{J}}_X$.*

Proof. The required properties of $\overline{\mathbb{J}}_X$ will be deduced from the analogous properties of the Hilbert scheme of X via the Abel map.

Denote by $\mathrm{Hilb}^d(X)$ the Hilbert scheme parametrizing subschemes D of X of finite length $d \geq 0$, or equivalently ideal sheaves $I \subset \mathcal{O}_X$ such that $\mathcal{O}_X/I$ is a finite scheme of length d. Note that if $[I \subset \mathcal{O}_X] \in \mathrm{Hilb}^d(X)$ then I is a torsion-free rank-1 sheaf on X. However, in general, I is neither simple (unless X is irreducible) nor a line bundle (unless X is smooth). We refer to [Est01, Exa. 38] for an example of $[I \subset \mathcal{O}_X] \in \mathrm{Hilb}^d(X)$ with I not simple. We introduce the following subschemes of $\mathrm{Hilb}^d(X)$:

$$\begin{cases} \mathrm{Hilb}^d(X)_s := \{[I \subset \mathcal{O}_X] \in \mathrm{Hilb}^d(X) \ : \ I \text{ is simple}\}, \\ \mathrm{Hilb}^d(X)_l := \{[I \subset \mathcal{O}_X] \in \mathrm{Hilb}^d(X) \ : \ I \text{ is a line bundle}\}. \end{cases}$$

By [AK80, Prop. 5.15], the natural inclusions

$$\mathrm{Hilb}^d(X)_l \subseteq \mathrm{Hilb}^d(X)_s \subseteq \mathrm{Hilb}^d(X)$$

are open inclusions.

If X is a reduced curve with locally planar singularities, then the Hilbert scheme $\mathrm{Hilb}^d(X)$ has the following properties (which are inherited by its open subscheme $\mathrm{Hilb}^d(X)_s$):

- (a) $\mathrm{Hilb}^d(X)$ (and hence $\mathrm{Hilb}^d(X)_s$) is reduced with locally complete intersection singularities by [AIK76, Cor. 7] (see also [BGS81, Prop. 1.4]);
- (b) $\mathrm{Hilb}^d(X)_l$ is dense in $\mathrm{Hilb}^d(X)$ (and hence in $\mathrm{Hilb}^d(X)_s$) by [AIK76, Thm. 8] (see also [BGS81, Prop. 1.4]);
- (c) $\mathrm{Hilb}^d(X)_l$ is the smooth locus of $\mathrm{Hilb}^d(X)$ (and hence also of $\mathrm{Hilb}^d(X)_s$) by [BGS81, Prop. 2.3].

Now, given a line bundle M on X, we define the M-twisted Abel map of degree d by

$$\begin{aligned} A_M^d : \mathrm{Hilb}^d(X)_s &\longrightarrow \overline{\mathbb{J}}_X, \\ [I \subset \mathcal{O}_X] &\mapsto I \otimes M. \end{aligned} \tag{2.4}$$

Note that, by definition, it follows that

$$(A_M^d)^{-1}(\mathbb{J}_X) = \mathrm{Hilb}^d(X)_l. \tag{2.5}$$

We claim that, locally on the target, the M-twisted Abel map of degree d is smooth and surjective (for suitable choices of $M \in \mathrm{Pic}(X)$ and $d \geq 0$), or more precisely

CLAIM: For every point $I_0 \in \overline{\mathbb{J}}_X$, there exists an open subset $U \subseteq \overline{\mathbb{J}}_X$ containing I_0, an open subset $V \subseteq \mathrm{Hilb}^d(X)_s$ for some $d \geq 0$ and a line bundle $M \in \mathrm{Pic}(X)$ such that $A_M^d(V) = U$ and $(A_M^d)_{|V}$ is smooth.

Before proving the claim, let us see how it implies the theorem. Observe that each of the three statements of the theorem is local in $\overline{\mathbb{J}}_X$, i.e. it is sufficient to check it on an open cover of $\overline{\mathbb{J}}_X$. Since the open subsets U of the claim, as I_0 varies in $\overline{\mathbb{J}}_X$, cover $\overline{\mathbb{J}}_X$, it is enough to prove the desired properties on an open subset U as in the claim.

Part (i): from the above property (a), it follows that V is reduced with locally complete intersection singularities. Since $(A_M^d)_{|V}$ is smooth and surjective, also U inherits the same properties.

Part (ii): from the above property (b), it follows that $\mathrm{Hilb}^d(X)_l \cap V$ is dense in V. From the surjectivity of $(A_M^d)_{|V}$ together with (2.5), it follows that $A_M^d(V \cap \mathrm{Hilb}^d(X)_l) = U \cap \mathbb{J}_X$ is dense in $A_M^d(V) = U$.

Part (iii): from the above property (c), it follows that $\mathrm{Hilb}^d(X)_l \cap V$ is the smooth locus of V. Since $(A_M^d)_{|V}$ is smooth and surjective and (2.5) holds, we infer that $A_M^d(V \cap \mathrm{Hilb}^d(X)_l) = U \cap \mathbb{J}_X$ is the smooth locus of $A_M^d(V) = U$.

It remains to prove the claim. Suppose that $I_0 \in \overline{\mathbb{J}}_X^e$ for a certain $e \in \mathbb{Z}$. Fix an ample invertible sheaf $\mathcal{O}_X(1)$ on X and let $a := \deg \mathcal{O}_X(1)$. Fix an integer $t >> 0$ such that, denoting by I_0^* the dual of I_0 (i.e. $I_0^* := \mathcal{H}om(I_0, \mathcal{O}_X)$) we have that

$$I_0^*(t) := I_0^* \otimes \mathcal{O}_X(t) \quad \text{is globally generated,} \tag{2.6}$$

$$\mathrm{Ext}^1(\mathrm{I}_0(-\mathrm{t}), \mathcal{O}_\mathrm{X}) = 0. \tag{2.7}$$

Let U be the locus of $\overline{\mathbb{J}}_X^e$ consisting of the sheaves $I \in \overline{\mathbb{J}}_X^e$ satisfying (2.6) and (2.7). Clearly, U is an open subset of $\overline{\mathbb{J}}_X^e$ containing I_0. Set $d := -e + at$ and $M := \mathcal{O}_X(t)$. Let V the open subset of $\mathrm{Hilb}^d(X)_s$ consisting of the points $[J \subset \mathcal{O}_X] \in \mathrm{Hilb}^d(X)_s$ such that

$$J^* \quad \text{is globally generated,} \tag{2.8}$$

$$\mathrm{Ext}^1(\mathrm{J}, \mathcal{O}_\mathrm{X}) = 0. \tag{2.9}$$

By the definitions of U and V, it is clear that $A_M^d(V) \subseteq U$. Let us show that $A_M^d(V) = U$ and that $(A_M^d)_{|V}$ is smooth.

Given a sheaf $I \in U$, using (2.6) we deduce that there exists an injective homomorphism $\mathcal{O}_X \hookrightarrow I^*(t)$. By taking duals, we get an injective homomorphism $I(-t) \hookrightarrow \mathcal{O}_X$. This defines an element $[I(-t) \subseteq \mathcal{O}_X] \in \mathrm{Hilb}^d(X)_s$, which moreover belongs to V since the properties (2.6) and (2.7) for I are equivalent to the properties (2.8) and (2.9) for $I(-t)$. Since clearly $A_M^d([I(-t) \subseteq \mathcal{O}_X]) = I(-t) \otimes M = I(-t) \otimes \mathcal{O}_X(t) = I$, we get that $I \in \mathrm{Im}(\mathrm{A}_\mathrm{M}^\mathrm{d})$, hence that $A_M^d(V) = U$.

According to [AK80, Thm. 5.18(ii)][3], the Abel map A_M^d is smooth along $(A_M^d)^{-1}(I)$ if $\mathrm{Ext}^1(\mathrm{I} \otimes \mathrm{M}^{-1}, \mathcal{O}_\mathrm{X}) = \mathrm{Ext}^1(\mathrm{I}(-\mathrm{t}), \mathcal{O}_\mathrm{X}) = 0$. Therefore, using (2.7), we get that A_M^d is smooth along $(A_M^d)^{-1}(U)$, hence the restriction $(A_M^d)_{|V}$ is smooth. □

Remark 2.4.

(i) Theorem 2.3 is well-known in the case where X is irreducible (and hence integral): parts (i) and (ii) are due to Altman-Iarrobino-Kleiman [AIK76, Thm. 9]; part (iii) is due to Kleppe [Kle81] (unpublished, for a proof see [Kas09, Prop. 6.4]). Note that, for X irreducible, part (ii) is equivalent to the irreducibility of $\overline{\mathbb{J}}_X^d$ for a certain $d \in \mathbb{Z}$ (hence for all $d \in \mathbb{Z}$).

(ii) The hypothesis that X has locally planar singularities is crucial in the above Theorem 2.3. For example, Altman-Iarrobino-Kleiman constructed in [AIK76, Exa. (13)] an integral curve without locally planar singularities (indeed, a curve which is a complete intersection in $\mathbb{P}^3$) for which $\overline{\mathbb{J}}_X^d$ is not irreducible. Later, Rego ([Reg80, Thm. A]) and Kleppe-Kleiman ([KK81, Thm. 1]) showed that, for X irreducible, $\overline{\mathbb{J}}_X^d$ is irreducible if and only if X has locally planar singularities.

2.2. **Fine compactified Jacobians.**

The functor $\overline{\mathbb{J}}_X$ is not of finite type nor separated over k (and similarly for $\mathbb{J}_X$) if X is not irreducible. However, it can be covered by open subsets that are proper (and even projective) over k: the fine compactified Jacobians of X. The fine compactified Jacobians depend on the choice of a polarization, whose definition is as follows (using the notations of [MV12]).

Definition 2.5. A *polarization* on a connected curve X is a tuple of rational numbers $\underline{q} = \{\underline{q}_{C_i}\}$, one for each irreducible component C_i of X, such that $|\underline{q}| := \sum_i \underline{q}_{C_i} \in \mathbb{Z}$. We call $|\underline{q}|$ the total degree of $\underline{q}$.

Given any subcurve $Y \subseteq X$, we set $\underline{q}_Y := \sum_j \underline{q}_{C_j}$ where the sum runs over all the irreducible components C_j of Y. Note that giving a polarization $\underline{q}$ is the same as giving an assignment $(Y \subseteq X) \mapsto \underline{q}_Y$ such that $\underline{q}_X \in \mathbb{Z}$ and which is additive on Y, i.e. such that if $Y_1, Y_2 \subseteq X$ are two subcurves of X without common irreducible components, then $\underline{q}_{Y_1 \cup Y_2} = \underline{q}_{Y_1} + \underline{q}_{Y_2}$.

Definition 2.6. A polarization $\underline{q}$ is called *integral* at a subcurve $Y \subseteq X$ if $\underline{q}_Z - \frac{\delta_Z}{2} \in \mathbb{Z}$ for any connected component Z of Y and of Y^c.

A polarization is called *general* if it is not integral at any proper subcurve $Y \subset X$.

Remark 2.7. It is easily seen that $\underline{q}$ is general if and only if $\underline{q}_Y - \frac{\delta_Y}{2} \notin \mathbb{Z}$ for any proper subcurve $Y \subset X$ such that Y and Y^c are connected.

---

[3] Note that in loc.cit., this is stated under the assumption that X is integral. However, a close inspection of the proof reveals that this continues to hold true under the assumption that X is only reduced. The irreducibility is only used in part (i) of [AK80, Thm. 5.18].

For each subcurve Y of X and each torsion-free sheaf I on X, the restriction $I_{|Y}$ of I to Y is not necessarily a torsion-free sheaf on Y. However, $I_{|Y}$ contains a biggest subsheaf, call it temporarily J, whose support has dimension zero, or in other words such that J is a torsion sheaf. We denote by I_Y the quotient of $I_{|Y}$ by J. It is easily seen that I_Y is torsion-free on Y and it is the biggest torsion-free quotient of $I_{|Y}$. Moreover, if I is torsion-free rank-1 then I_Y is torsion-free rank-1. We let $\deg_Y(I)$ denote the degree of I_Y on Y, that is, $\deg_Y(I) := \chi(I_Y) - \chi(\mathcal{O}_Y)$.

Definition 2.8. Let $\underline{q}$ be a polarization on X. Let I be a torsion-free rank-1 sheaf on X of degree $d = |\underline{q}|$ (not necessarily simple).

(i) We say that I is *semistable* with respect to $\underline{q}$ (or $\underline{q}$-semistable) if for every proper subcurve $Y \subset X$, we have that

$$\deg_Y(I) \geq \underline{q}_Y - \frac{\delta_Y}{2} \tag{2.10}$$

(ii) We say that I is *stable* with respect to $\underline{q}$ (or $\underline{q}$-stable) if it is semistable with respect to $\underline{q}$ and if the inequality (2.10) is always strict.

Remark 2.9. It is easily seen that a torsion-free rank-1 sheaf I is $\underline{q}$-semistable (resp. $\underline{q}$-stable) if and only if (2.10) is satisfied (resp. is satisfied with strict inequality) for any subcurve $Y \subset X$ such that Y and Y^c are connected.

Lemma 2.10.

(i) If I is stable with respect to a polarization $\underline{q}$ on X then I is simple.

(ii) For a polarization $\underline{q}$ on X, the following conditions are equivalent

  (a) $\underline{q}$ is general.

  (b) Every rank-1 torsion-free sheaf which is $\underline{q}$-semistable is also $\underline{q}$-stable.

  (c) Every simple rank-1 torsion-free sheaf which is $\underline{q}$-semistable is also $\underline{q}$-stable.

  (d) Every line bundle which is $\underline{q}$-semistable is also $\underline{q}$-stable.

Proof. Let us first prove part (i). By contradiction, suppose that I is $\underline{q}$-stable and not simple. Since I is not simple, we can find, according to [Est01, Prop. 1], a proper subcurve $Y \subset X$ such that the natural map $I \to I_Y \oplus I_{Y^c}$ is an isomorphism. We claim that this implies that

$$\deg I = \deg_Y(I) + \deg_{Y^c}(I) + \delta_Y. \tag{2.11}$$

Indeed, since $I \cong I_Y \oplus I_{Y^c}$, we have that $\chi(I) = \chi(I_Y) + \chi(I_{Y^c})$. Moreover, from the exact sequence

$$0 \to \mathcal{O}_X \to \mathcal{O}_Y \oplus \mathcal{O}_{Y^c} \to \mathcal{O}_{Y \cap Y^c} \to 0,$$

we deduce that $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_Y) + \chi(\mathcal{O}_{Y^c}) - \chi(\mathcal{O}_{Y\cap Y^c}) = \chi(\mathcal{O}_Y) + \chi(\mathcal{O}_{Y^c}) - \delta_Y$. Using now the definition of degree, we compute

$$\begin{aligned}\deg(I) = \chi(I) - \chi(\mathcal{O}_X) = \chi(I_Y) + \chi(I_{Y^c}) - \chi(\mathcal{O}_Y) - \chi(\mathcal{O}_{Y^c}) + \delta_Y = \\ = \deg(I_Y) + \deg(I_{Y^c}) + \delta_Y,\end{aligned}$$

which gives (2.11).

Since I is $\underline{q}$-semistable, we get from (2.10), the two inequalities

$$\begin{cases} \deg_Y(I) \geq \underline{q}_Y - \dfrac{\delta_Y}{2}, \\ \deg_{Y^c}(I) \geq \underline{q}_{Y^c} - \dfrac{\delta_{Y^c}}{2}. \end{cases} \tag{2.12}$$

Summing up these two inequalities and using that $\deg(I) = |\underline{q}| = \underline{q}_Y + \underline{q}_{Y^c}$ and that $\delta_Y = \delta_{Y^c}$, we get

$$\deg_Y(I) + \deg_{Y^c}(I) \geq \deg(I) - \delta_Y.$$

Equation (2.11) tells us that the above inequality is actually an equality; hence equalities must hold in each of the inequalities in (2.12) and this contradicts the fact that I is $\underline{q}$-stable.

Let us now prove part (ii).

(a) ⇒ (b): Since $\underline{q}$ is general, from Remark 2.7 it follows that if $Y \subset X$ is a subcurve of X such that Y and Y^c are connected then $\underline{q}_Y - \dfrac{\delta_Y}{2} \notin \mathbb{Z}$. Therefore, the right hand side of (2.10) is not an integer for such subcurves, hence the inequality is a fortiori always strict. This is enough to guarantee that a torsion-free rank-1 sheaf that is $\underline{q}$-semistable is also $\underline{q}$-stable, by Remark 2.9.

(b) ⇒ (c) ⇒ (d) are trivial.

(d) ⇒ (a): the proof of [MV12, Lemma 6.2] for nodal curves extends easily to any curve X. ☐

Given a polarization $\underline{q}$ on X, we denote by $\overline{J}_X^{ss}(\underline{q})$ (resp. $\overline{J}_X^{s}(\underline{q})$) the subscheme of $\overline{\mathbb{J}}_X$ parametrizing simple rank-1 torsion-free sheaves I on X which are $\underline{q}$-semistable (resp. $\underline{q}$-stable). By [Est01, Prop. 34], the inclusions

$$\overline{J}_X^{s}(\underline{q}) \subseteq \overline{J}_X^{ss}(\underline{q}) \subset \overline{\mathbb{J}}_X$$

are open.

Fact 2.11 (Esteves). *Let X be a connected curve.*

- *(i) $\overline{J}_X^{s}(\underline{q})$ is a quasi-projective scheme over k (not necessarily reduced). In particular, $\overline{J}_X^{s}(\underline{q})$ is a scheme of finite type and separated over k.*
- *(ii) $\overline{J}_X^{ss}(\underline{q})$ is a k-scheme of finite type and universally closed over k.*
- *(iii) If $\underline{q}$ is general then $\overline{J}_X^{ss}(\underline{q}) = \overline{J}_X^{s}(\underline{q})$ is a projective scheme over k (not necessarily reduced).*
- *(iv) $\overline{\mathbb{J}}_X = \bigcup\limits_{\underline{q} \text{ general}} \overline{J}_X^{s}(\underline{q})$.*

Proof. Part (i) follows from [Est01, Thm. A(2) and Thm. C(4)].

Part (ii) follows from [Est01, Thm. A(1)].

Part (iii): the fact that $\overline{J}_X^{ss}(\underline{q}) = \overline{J}_X^{s}(\underline{q})$ follows from Lemma 2.10. Its projectivity follows from (i) and (ii) since a quasi-projective scheme over k which is universally closed over k must be a projective scheme over k.

Part (iv) follows from [Est01, Prop. 13] together with [Est09, Rmk. 2.5].

☐

If $\underline{q}$ is general, we set $\overline{J}_X(\underline{q}) := \overline{J}_X^{ss}(\underline{q}) = \overline{J}_X^{s}(\underline{q})$ and we call it the *fine compactified Jacobian* with respect to the polarization $\underline{q}$. We denote by $J_X(\underline{q})$ the open subset of $\overline{J}_X(\underline{q})$ parametrizing line bundles on X. Note that $J_X(\underline{q})$ is isomorphic to the disjoint union of a certain number of copies of the generalized Jacobian $J(X) = \mathrm{Pic}^{\underline{0}}(X)$ of X.

Since, for $\underline{q}$ general, $J_X(\underline{q})$ is an open subset of $\overline{\mathbb{J}}_X$, the above Theorem 2.3 immediately yields the following properties for the fine compactified Jacobians of curves with locally planar singularities.

Corollary 2.12. *Let X be a connected curve with locally planar singularities and $\underline{q}$ a general polarization on X. Then*

- *(i) $\overline{J}_X(\underline{q})$ is a reduced scheme with locally complete intersection singularities.*
- *(ii) $J_X(\underline{q})$ is dense in $\overline{J}_X(\underline{q})$. In particular, $\overline{J}_X(\underline{q})$ has pure dimension equal to the arithmetic genus $p_a(X)$ of X.*
- *(iii) $J_X(\underline{q})$ is the smooth locus of $\overline{J}_X(\underline{q})$.*

Later, we will prove that $\overline{J}_X(\underline{q})$ is connected (see Corollary 4.4) and we will give a formula for the number of its irreducible components in terms solely of the combinatorics of the curve X (see Corollary 4.12).

Fine compactified Jacobians of X depend on the choice of a polarization and there are clearly infinitely many polarizations. However, we are now going to show that there are finitely many isomorphism classes of fine compactified Jacobians. The simplest way to show that two fine compactified Jacobians are isomorphic is to show that there is translation that sends one into the other.

Definition 2.13. Let X be a connected curve. We say that two compactified Jacobians $\overline{J}_X(\underline{q})$ and $\overline{J}_X(\underline{q}')$ are *equivalent by translation* if there exists a line bundle L on X inducing an isomorphism

$$\begin{aligned} \overline{J}_X(\underline{q}) &\xrightarrow{\cong} \overline{J}_X(\underline{q}'), \\ I &\mapsto I \otimes L. \end{aligned}$$

Note however that, in general, there could be fine compactified Jacobians that are isomorphic without being equivalent by translation, see Section 7 for some explicit examples.

Lemma 2.14. *Let X be a connected curve. There is a finite number of fine compactified Jacobians up to equivalence by translation.*

Proof. If two generic polarizations $\underline{q}$ and $\underline{q}'$ are such that $\underline{q}-\underline{q}'\in\mathbb{Z}^{\gamma(X)}$, then the multiplication by a line bundle of multidegree $\underline{q}-\underline{q}'$ gives an isomorphism between $\overline{J}_X(\underline{q}')$ and $\overline{J}_X(\underline{q})$. Therefore, any fine compactified Jacobian of X is equivalent by translation to a fine compactified Jacobian $\overline{J}_X(\underline{q})$ such that $0\le \underline{q}_{C_i}<1$ for any irreducible component C_i of X.

Moreover, if $\underline{q}$ and $\underline{q}'$ are two generic polarizations on X such that $\lceil \underline{q}_Y-\frac{\delta_Y}{2}\rceil=\lceil \underline{q}'_Y-\frac{\delta_Y}{2}\rceil$ for any subcurve $Y\subseteq X$, then a sheaf $I\in\overline{\mathbb{J}}_X$ is $\underline{q}$-stable if and only if it is $\underline{q}'$-stable. Therefore, the arrangement of hyperplanes in $\mathbb{R}^{\gamma(X)}$

$$\left\{\sum_{C_i\subseteq Y}x_i-\frac{\delta_Y}{2}=n\right\}_{Y\subseteq X,n\in\mathbb{Z}}$$

subdivides the unitary cube $[0,1)^{\gamma(X)}\subset\mathbb{R}^{\gamma(X)}$ into finitely many chambers such that if $\underline{q}$ and $\underline{q}'$ belong to the same chamber then $\overline{J}_X(\underline{q})=\overline{J}_X(\underline{q}')$. The assertion now follows. ☐

3. DEFORMATION THEORY

The aim of this section is to study the deformation theory and the semiuniversal deformation space of a pair (X,I) where X is a (reduced) connected curve and I is rank-1 torsion-free simple sheaf on X. For basic facts on deformation theory, we refer to the book of Sernesi [Ser06].

3.1. **Deformation theory of X.**

The aim of this subsection is to recall some well-known facts about the deformation theory of a (reduced) curve X.

Let $\mathrm{Def_X}$ (resp. $\mathrm{Def'_X}$) be the local moduli functor of X (resp. the locally trivial moduli functor) of X in the sense of [Ser06, Sec. 2.4.1]. Moreover, for any $p\in X_{\rm sing}$, we denote by $\mathrm{Def_{X,p}}$ the deformation functor of the complete local k-algebra $\mathcal{O}_{X,p}$ in the sense of [Ser06, Sec. 1.2.2]. The above deformation functors are related by the following natural morphisms:

$$\mathrm{Def'_X}\to\mathrm{Def_X}\to\mathrm{Def_X^{loc}}:=\prod_{\mathrm{p\in X_{sing}}}\mathrm{Def_{X,p}}. \tag{3.1}$$

Since X is reduced, the tangent spaces to $\mathrm{Def'_X}$, $\mathrm{Def_X}$ and $\mathrm{Def_{X,p}}$ where $p\in X_{\rm sing}$ are isomorphic to (see [Ser06, Cor. 1.1.11, Thm. 2.4.1])

$$\begin{aligned} T\mathrm{Def'_X}&:=\mathrm{Def'_X(k[\epsilon])=H^1(X,T_X)},\\ T\mathrm{Def_X}&:=\mathrm{Def_X(k[\epsilon])=Ext^1(\Omega^1_X,\mathcal{O}_X)},\\ T\mathrm{Def_{(X,p)}}&:=\mathrm{Def_{(X,p)}(k[\epsilon])=(T^1_X)_p},\end{aligned} \tag{3.2}$$

where Ω^1_X is the sheaf of Kähler differentials on X, $T_X:=\mathcal{H}om(\Omega^1_X,\mathcal{O}_X)$ is the tangent sheaf of X and $T^1_X=\mathcal{E}xt^1(\Omega^1_X,\mathcal{O}_X)$ is the first cotangent sheaf of X, which is a sheaf supported on $X_{\rm sing}$ by [Ser06, Prop. 1.1.9(ii)].

The usual local-to-global spectral sequence gives a short exact sequence

$$\begin{aligned} &0\to H^1(X,T_X)=T\mathrm{Def'_X}\to\mathrm{Ext^1(\Omega^1_X,\mathcal{O}_X)=TDef_X}\to\\ &\to H^0(X,\mathcal{E}xt^1(\Omega^1_X,\mathcal{O}_X))=\bigoplus_{p\in X_{\rm sing}}\mathcal{E}xt^1(\Omega^1_X,\mathcal{O}_X)_p=T\mathrm{Def_X^{loc}}\to\mathrm{H^2(X,T_X)}=0,\end{aligned} \tag{3.3}$$

which coincides with the exact sequence on the tangent spaces induced by (3.1).

By looking at the obstruction spaces of the above functors, one can give criteria under which the above deformation functors are smooth (in the sense of [Ser06, Def. 2.2.4]).

Fact 3.1.

(i) $\mathrm{Def'_X}$ *is smooth;*
(ii) If X *has l.c.i. singularities at* $p\in X_{\rm sing}$ *then* $\mathrm{Def_{X,p}}$ *is smooth;*
(iii) If X *has l.c.i. singularities, then* $\mathrm{Def_X}$ *is smooth and the morphism* $\mathrm{Def_X}\to\mathrm{Def_X^{loc}}$ *is smooth.*

Proof. Part (i): an obstruction space for $\mathrm{Def'_X}$ is $H^2(X,T_X)$ by [Ser06, Prop. 2.4.6] and $H^2(X,T_X)=0$ because $\dim X=1$. Therefore, $\mathrm{Def'_X}$ is smooth.

Part (ii) follows from [Ser06, Cor. 3.1.13(ii)].

Part (iii): by [Ser06, Prop. 2.4.8], an obstruction space for $\mathrm{Def_X}$ is $\mathrm{Ext^2(\Omega^1_X,\mathcal{O}_X)}$, which is zero by [Ser06, Example 2.4.9]. Therefore we get that $\mathrm{Def_X}$ is smooth.

Since $\mathrm{Def}_{\mathrm{X}}^{\mathrm{loc}}$ is smooth by part (ii) and the map of tangent spaces $T\mathrm{Def}_{\mathrm{X}} \to \mathrm{TDef}_{\mathrm{X}}^{\mathrm{loc}}$ is surjective by (3.3), the smoothness of the morphism $\mathrm{Def}_{\mathrm{X}} \to \mathrm{Def}_{\mathrm{X}}^{\mathrm{loc}}$ follows from the criterion [Ser06, Prop. 2.3.6]. □

3.2. **Deformation theory of the pair** (X, I)**.** The aim of this subsection is to review some fundamental results due to Fantechi-Göttsche-vanStraten [FGvS99] on the deformation theory of a pair (X, I), where X is a (reduced) curve and I is a rank-1 torsion-free sheaf on X (not necessarily simple).

Let $\mathrm{Def}_{(\mathrm{X},\mathrm{I})}$ be the deformation functor of the pair (X, I) and, for any $p \in X_{\mathrm{sing}}$, we denote by $\mathrm{Def}_{(\mathrm{X},\mathrm{I}),\mathrm{p}}$ the deformation functor of the pair $(\mathcal{O}_{X,p}, I_p)$. We have a natural commutative diagram

$$\begin{tikzcd}
\mathrm{Def}_{(\mathrm{X},\mathrm{I})} \arrow[r] \arrow[d] & \mathrm{Def}_{(\mathrm{X},\mathrm{I})}^{\mathrm{loc}} := \prod_{\mathrm{p}\in\mathrm{X}_{\mathrm{sing}}} \mathrm{Def}_{(\mathrm{X},\mathrm{I}),\mathrm{p}} \arrow[d] \\
\mathrm{Def}_{\mathrm{X}} \arrow[r] & \mathrm{Def}_{\mathrm{X}}^{\mathrm{loc}} := \prod_{\mathrm{p}\in\mathrm{X}_{\mathrm{sing}}} \mathrm{Def}_{\mathrm{X},\mathrm{p}}.
\end{tikzcd} \tag{3.4}$$

Under suitable hypothesis, the deformation functors appearing in the above diagram (3.4) are smooth and the horizontal morphisms are smooth as well.

Fact 3.2 (Fantechi-Göttsche-vanStraten)**.**

(i) The natural morphism

$$\mathrm{Def}_{(\mathrm{X},\mathrm{I})} \to \mathrm{Def}_{(\mathrm{X},\mathrm{I})}^{\mathrm{loc}} \times_{\mathrm{Def}_{\mathrm{X}}^{\mathrm{loc}}} \mathrm{Def}_{\mathrm{X}}$$

is smooth. In particular, if X has l.c.i. singularities then the morphism $\mathrm{Def}_{(\mathrm{X},\mathrm{I})} \to \mathrm{Def}_{(\mathrm{X},\mathrm{I})}^{\mathrm{loc}}$ *is smooth.*

(ii) If X has locally planar singularities at $p \in X_{\mathrm{sing}}$ then $\mathrm{Def}_{(\mathrm{X},\mathrm{I}),\mathrm{p}}$ *is smooth. In particular, if X has locally planar singularities then* $\mathrm{Def}_{(\mathrm{X},\mathrm{I})}^{\mathrm{loc}}$ *and* $\mathrm{Def}_{(\mathrm{X},\mathrm{I})}$ *are smooth.*

Proof. Part (i): the first assertion follows from [FGvS99, Prop. A.1]. The second assertion follows from the first one together with Fact 3.1(iii) which implies that the morphism $\mathrm{Def}_{(\mathrm{X},\mathrm{I})}^{\mathrm{loc}} \times_{\mathrm{Def}_{\mathrm{X}}^{\mathrm{loc}}} \mathrm{Def}_{\mathrm{X}} \to \mathrm{Def}_{(\mathrm{X},\mathrm{I})}^{\mathrm{loc}}$ is smooth.

Part (ii): the first assertion follows from [FGvS99, Prop. A.3]. The second assertion follows from the first together with part (i). □

3.3. **Semiuniversal deformation space.** The aim of this subsection is to describe and study the semiuniversal deformation spaces for the deformation functors $\mathrm{Def}_{\mathrm{X}}$ and $\mathrm{Def}_{(\mathrm{X},\mathrm{I})}$.

According to [Ser06, Cor. 2.4.2], the functor $\mathrm{Def}_{\mathrm{X}}$ admits a semiuniversal [4] formal couple $(R_X, \overline{\mathcal{X}})$, where R_X is a Noetherian complete local k-algebra with maximal ideal $\mathfrak{m}_X$ and residue field k and

$$\overline{\mathcal{X}} \in \widehat{\mathrm{Def}_{\mathrm{X}}}(R_X) := \varprojlim \mathrm{Def}_{\mathrm{X}}\left(\frac{\mathrm{R}_{\mathrm{X}}}{\mathfrak{m}_{\mathrm{X}}^{\mathrm{n}}}\right)$$

is a formal deformation of X over R_X. Recall that this means that the morphism of functors

$$h_{R_X} := \mathrm{Hom}(\mathrm{R}_{\mathrm{X}}, -) \longrightarrow \mathrm{Def}_{\mathrm{X}} \tag{3.5}$$

determined by $\overline{\mathcal{X}}$ is smooth and induces an isomorphism of tangent spaces $TR_X := (\mathfrak{m}_X/\mathfrak{m}_X^2)^\vee \xrightarrow{\cong} T\mathrm{Def}_{\mathrm{X}}$ (see [Ser06, Sec. 2.2]). The formal couple $(R_X, \overline{\mathcal{X}})$ can be also viewed as a flat morphism of formal schemes

$$\overline{\pi} : \overline{\mathcal{X}} \to \mathrm{Spf}\ \mathrm{R}_{\mathrm{X}}, \tag{3.6}$$

where Spf denotes the formal spectrum, such that the fiber over $[\mathfrak{m}_X] \in \mathrm{Spf}\,\mathrm{R}_{\mathrm{X}}$ is isomorphic to X (see [Ser06, p. 77]). Note that the semiuniversal formal couple $(R_X, \overline{\mathcal{X}})$ is unique by [Ser06, Prop. 2.2.7].

Since X is projective and $H^2(X, \mathcal{O}_X) = 0$, Grothendieck's existence theorem (see [Ser06, Thm. 2.5.13]) gives that the formal deformation (3.6) is *effective*, i.e. there exists a deformation $\pi : \mathcal{X} \to \mathrm{Spec}\,\mathrm{R}_{\mathrm{X}}$ of

---

[4] Some authors use the word miniversal instead of semiuniversal. We prefer to use the word semiuniversal in order to be coherent with the terminology of the book of Sernesi [Ser06].

X over $\mathrm{Spec}\,\mathrm{R_X}$ whose completion along $X = \pi^{-1}([\mathfrak{m}_X])$ is isomorphic to (3.6). In other words, we have a Cartesian diagram

$$\begin{tikzcd}
X \arrow[r, hook] \arrow[d] \arrow[dr, phantom, "\square"] & \overline{\mathcal{X}} \arrow[r] \arrow[d, "\overline{\pi}"] \arrow[dr, phantom, "\square"] & \mathcal{X} \arrow[d, "\pi"] \\
\mathrm{Spec}\,\mathrm{k} \cong [\mathfrak{m}_\mathrm{X}] \arrow[r, hook] & \mathrm{Spf}\,\mathrm{R_X} \arrow[r] & \mathrm{Spec}\,\mathrm{R_X}.
\end{tikzcd} \tag{3.7}$$

Note also that the deformation π is unique by [Ser06, Thm. 2.5.11].

The space $\mathrm{Spec}\,\mathrm{R_X}$ admits two stratifications into closed subsets according to either the arithmetic genus or the geometric genus of the normalization of the fibers of the family π. More precisely, using the notation introduced in 1.2, we have two functions

$$\begin{aligned} p_a^\nu : \mathrm{Spec}\,\mathrm{R_X} &\longrightarrow \mathbb{N}, & g^\nu : \mathrm{Spec}\,\mathrm{R_X} &\longrightarrow \mathbb{N}, \\ s &\mapsto p_a^\nu(\mathcal{X}_s) := p_a(\mathcal{X}_s^\nu), & s &\mapsto g^\nu(\mathcal{X}_s) = g^\nu(\mathcal{X}_s^\nu), \end{aligned} \tag{3.8}$$

where $\mathcal{X}_s$ denotes the fiber of π over the point $s \in \mathrm{Spec}\,\mathrm{R_X}$. Since the number of connected components of $\mathcal{X}_s^\nu$ is the number $\gamma(\mathcal{X}_s)$ of irreducible components of $\mathcal{X}_s$, we have the relation

$$p_a^\nu(\mathcal{X}_s) = g^\nu(\mathcal{X}_s) - \gamma(\mathcal{X}_s) + 1 \leq g^\nu(\mathcal{X}_s). \tag{3.9}$$

The functions p_a^ν and g^ν are lower semi-continuous since the arithmetic genus and the geometric genus of the normalization cannot increase by specialization. Using this lower-semicontinuity, (3.9) and the fact that the arithmetic genus p_a stays constant in the family π because of flatness, we get that

$$p_a(X^\nu) = p_a^\nu(X) \leq p_a^\nu(\mathcal{X}_s) \leq g^\nu(\mathcal{X}_s) \leq p_a(\mathcal{X}_s) = p_a(X).$$

Therefore for any $p_a(X^\nu) \leq k \leq p_a(X)$ we have two closed subsets of $\mathrm{Spec}\,\mathrm{R_X}$:

$$(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^\nu \leq \mathrm{k}} := \{\mathrm{s} \in \mathrm{Spec}\,\mathrm{R_X} \ : \ \mathrm{g}^\nu(\mathcal{X}_\mathrm{s}) \leq \mathrm{k}\} \subseteq (\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{k}} := \{\mathrm{s} \in \mathrm{Spec}\,\mathrm{R_X} \ : \ \mathrm{p}_\mathrm{a}^\nu(\mathcal{X}_\mathrm{s}) \leq \mathrm{k}\}. \tag{3.10}$$

If X has locally planar singularities, then the second closed subset behaves particularly well due to the following results of Teissier ([Tei80]) and Diaz-Harris ([DH88]).

Fact 3.3 (Teissier, Diaz-Harris)**.** *Assume that X has locally planar singularities. Then, for any $p_a(X^\nu) \leq k \leq p_a(X)$, we have that:*

(i) The codimension of the closed subset $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{k}}$ inside $\mathrm{Spec}\,\mathrm{R_X}$ is equal to $p_a(X) - k$.
(ii) Each generic point s of $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{k}}$ is such that $\mathcal{X}_s$ is a nodal curve with $p_a(X) - k$ nodes.

Proof. Let us first prove the two statements for $k = p_a(X^\nu)$. The arithmetic genus of the curve $\mathcal{X}_s$ and the one of its normalization $\mathcal{X}_s^\nu$ are related by the well-known formula

$$p_a(\mathcal{X}_s) = p_a(\mathcal{X}_s^\nu) + \delta(\mathcal{X}_s), \tag{3.11}$$

where

$$\delta(\mathcal{X}_s) := \sum_{q \in (\mathcal{X}_s)_{\mathrm{sing}}} \delta(X, q) = \sum_{q \in (\mathcal{X}_s)_{\mathrm{sing}}} \mathrm{length}(\mathcal{O}_{\mathcal{X}_s, q}^\nu / \mathcal{O}_{\mathcal{X}_s, q})$$

is the total delta invariant of $\mathcal{X}_s$. Using the equation (3.11) and the fact that $p_a(\mathcal{X}_s) = p_a(X)$ for every $s \in \mathrm{Spec}\,\mathrm{R_X}$, we deduce that

$$p_a(\mathcal{X}_s^\nu) = p_a(X^\nu) \Leftrightarrow \delta(\mathcal{X}_s) = \delta(X).$$

In oder words, $(\mathrm{Spec}\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{p}_\mathrm{a}(\mathrm{X}^\nu)}$ is the closed subset of $\mathrm{Spec}\mathrm{R_X}$ where the delta invariant stays constant. Since the total δ-invariant is the sum of the local δ-invariants, we conclude that $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{p}_\mathrm{a}(\mathrm{X}^\nu)}$ is the inverse image via the natural morphism

$$F : h_{R_X} \to \mathrm{Def_X} \to \mathrm{Def}_\mathrm{X}^{\mathrm{loc}} = \prod_{\mathrm{p} \in \mathrm{X}_{\mathrm{sing}}} \mathrm{Def}_{\mathrm{X,p}}$$

of the product of the δ-constant (or equigeneric) loci EG_p inside each $\mathrm{Def}_{\mathrm{X,p}}$. For any $p \in X_{\mathrm{sing}}$, the δ-constant locus EG_p has codimension $\delta(X, p)$ inside $\mathrm{Def}_{\mathrm{X,p}}$ by [DH88, p. 435] and, moreover, each generic point of EG_p corresponds to a deformation of the singularity $\widehat{O}_{X,p}$ having $\delta(X, p)$ nodes (and no other singularities) by [Tei80, p. 10]. Since the morphism F is smooth by Fact 3.1(iii), we conclude that

$$\mathrm{codim}(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{p}_\mathrm{a}(\mathrm{X}^\nu)} = \sum_{\mathrm{p} \in \mathrm{X}_{\mathrm{sing}}} \mathrm{codim}(\mathrm{EG_p}) = \sum_{\mathrm{p} \in \mathrm{X}_{\mathrm{sing}}} \delta(\mathrm{X}, \mathrm{p}) = \delta(\mathrm{X}) = \mathrm{p_a}(\mathrm{X}) - \mathrm{p_a}(\mathrm{X}^\nu),$$

and, moreover, that the fiber of $\mathcal{X}$ over every generic point of $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{p}_\mathrm{a}^\nu \leq \mathrm{p}_\mathrm{a}(\mathrm{X}^\nu)}$ is a nodal curve.

The case of an arbitrary k follows from the previous case using the openess of versality, which amounts to say that for any $s \in \operatorname{Spec} \mathrm{R_X}$ a versal deformation ring for $\mathcal{X}_s$ is provided by the completion $\widehat{(R_X)_s}$ of the localization of R_X at s together with the image of the family $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R_X}$. $\square$

From the above Fact 3.3 together with the inclusion in (3.10), we get the following

Corollary 3.4. *Assume that X has locally planar singularities. Then, for any $p_a(X^\nu) \leq k \leq p_a(X)$, the codimension of the closed subset* $(\operatorname{Spec} \mathrm{R_X})^{\mathrm{g}^\nu \leq \mathrm{k}}$ *inside* $\operatorname{Spec} \mathrm{R_X}$ *is at least equal to $p_a(X) - k$.*

Another useful result that we will need in the sequel is contained in the following

Lemma 3.5. *Assume that X has locally planar singularities. Let U be the open subset of* $\operatorname{Spec} \mathrm{R_X}$ *consisting of all the (schematic) points $s \in$* $\operatorname{Spec} \mathrm{R_X}$ *such that the fiber $\mathcal{X}_s$ of the universal family $\pi : \mathcal{X} \to$* $\operatorname{Spec} \mathrm{R_X}$ *is smooth or has a unique singular point which is a node. Then the codimension of the complement of U inside* $\operatorname{Spec} \mathrm{R_X}$ *is at least two.*

Proof. Since the natural morphism (see (3.1))

$$\mathrm{Def_X} \to \mathrm{Def_X^{loc}} := \prod_{\mathrm{p} \in \mathrm{X_{sing}}} \mathrm{Def_{X,p}}$$

is smooth by Fact 3.1(iii), it is enough to show that if $\mathrm{Def_{X,p}}$ has dimension at most one then $p \in X_{\mathrm{sing}}$ is a node of X. Since X has locally planar singularities, this is exactly the content of [Ser06, Prop. 3.1.5.]. $\square$

Consider now the functor

$$\overline{\mathbb{J}}_{\mathcal{X}}^* : \{\operatorname{Spec} \mathrm{R_X} - \text{schemes}\} \longrightarrow \{\text{Sets}\}$$

which sends a scheme $T \to \operatorname{Spec} \mathrm{R_X}$ to the set of isomorphism classes of T-flat, coherent sheaves on $\mathcal{X}_T := T \times_{\operatorname{Spec} \mathrm{R_X}} \mathcal{X}$ whose fibers over T are simple rank-1 torsion-free sheaves. The functor $\overline{\mathbb{J}}_{\mathcal{X}}^*$ contains the open subfunctor

$$\mathbb{J}_{\mathcal{X}}^* : \{\operatorname{Spec} \mathrm{R_X} - \text{schemes}\} \longrightarrow \{\text{Sets}\}$$

which sends a scheme $T \to \operatorname{Spec} \mathrm{R_X}$ to the set of isomorphism classes of line bundles on $\mathcal{X}_T$.

Analogously to Fact 2.2, we have the following

Fact 3.6 (Altman-Kleiman, Esteves)**.**

(i) The étale sheafification of $\overline{\mathbb{J}}_{\mathcal{X}}^*$ is represented by a scheme $\overline{\mathbb{J}}_{\mathcal{X}}$ endowed with a morphism $u : \overline{\mathbb{J}}_{\mathcal{X}} \to$ $\operatorname{Spec} \mathrm{R_X}$*, which is locally of finite type and universally closed. The scheme $\overline{\mathbb{J}}_{\mathcal{X}}$ contains an open subset $\mathbb{J}_{\mathcal{X}}$ which represents the étale sheafification of $\mathbb{J}_{\mathcal{X}}^*$ and the restriction $u : \mathbb{J}_{\mathcal{X}} \to$* $\operatorname{Spec} \mathrm{R_X}$ *is smooth.*
Moreover, the fiber of $\overline{\mathbb{J}}_{\mathcal{X}}$ (resp. of $\mathbb{J}_{\mathcal{X}}$) over the closed point $[\mathfrak{m}_X] \in$ $\operatorname{Spec} \mathrm{R_X}$ *is isomorphic to $\overline{\mathbb{J}}_X$ (resp. $\mathbb{J}_X$).*

(ii) There exists a sheaf $\widehat{\mathcal{I}}$ on $\mathcal{X} \times_{\operatorname{Spec} \mathrm{R_X}} \overline{\mathbb{J}}_{\mathcal{X}}$ such for every $\mathcal{F} \in \overline{\mathbb{J}}_{\mathcal{X}}^*(T)$ there exists a unique $\operatorname{Spec} \mathrm{R_X}$*-map $\alpha_{\mathcal{F}} : T \to \overline{\mathbb{J}}_{\mathcal{X}}$ with the property that $\mathcal{F} = (\mathrm{id}_{\mathcal{X}} \times \alpha_{\mathcal{F}})^*(\widehat{\mathcal{I}}) \otimes \pi_2^*(\mathrm{N})$ for some $N \in \operatorname{Pic}(T)$, where $\pi_2 : \mathcal{X} \times_{\operatorname{Spec} \mathrm{R_X}} T \to T$ is the projection onto the second factor. The sheaf $\widehat{\mathcal{I}}$ is uniquely determined up to tensor product with the pullback of an invertible sheaf on $\overline{\mathbb{J}}_{\mathcal{X}}$ and it is called a* universal sheaf *on $\overline{\mathbb{J}}_{\mathcal{X}}$.*
Moreover, the restriction of $\widehat{\mathcal{I}}$ to $X \times \overline{\mathbb{J}}_X$ is equal to a universal sheaf as in Fact 2.2(iii).

Proof. Part (i): the representability of the étale sheafification of $\overline{\mathbb{J}}_{\mathcal{X}}^*$ by an algebraic space $\overline{\mathbb{J}}_{\mathcal{X}}$ locally of finite type over $\operatorname{Spec} \mathrm{R_X}$ follows from [AK80, Thm. 7.4]. From [Est01, Cor. 52], it follows that $\overline{\mathbb{J}}_{\mathcal{X}}$ becomes a scheme after an étale cover of $\operatorname{Spec} \mathrm{R_X}$. However, since R_X is strictly henselian (being a complete local ring with algebraically closed residue field), then $\operatorname{Spec} \mathrm{R_X}$ does not admit non trivial étale covers (see [BLR90, Sec. 2.3]); hence $\overline{\mathbb{J}}_{\mathcal{X}}$ is a scheme. The fact that $\overline{\mathbb{J}}_{\mathcal{X}}$ is universally closed follows from [Est01, Thm. 32]. Since $\mathbb{J}_{\mathcal{X}}^*$ is an open subfunctor of $\overline{\mathbb{J}}_{\mathcal{X}}^*$, it follows that $\overline{\mathbb{J}}_{\mathcal{X}}$ contains an open subscheme $\mathbb{J}_{\mathcal{X}}$ which represents the étale sheafification of $\mathbb{J}_{\mathcal{X}}^*$. The smoothness of $\mathbb{J}_{\mathcal{X}}$ over $\operatorname{Spec} \mathrm{R_X}$ follows from [BLR90, Prop. 2]. The last assertion of part (i) is obvious.

Part (ii) follows from [AK79, Thm. 3.4] once we prove that the morphism $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R_X}$ admits a section through its smooth locus. Indeed, let U be the smooth locus of the morphism π and denote by $\pi' : U \to \operatorname{Spec} \mathrm{R_X}$ the restriction of π to U. Since X is assumed to be reduced and all the fibers of π are smoothings of X and hence reduced as well, we deduce that for every $s \in \operatorname{Spec} \mathrm{R_X}$ the open subset $\pi'^{-1}(s)$ is dense in $\pi^{-1}(s)$. Now, since R_X is a strictly henselian ring, given any point $p \in \pi'^{-1}([\mathfrak{m}_X])$, we can find a section of $\pi' : U \to \mathrm{Spec\, R_X}$ passing through p (see [BLR90, Sec. 2.2, Prop. 14]), q.e.d. The last assertion of part (ii) is obvious.

$\square$

Let now I be a simple rank-1 torsion-free sheaf I on X, i.e. $I \in \overline{\mathbb{J}}_X \subset \overline{\mathbb{J}}_{\mathcal{X}}$. If we denote by $R_{(X,I)} := \widehat{\mathcal{O}}_{\overline{\mathbb{J}}_{\mathcal{X}},I}$ the completion of the local ring of $\overline{\mathbb{J}}_{\mathcal{X}}$ at I and by $\mathfrak{m}_{(X,I)}$ its maximal ideal, then there is a natural map $f : \mathrm{Spec\, R_{(X,I)}} \to \overline{\mathbb{J}}_{\mathcal{X}}$ which fits into the following Cartesian diagram

$$\begin{tikzcd}
\mathrm{(id \times f)^*}(\widehat{\mathcal{I}}) \arrow[d, dash] & \widehat{\mathcal{I}} \arrow[d, dash] & \\
\mathcal{X} \times_{\mathrm{Spec\, R_X}} \mathrm{Spec\, R_{(X,I)}} \arrow[r, "\mathrm{id}\times f"] \arrow[d, "\pi\times\mathrm{id}"'] \arrow[dr, phantom, "\square"] & \mathcal{X} \times_{\mathrm{Spec\, R_X}} \overline{\mathbb{J}}_{\mathcal{X}} \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & \mathcal{X} \arrow[d, "\pi"] \\
\mathrm{Spec\, R_{(X,I)}} \arrow[r, "j"] & \overline{\mathbb{J}}_{\mathcal{X}} \arrow[r, "u"] & \mathrm{Spec\, R_X}.
\end{tikzcd} \tag{3.12}$$

Since $I \in \overline{\mathbb{J}}_X \subset \overline{\mathbb{J}}_{\mathcal{X}}$, the map $u \circ j$ sends the closed point $[\mathfrak{m}_{X,I}] \in \mathrm{Spec}\, \widehat{\mathcal{O}}_{\overline{\mathbb{J}}_{\mathcal{X}},\mathrm{I}}$ into the closed point $[\mathfrak{m}_X] \in \mathrm{Spec\, R_X}$. In particular, we have that $(\pi \times \mathrm{id})^{-1}(\mathfrak{m}_{\mathrm{(X,I)}}) = \pi^{-1}(\mathfrak{m}_{\mathrm{X}}) = \mathrm{X}$ and the restriction of $\mathrm{(id \times j)^*}(\widehat{\mathcal{I}})$ to $(\pi \times \mathrm{id})^{-1}(\mathfrak{m}_{\mathrm{(X,I)}}) = \mathrm{X}$ is isomorphic to I by the universal property in Fact 3.6(ii). The completion of the family $\pi \times \mathrm{id}$ along the central fiber X together with the completion of the sheaf $\mathrm{(id \times j)^*}(\widehat{\mathcal{I}})$ determine a formal deformation of (X, I) over $R_{(X,I)}$, or equivalently, a morphism of deformation functors

$$h_{R_{(X,I)}} := \mathrm{Hom(R_{(X,I)}}, -) \longrightarrow \mathrm{Def_{(X,I)}}. \tag{3.13}$$

We can now prove the main result of this section.

Theorem 3.7. *Let X be a (reduced) curve and I a rank-1 torsion-free simple sheaf on X.*

- *(i) There exists a Cartesian diagram of deformation functors*

$$\begin{tikzcd}
h_{R_{(X,I)}} \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & \mathrm{Def_{(X,I)}} \arrow[d] \\
h_{R_X} \arrow[r] & \mathrm{Def_X},
\end{tikzcd} \tag{3.14}$$

where the horizontal arrows realize $R_{(X,I)}$ and R_X as the semiuniversal deformation rings for $\mathrm{Def_{(X,I)}}$ and $\mathrm{Def_X}$, respectively.
- *(ii) If X has l.c.i. singularities then R_X is smooth (i.e. it is a power series ring over k).*
- *(iii) If X has locally planar singularities then $R_{(X,I)}$ is smooth. In particular, the scheme $\overline{\mathbb{J}}_{\mathcal{X}}$ is smooth.*

Proof. Part (i): the fact that the diagram (3.14) is commutative follows from the definition of the map (3.13) and the commutativity of the diagram (3.12).

Let us check that the above diagram (3.14) is Cartesian. Let A be an Artinian local k-algebra with maximal ideal $\mathfrak{m}_A$. Suppose that there exists a deformation $(\widetilde{X}, \widetilde{I}) \in \mathrm{Def_{(X,I)}(A)}$ of (X, I) over A and a homomorphism $\phi \in \mathrm{Hom(R_X, A) = h_{R_X}(A)}$ that have the same image in $\mathrm{Def_X(A)}$. We have to find a homomorphism $\eta \in \mathrm{Hom(R_{(X,I)}, A) = h_{R_{(X,I)}}(A)}$ that maps into $\phi \in h_{R_X}(A)$ and $(\widetilde{X}, \widetilde{I}) \in \mathrm{Def_{(X,I)}(A)}$ via the maps of diagram (3.14). The assumption that the elements $(\widetilde{X}, \widetilde{I}) \in \mathrm{Def_{(X,I)}(A)}$ and $\phi \in h_{R_X}(A)$ have the same image in $\mathrm{Def_X(A)}$ is equivalent to the fact that $\widetilde{X}$ is isomorphic to $\mathcal{X}_A := \mathcal{X} \times_{\mathrm{Spec\, R_X}} \mathrm{Spec\, A}$ with respect to the natural morphism $\mathrm{Spec\, A \to Spec\, R_X}$ induced by ϕ. Therefore the sheaf $\widetilde{I}$ can be seen as an element of $\overline{\mathbb{J}}^*_{\mathcal{X}}(\mathrm{Spec\ A})$. Fact 3.6(ii) gives a map $\alpha_{\widetilde{I}} : \mathrm{Spec\ A} \to \overline{\mathbb{J}}_{\mathcal{X}}$ such that $\widetilde{I} = (\mathrm{id}_{\mathcal{X}} \times \alpha_{\widetilde{\mathrm{I}}})^*(\widehat{\mathcal{I}})$, because $\mathrm{Pic(Spec\ A)} = 0$. Clearly the map $\alpha_{\widetilde{I}}$ sends $[\mathfrak{m}_A]$ into $I \in \overline{\mathbb{J}}_X \subset \overline{\mathbb{J}}_{\mathcal{X}}$ and therefore it factors through a map $\beta : \mathrm{Spec\ A \to Spec\, R_{(X,I)}}$ followed by the map j of (3.12). The morphism β determines the element $\eta \in \mathrm{Hom(R_{(X,I)}, A) = h_{R_{(X,I)}}(A)}$ we were looking for.

Finally, the bottom horizontal morphism realizes the ring R_X as the semiuniversal deformation ring for $\mathrm{Def_X}$ by the very definition of R_X. Since the diagram (3.14) is Cartesian, the same is true for the top horizontal arrow.

Part (ii): R_X is smooth since the morphism $h_{R_X} \to \mathrm{Def_X}$ is smooth and $\mathrm{Def_X}$ is smooth by Fact 3.1(iii).

Part (iii): $R_{(X,I)}$ is smooth since the morphism $h_{R_{(X,I)}} \to \mathrm{Def}_{(\mathrm{X},\mathrm{I})}$ is smooth and $\mathrm{Def}_{(\mathrm{X},\mathrm{I})}$ is smooth by Fact 3.2(ii). We deduce that the open subset U of smooth points of $\overline{\mathbb{J}}_{\mathcal{X}}$ contains the central fiber $u^{-1}([\mathfrak{m}_X]) = \overline{\mathbb{J}}_X$, which implies that $U = \overline{\mathbb{J}}_{\mathcal{X}}$ because $u^{-1}([\mathfrak{m}_X])$ contains all the closed points of $\overline{\mathbb{J}}_{\mathcal{X}}$; hence $\overline{\mathbb{J}}_{\mathcal{X}}$ is smooth.

□

4. The universal fine compactified Jacobians

The aim of this section is to introduce and study the universal fine compactified Jacobians relative to the semiuniversal deformation $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$ introduced in 3.3.

The universal fine compactified Jacobian will depend on a general polarization $\underline{q}$ on X as in Definition 2.6. Indeed, the polarization $\underline{q}$ induces a polarization on each fiber of the effective semiuniversal deformation family $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$, in the following way. For any (schematic) point $s \in \mathrm{Spec}\, \mathrm{R_X}$, denote by $\mathcal{X}_s := \pi^{-1}(s)$ the fiber of π over s. There is a natural map

$$\begin{aligned} \Sigma_s : \{\text{Subcurves of } \mathcal{X}_s\} &\longrightarrow \{\text{Subcurves of } X\} \\ \mathcal{X}_s \supseteq Z &\mapsto \overline{Z} \cap X \subseteq X, \end{aligned} \tag{4.1}$$

where $\overline{Z}$ denotes the Zariski closure of Z inside $\mathcal{X}$.

Lemma-Definition 4.1. *Let $s \in \mathrm{Spec}\, \mathrm{R_X}$ and let $\underline{q}$ be a polarization on X. The polarization $\underline{q}^s$ induced by $\underline{q}$ on the fiber $\mathcal{X}_s$ is defined by*

$$\underline{q}^s_Z := \underline{q}_{\Sigma_s(Z)} \in \mathbb{Q}$$

for every subcurve $Z \subseteq \mathcal{X}_s$. If $\underline{q}$ is general then $\underline{q}^s$ is general.

Proof. Let us first check that $\underline{q}^s$ is well-defined. i.e. that $|\underline{q}^s| \in \mathbb{Z}$ and that $(Z \subseteq \mathcal{X}_s) \mapsto \underline{q}^s_Z$ is additive (see the discussion after Definition 2.5). Clearly we have that $|\underline{q}^s| = \underline{q}^s_{\mathcal{X}_s} = \underline{q}_X = |\underline{q}| \in \mathbb{Z}$. Moreover, if Z_1, Z_2 are two subcurves of $\mathcal{X}_s$ without common irreducible components then the subcurves $\Sigma_s(Z_1)$ and $\Sigma_s(Z_2)$ of X are without common irreducible components and therefore the additivity of $\underline{q}^s$ follows from the additivity of $\underline{q}$.

The last assertion is implied, using Remark 2.7, by the following easily checked properties of the map Σ_s:

$$\begin{cases} Z \text{ connected} \Rightarrow \Sigma_s(Z) \text{ connected}, \\ \Sigma_s(Z^c) = \Sigma_s(Z)^c, \\ \delta_Z \equiv \deg_Z(\omega_{\mathcal{X}_s}) = \deg_Z(\omega_{\mathcal{X}}) = \deg_{\Sigma_s(Z)}(\omega_{\mathcal{X}}) = \deg_{\Sigma_s(Z)}(\omega_X) \equiv \delta_{\Sigma_s(Z)}, \end{cases} \tag{4.2}$$

where $Z \subseteq \mathcal{X}_s$ is any subcurve of $\mathcal{X}_s$ and $\equiv$ denotes congruence modulo 2. □

Given a general polarization $\underline{q}$ on X, we get an open subset of $\overline{\mathbb{J}}_X$ which is proper over $\mathrm{Spec}\, \mathrm{R_X}$.

Theorem 4.2. *Let $\underline{q}$ be a general polarization on X. Then there exists an open subscheme $\overline{J}_{\mathcal{X}}(\underline{q}) \subseteq \overline{\mathbb{J}}_{\mathcal{X}}$ which is projective over $\mathrm{SpecR_X}$ and such that the fiber of $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{SpecR_X}$ over a point $s \in \mathrm{SpecR_X}$ is isomorphic to $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$. In particular, the fiber of $\overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{Spec}\, \mathrm{R_X}$ over the closed point $[\mathfrak{m}_X] \in \mathrm{Spec}\, \mathrm{R_X}$ is isomorphic to $\overline{J}_X(\underline{q})$.*

We call the scheme $\overline{J}_{\mathcal{X}}(\underline{q})$ the *universal fine compactified Jacobian* of X with respect to the polarization $\underline{q}$. We denote by $J_{\mathcal{X}}(\underline{q})$ the open subset of $\overline{J}_{\mathcal{X}}(\underline{q})$ parametrizing line bundles, i.e. $J_{\mathcal{X}}(\underline{q}) = \overline{J}_{\mathcal{X}}(\underline{q}) \cap \mathbb{J}_{\mathcal{X}} \subseteq \overline{\mathbb{J}}_{\mathcal{X}}$.

Proof. This statement follows by applying to the effective semiuniversal family $\mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$ a general result of Esteves ([Est01, Thm. A]). In order to connect our notations with the notations of loc. cit., take $r > 0$ a sufficiently divisible natural number such that, for every subcurve $W \subseteq X$, it holds that

$$r\left(\underline{q}_W - \frac{\deg_W(\omega_X)}{2}\right) \in \mathbb{Z}.$$

Consider a vector bundle E on X of rank r such that, for every subcurve $W \subseteq X$ (or equivalently for every irreducible component of X), the degree of E restricted to W is equal to

$$-\deg(E_{|W}) = r\left(\underline{q}_W - \frac{\deg_W(\omega_X)}{2}\right). \tag{4.3}$$

As it was already observed in [MV12, Rmk. 1.16(i)], it is easy to check that a torsion-free rank-1 sheaf I on X is (semi)stable with respect to $\underline{q}$ in the sense of Definition 2.8 if and only if it is (semi)stable with respect to E in the sense of [Est01, Sec. 1.2]. Therefore, our fine compactified Jacobian $\overline{J}_X(\underline{q})$ coincides with the variety $J_E^s = J_E^{ss}$ in [Est01, Sec. 4].

Since an obstruction space for the functor of deformations of E is $H^2(X, E \otimes E^\vee)$ (see e.g. [FGA05, Thm. 8.5.3(b)]) and since this latter group is zero because X is a curve, we get that E can be extended to a vector bundle $\overline{\mathcal{E}}$ on the formal semiuniversal deformation $\overline{\mathcal{X}} \to \mathrm{Spec}\mathrm{R}_\mathrm{X}$ of X. However, by Grothendieck's algebraization theorem for coherent sheaves (see [FGA05, Thm. 8.4.2]), the vector bundle $\overline{\mathcal{E}}$ is the completion of a vector bundle $\mathcal{E}$ on the effective semiuniversal deformation $\pi : \mathcal{X} \to \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ of X. Note that the restriction of $\mathcal{E}$ to the central fiber of π is isomorphic to the vector bundle E on X. Denote by $\mathcal{E}_s$ the restriction of $\mathcal{E}$ to the fiber $\mathcal{X}_s$. By construction, it is easy to see that for any $s \in \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ and any subcurve $Z \subseteq \mathcal{X}_s$, we have that

$$\deg_Z(\mathcal{E}_s) = \deg_Z(\mathcal{E}) = \deg_{\Sigma_s(Z)}(\mathcal{E}) = \deg_{\Sigma_s(Z)}(E).$$

Combining the previous equality with the equality $\deg_Z(\omega_{\mathcal{X}_s}) = \deg_{\Sigma_s(Z)}(\omega_X)$ (see (4.2)) and tha fact that $\underline{q}^s_Z = \underline{q}_{\Sigma_s(Z)}$ (see Lemma-Definition 4.1), and applying (4.3) to the subcurve $\Sigma_s(Z) \subseteq X$, it follows that

$$-\deg((\mathcal{E}_s)_{|Z}) = r\left(\underline{q}^s_Z - \frac{\deg_Z(\omega_{\mathcal{X}_s})}{2}\right).$$

Therefore, exactly as before, we get that a torsion-free rank-1 sheaf $\mathcal{I}$ on $\mathcal{X}$, flat on $\mathrm{Spec}\mathrm{R}_\mathrm{X}$, is (semi)stable with respect to $\mathcal{E}$ in the sense of [Est01, Sec. 1.4] if and only if for every $s \in \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ the restriction $\mathcal{I}_s$ of $\mathcal{I}$ to $\mathcal{X}_s$ is (semi)stable with respect to $\underline{q}^s$ in the sense of Definition 2.8. Since all the polarizations $\underline{q}^s$ are general by Lemma-Definition 4.1, we get that the open subscheme $\overline{J}_{\mathcal{X}}(\underline{q}) := J^{\mathrm{s}}_{\mathcal{E}} = J^{\mathrm{ss}}_{\mathcal{E}} \subset \mathbb{J}_{\mathcal{X}}$ parametrizing sheaves $\mathcal{I} \in \mathbb{J}_{\mathcal{X}}$ whose restriction to $\mathcal{X}_s$ is $\underline{q}^s$-semistable (or equivalently $\underline{q}^s$-stable) is a proper scheme over $\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ by [Est01, Thm. A]. Moreover, $J^{\mathrm{s}}_{\mathcal{E}}$ is quasi-projective over $\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ by [Est01, Thm. C]; hence it is projective over $\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$. The description of the fibers of $\overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ is now clear from the definition of $\overline{J}_{\mathcal{X}}(\underline{q})$.

□

If the curve X has locally planar singularities, then the universal fine compactified Jacobians of X have several nice properties that we collect in the following statement.

Theorem 4.3. *Assume that X has locally planar singularities and let $\underline{q}$ be a general polarization on X. Then we have:*

(i) The scheme $\overline{J}_{\mathcal{X}}(\underline{q})$ is smooth and irreducible.
(ii) The surjective map $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ is projective and flat of relative dimension $p_a(X)$.
(iii) The smooth locus of u is $J_{\mathcal{X}}(\underline{q})$.

Proof. The smoothness of $\overline{J}_{\mathcal{X}}(\underline{q})$ follows from Theorem 3.7(iii).

Since also $\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ is smooth by Theorem 3.7(ii), the flatness of the map $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ will follow if we show that all the fibers are equi-dimensional of the same dimension (see [Mat89, Cor. of Thm 23.1, p. 179]). By Theorem 4.2, the fiber of u over $s \in \mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ is isomorphic to $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ which has pure dimension equal to $h^1(\mathcal{X}_s, \mathcal{O}_{\mathcal{X}_s}) = h^1(X, \mathcal{O}_X) = p_a(X)$ by Corollary 2.12.

The map u is projective by Theorem 4.2 and the fact that its smooth locus is equal to $J_{\mathcal{X}}(\underline{q})$ follows from Corollary 2.12.

Finally, in order to show that $\overline{J}_{\mathcal{X}}(\underline{q})$ is irreducible, it is enough to show that it is connected since we know already that $\overline{J}_{\mathcal{X}}(\underline{q})$ is smooth. Since the open subset $J_{\mathcal{X}}(\underline{q})$ is dense by Corollary 2.12, it is enough to prove that $J_{\mathcal{X}}(\underline{q})$ is connected. However, this follows easily from the fact that $J_{\mathcal{X}}(\underline{q})$ is smooth over $\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ and its generic fiber is the Jacobian of degree $|\underline{q}|$ of a smooth curve, hence it is connected.

□

The above result on the universal fine compactified Jacobians of X has also some very important consequences for the fine compactified Jacobians of X, that we collect in the following two corollaries.

Corollary 4.4. *Assume that X has locally planar singularities and let $\underline{q}$ be a general polarization on X. Then $\overline{J}_X(\underline{q})$ is connected.*

Proof. Consider the universal fine compactified Jacobian $\overline{J}_{\mathcal{X}}(\underline{q})$ and the natural surjective morphism $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R_X}$. According to Theorem 3.7 and Theorem 4.3, u is a projective morphism between two smooth and irreducible Noetherian schemes. Consider the Stein factorization of u:

$$u : \overline{J}_{\mathcal{X}}(\underline{q}) \xrightarrow{f} Z \xrightarrow{g} \operatorname{Spec} \mathrm{R_X},$$

where f has connected fibers and g is a finite surjective morphism. Note that the generic fiber of u is the Jacobian of degree $|\underline{q}|$ of a smooth curve, hence it is connected. This implies that g is generically injective, hence birational. Since $\operatorname{Spec} \mathrm{R_X}$ is normal, Zariski's main theorem gives that g is an isomorphism. We conclude that $u = f$ has connected fibers; in particular, $\overline{J}_X(\underline{q}) = u^{-1}[\mathfrak{m}_X]$ is connected.

□

Corollary 4.5. *Assume that X has locally planar singularities and let $\underline{q}$ be a general polarization on X. Then the universal fine compactified Jacobian $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R_X}$ (with respect to the polarization $\underline{q}$) has trivial relative dualizing sheaf. In particular, $\overline{J}_X(\underline{q})$ has trivial dualizing sheaf.*

Proof. Observe that the relative dualizing sheaf, call it ω_u, of the universal fine compactified Jacobian $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R_X}$ is a line bundle because the fibers of u have l.c.i. singularities by Theorem 4.2 and Corollary 2.12.

CLAIM 1: $(\omega_u)_{|J_{\mathcal{X}}(\underline{q})} = \mathcal{O}_{J_{\mathcal{X}}(\underline{q})}$.

Indeed, Theorem 4.2 implies that the fiber of $J_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R_X}$ over a point s is isomorphic $J_{\mathcal{X}_s}(\underline{q}^s)$ which is a disjoint union of a certain number of copies of the generalized Jacobian $J(\mathcal{X}_s)$. In particular, the fibers of the map $J_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R_X}$ have trivial canonical sheaf. Therefore, Claim 1 follows from the see-saw principle (see e.g. [Ser06, Sec. 4.2, Note 2]) together with the fact that $\operatorname{Pic}(\operatorname{Spec} \mathrm{R_X}) = 0$.

CLAIM 2: The codimension of $\overline{J}_{\mathcal{X}}(\underline{q}) \setminus J_{\mathcal{X}}(\underline{q})$ in $\overline{J}_{\mathcal{X}}(\underline{q})$ is at least two.

Indeed, for every $s \in \operatorname{Spec} \mathrm{R_X}$, the codimension of $\overline{J}_{\mathcal{X}_s}(\underline{q}^s) \setminus J_{\mathcal{X}_s}(\underline{q}^s)$ inside $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ is at least one since $J_{\mathcal{X}_s}(\underline{q}^s)$ is dense in $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ by Corollary 2.12. Moreover, for the general point $s_0 \in \operatorname{Spec} \mathrm{R_X}$, the fiber $\mathcal{X}_{s_0}$ is smooth since $\mathrm{Def}_{\mathrm{X}}^{\mathrm{loc}}$ is smooth by Fact 3.1(ii). Therefore, clearly we have that $J_{\mathcal{X}_{s_0}}(\underline{q}^{s_0}) = \overline{J}_{\mathcal{X}_{s_0}}(\underline{q}^{s_0})$. Claim 2 now follows.

Claim 1 and Claim 2 imply that ω_u and $\mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}$ agree on an open subset whose complement has codimension at least two. Since $\overline{J}_{\mathcal{X}}(\underline{q})$ is smooth (hence S_2) by Theorem 4.3, this implies that $\omega_u = \mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}$.

The second assertion follows now by restricting the equality $\omega_u = \mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}$ to the fiber $\overline{J}_X(\underline{q})$ of u over the closed point $[\mathfrak{m}_X] \in \operatorname{Spec} \mathrm{R_X}$. □

Note that a statement similar to Corollary 4.5 was proved by Arinkin in [Ari11, Cor. 9] for the universal compactified Jacobian over the moduli stack of *integral* curves with locally planar singularities.

Finally, note that the universal fine compactified Jacobians are acted upon by the universal generalized Jacobian, whose properties are collected into the following

Fact 4.6 (Bosch-Lütkebohmert-Raynaud). *There is an open subset of $\mathbb{J}_{\mathcal{X}}$, called the* universal generalized Jacobian *of $\pi : \mathcal{X} \to \operatorname{Spec}\mathrm{R_X}$ and denoted by $v : J(\mathcal{X}) \to \operatorname{Spec}\mathrm{R_X}$, whose fiber over any point $s \in \operatorname{Spec}\mathrm{R_X}$ is the generalized Jacobian $J(\mathcal{X}_s)$ of the fiber $\mathcal{X}_s$ of π over s.*

The morphism v makes $J(\mathcal{X})$ into a smooth and separated group scheme of finite type over $\operatorname{Spec} \mathrm{R_X}$.

Proof. The existence of a group scheme $v : J(\mathcal{X}) \to \operatorname{Spec} \mathrm{R_X}$ whose fibers are the generalized Jacobians of the fibers of $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R_X}$ follows by [BLR90, Sec. 9.3, Thm. 7], which can be applied since $\operatorname{Spec} \mathrm{R_X}$ is a strictly henselian local scheme (because R_X is a complete local ring) and the fibers of $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R_X}$ are geometrically reduced and connected since X is assumed to be so. The result of loc. cit. gives also that the map v is smooth, separated and of finite type. □

We denote by $\zeta : \operatorname{Spec} \mathrm{R_X} \to \mathrm{J}(\mathcal{X})$ the zero section of the group scheme $v : J(\mathcal{X}) \to \operatorname{Spec} \mathrm{R_X}$; in other words, ζ is the morphism which sends every point $s \in \operatorname{Spec} \mathrm{R_X}$ into the trivial line bundle on the fiber $\mathcal{X}_s$ of $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R_X}$ over s.

4.1. **1-parameter regular smoothings of X.** The aim of this subsection is to study relative fine compactified Jacobians associated to a 1-parameter smoothing of a curve X and their relationship with the Néron models of the Jacobians of the generic fiber. As a corollary, we will get a combinatorial formula for the number of irreducible components of a fine compactified Jacobian of a curve with locally planar singularities.

Let us start with the definition of 1-parameter regular smoothings of a curve X.

Definition 4.7. A *1-parameter regular smoothing* of X is a morphism $f : \mathcal{S} \to B = \mathrm{Spec\,R}$ where R is a complete discrete valuation domain (DVR for short) with residue field k and quotient field K and $\mathcal{S}$ is a regular surface and such that the special fiber $\mathcal{S}_k$ is isomorphic to X and the generic fiber $\mathcal{S}_K$ is a smooth curve.

The natural question one may ask is the following: which (reduced) curves X admit a 1-parameter regular smoothing? Of course, if X admits a 1-parameter regular smoothing $f : \mathcal{S} \to \mathrm{Spec\,R}$, then X is a divisor inside a regular surface $\mathcal{S}$, which implies that X has locally planar singularities. Indeed, it is well known to the experts that this necessary condition turns out to be also sufficient. We include a proof here since we couldn't find a suitable reference.

Lemma 4.8. *A (reduced) curve X admits a 1-parameter regular smoothing if and only if X has locally planar singularities. More precisely, if X has locally planar singularities then there exists a complete discrete valuation domain R and a morphism $\alpha : \mathrm{Spec\,R} \to \mathrm{Spec\,R_X}$ such that the pull-back*

$$\begin{tikzcd} \mathcal{S} \arrow[r] \arrow[d, "f"'] \arrow[dr, phantom, "\square"] & \mathcal{X} \arrow[d, "\pi"] \\ B = \mathrm{Spec\,R} \arrow[r, "\alpha"] & \mathrm{Spec\,R_X} \end{tikzcd} \tag{4.4}$$

is a 1-parameter regular smoothing of X.

Proof. We have already observed that the only if condition is trivially satisfied. Conversely, assume that X has locally planar singularities, and let us prove that X admits a 1-parameter regular smoothing.

Consider the natural morphisms of deformation functors

$$F : h_{R_X} \to \mathrm{Def_X} \to \mathrm{Def_X^{loc}} = \prod_{\mathrm{p}\in \mathrm{X_{sing}}} \mathrm{Def_{X,p}},$$

obtained by composing the morphism (3.1) with the morphism (3.5). Observe that F is smooth because the first morphism is smooth since R_X is a semiuniversal deformation ring for $\mathrm{Def_X}$ and the second morphism is smooth by Fact 3.1(iii).

Given an element $\alpha \in h_{R_X}(R) = \mathrm{Hom}(\mathrm{Spec\,R}, \mathrm{Spec\,R_X})$ associated to a Cartesian diagram like in (4.4), the image of α into $\mathrm{Def_{X,p}(R)}$ corresponds to the formal deformation of (X, p) given by

$$\begin{tikzcd} \mathrm{Spf}\,\widehat{\mathcal{O}}_{\mathrm{X,p}} \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & \mathrm{Spf}\,\widehat{\mathcal{O}}_{\mathcal{S},\mathrm{p}} \arrow[d] \\ \mathrm{Spf\,k} \arrow[r] & \mathrm{Spf\,R.} \end{tikzcd}$$

Clearly, in order to check that $f : \mathcal{S} \to B = \mathrm{Spec R}$ is a 1-parameter regular smoothing of X it is sufficient (and necessary) to check that the ring $\widehat{\mathcal{O}}_{\mathcal{S},p}$ is regular and that $\widehat{\mathcal{O}}_{\mathcal{S},p} \otimes_R K$ is K-smooth, where K is the quotient field of R. Conversely, if we find for each $p \in X_{\mathrm{sing}}$ an element of $\mathrm{Def_{X,p}(R)}$ corresponding to a formal deformation

$$\begin{tikzcd} \mathrm{Spf}\,\widehat{\mathcal{O}}_{\mathrm{X,p}} \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & \mathrm{Spf}\,\mathcal{A} \arrow[d] \\ \mathrm{Spf\,k} \arrow[r] & \mathrm{Spf\,R.} \end{tikzcd}$$

such that $\mathcal{A}$ is regular and $\mathcal{A} \otimes_R K$ is K-smooth, then using the smoothness of F we can lift this element to an element $\alpha \in h_{R_X}(R)$ which will have the desired properties.

Let us now check this local statement. Since X has locally planar singularities at $p \in X_{\mathrm{sing}}$, we can write

$$\widehat{O}_{X,p} = \frac{k[[x, y]]}{(f)},$$

where $f = f(x, y) \in k[[x, y]]$ is reduced and such that $(0, 0)$ is the unique singular point of $\{f = 0\}$. This is equivalent to say that $(0, 0)$ is the unique solution of the system $f = \dfrac{\partial f}{\partial x} = \dfrac{\partial f}{\partial y} = 0$. Take now $R := k[[t]]$ and set $\mathcal{A} := \dfrac{k[[x, y, t]]}{(f + t)}$. The ring $\mathcal{A}$ is regular since $\dfrac{\partial (f + t)}{\partial t} = 1$ does not vanish anywhere. The ring

$\mathcal{A} \otimes_R K = \dfrac{K[[x,y]]}{(f+t)}$ is K-smooth since the system of equations $(f+t) = \dfrac{\partial(f+t)}{\partial x} = \dfrac{\partial(f+t)}{\partial y} = 0$ does not have any solution. This concludes the proof of the local statement and along with it the proof of the Lemma. ☐

From now till the end of this subsection, we fix a 1-parameter regular smoothing $f : \mathcal{S} \to B = \operatorname{Spec} \mathrm{R}$ of X as in Lemma 4.8. Let $\mathcal{P}ic_f$ denote the *relative Picard functor* of f (often denoted $\mathcal{P}ic_{\mathcal{S}/B}$ in the literature, see [BLR90, Chap. 8] for the general theory) and let $\mathcal{P}ic_f^d$ be the subfunctor parametrizing line bundles of relative degree $d \in \mathbb{Z}$. The functor $\mathcal{P}ic_f$ (resp. $\mathcal{P}ic_f^d$) is represented by a scheme Pic_f (resp. Pic_f^d) locally of finite presentation over B (see [BLR90, Sec. 8.2, Thm. 2]) and smooth over B (by [BLR90, Sec. 8.4, Prop. 2]). The generic fiber of Pic_f (resp. Pic_f^d) is isomorphic to $\operatorname{Pic}(\mathcal{S}_K)$ (resp. $\operatorname{Pic}^d(\mathcal{S}_K)$).

Note that Pic_f^d is not separated over B whenever X is reducible. The biggest separated quotient of Pic_f^d coincides with the *Néron model* $N(\operatorname{Pic}^d \mathcal{S}_K)$ of $\operatorname{Pic}^d(\mathcal{S}_K)$, as proved by Raynaud in [Ray70, Sec. 8] (see also [BLR90, Sec. 9.5, Thm. 4]). Recall that $N(\operatorname{Pic}^d \mathcal{S}_K)$ is smooth and separated over B, the generic fiber $N(\operatorname{Pic}^d \mathcal{S}_K)_K$ is isomorphic to $\operatorname{Pic}^d \mathcal{S}_K$ and $N(\operatorname{Pic}^d \mathcal{S}_K)$ is uniquely characterized by the following universal property (the Néron mapping property, cf. [BLR90, Def. 1]): every K-morphism $q_K : Z_K \to N(\operatorname{Pic}^d \mathcal{S}_K)_K = \operatorname{Pic}^d \mathcal{S}_K$ defined on the generic fiber of some scheme Z smooth over B admits a unique extension to a B-morphism $q : Z \to N(\operatorname{Pic}^d \mathcal{S}_K)$. Moreover, $N(\operatorname{Pic}^0 \mathcal{S}_K)$ is a B-group scheme while, for every $d \in \mathbb{Z}$, $N(\operatorname{Pic}^d S_K)$ is a torsor under $N(\operatorname{Pic}^0 \mathcal{S}_K)$.

Fix now a general polarization $\underline{q}$ on X and consider the Cartesian diagram

(4.5)
$$\begin{tikzcd}
\overline{J}_f(\underline{q}) \arrow[r] \arrow[d] \arrow[dr, phantom, "\square"] & \overline{J}_{\mathcal{X}}(\underline{q}) \arrow[d, "u"] \\
B = \operatorname{Spec} \mathrm{R} \arrow[r, "\alpha"] & \operatorname{Spec} \mathrm{R}_X
\end{tikzcd}$$

We call the scheme $\overline{J}_f(\underline{q})$ the *f-relative fine compactified Jacobian* with respect to the polarization $\underline{q}$. Similarly, by replacing $\overline{J}_{\mathcal{X}}(\underline{q})$ with $J_{\mathcal{X}}(\underline{q})$ in the above diagram, we define the open subset $J_f(\underline{q}) \subset \overline{J}_f(\underline{q})$. Note that the generic fibers of $\overline{J}_f(\underline{q})$ and $J_f(\underline{q})$ coincide and are equal to $\overline{J}_f(\underline{q})_K = J_f(\underline{q})_K = \operatorname{Pic}^{|\underline{q}|}(\mathcal{S}_K)$ while their special fibers are equal to $\overline{J}_f(\underline{q})_k = \overline{J}_X(\underline{q})$ and $J_f(\underline{q})_k = J_X(\underline{q})$, respectively. From Theorem 4.3, we get that the morphism $\overline{J}_f(\underline{q}) \to B$ is flat and its smooth locus is $J_f(\underline{q})$. Therefore, the universal property of the Néron model $N(\operatorname{Pic}^{|\underline{q}|} \mathcal{S}_K)$ gives a unique B-morphism $q_f : J_f(\underline{q}) \to N(\operatorname{Pic}^{|\underline{q}|} \mathcal{S}_K)$ which is the identity on the generic fiber. Indeed, J. L. Kass proved in [Kas09, Thm. A] that the above map is an isomorphism.

Fact 4.9 (Kass)**.** *For a 1-paramater regular smoothing $f : \mathcal{X} \to B = \operatorname{Spec} \mathrm{R}$ as above and any general polarization $\underline{q}$ on X, the natural B-morphism*

$$q_f : J_f(\underline{q}) \to N(\operatorname{Pic}^{|\underline{q}|} \mathcal{S}_K)$$

is an isomorphism.

From the above isomorphism, we can deduce a formula for the number of irreducible components of $\overline{J}_X(\underline{q})$. We first need to recall the description due to Raynaud of the group of connected components of the Néron model $N(\operatorname{Pic}^0(\mathcal{S}_K))$ (see [BLR90, Sec. 9.6] for a detailed exposition).

Denote by $C_1, \ldots, C_\gamma$ the irreducible components of X. A *multidegree* on X is an ordered γ-tuple of integers

$$\underline{d} = (\underline{d}_{C_1}, \ldots, \underline{d}_{C_\gamma}) \in \mathbb{Z}^\gamma.$$

We denote by $|\underline{d}| = \sum_{i=1}^{\gamma} \underline{d}_{C_i}$ the total degree of $\underline{d}$. We now define an equivalence relation on the set of multidegrees on X. For every irreducible component C_i of X, consider the multidegree $\underline{C_i} = ((\underline{C_i})_1, \ldots, (\underline{C_i})_\gamma)$ of total degree 0 defined by

$$(\underline{C_i})_j = \begin{cases} |C_i \cap C_j| & \text{if } i \neq j, \\ -\displaystyle\sum_{k \neq i} |C_i \cap C_k| & \text{if } i = j, \end{cases}$$

where $|C_i \cap C_j|$ denotes the length of the scheme-theoretic intersection of C_i and C_j. Clearly, if we take a 1-parameter regular smoothing $f : \mathcal{S} \to B$ of X as in Lemma 4.8, then $|C_i \cap C_j|$ is also equal to the intersection product of the two divisors C_i and C_j inside the regular surface $\mathcal{S}$.

Denote by $\Lambda_X \subseteq \mathbb{Z}^\gamma$ the subgroup of $\mathbb{Z}^\gamma$ generated by the multidegrees $\underline{C_i}$ for $i = 1, \ldots, \gamma$. It is easy to see that $\sum_i \underline{C_i} = 0$ and this is the only relation among the multidegrees $\underline{C_i}$. Therefore, Λ_X is a free abelian group of rank $\gamma - 1$.

Definition 4.10. Two multidegrees $\underline{d}$ and $\underline{d}'$ are said to be *equivalent*, and we write $\underline{d} \equiv \underline{d}'$, if $\underline{d} - \underline{d}' \in \Lambda_X$. In particular, if $\underline{d} \equiv \underline{d}'$ then $|\underline{d}| = |\underline{d}'|$.

For every $d \in \mathbb{Z}$, we denote by Δ_X^d the set of equivalence classes of multidegrees of total degree $d = |\underline{d}|$. Clearly Δ_X^0 is a finite group under component-wise addition of multidegrees (called the *degree class group* of X) and each Δ_X^d is a torsor under Δ_X^0. The cardinality of Δ^0 is called the *degree class number* or the *complexity* of X, and it is denoted by $c(X)$.

The name degree class group was first introduced by L. Caporaso in [Cap94, Sec. 4.1]. The name complexity comes from the fact that if X is a nodal curve then $c(X)$ is the complexity of the dual graph Γ_X of X, i.e. the number of spanning trees of Γ_X (see e.g. [MV12, Sec. 1.2]).

Fact 4.11 (Raynaud). *The group of connected component of the B-group scheme $N(\operatorname{Pic}^0 \mathcal{S}_K)$ is isomorphic to Δ_X^0. In particular, the special fiber of $N(\operatorname{Pic}^d \mathcal{S}_K)$ for any $d \in \mathbb{Z}$ is isomorphic to the disjoint union of $c(X)$ copies of the generalized Jacobian $J(X)$ of X.*

For a proof, see the original paper of Raynaud [Ray70, Prop. 8.1.2] or [BLR90, Sec. 9.6].

Finally, by combining Corollary 2.12, Fact 4.9 and Fact 4.11, we obtain a formula for the number of irreducible components of a fine compactified Jacobian.

Corollary 4.12. *Assume that X has locally planar singularities and let $\underline{q}$ be any general polarization on X. Then $\overline{J}_X(\underline{q})$ has $c(X)$ irreducible components.*

5. The Picard scheme of the universal fine compactified Jacobians

The aim of this section is to discuss the properties of the Picard scheme of the universal fine compactified Jacobians $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R_X}$, introduced in Section 4. Following [FGA05, Chap. 9, Sec. 2] and [BLR90, Sec. 8.1], we define the *relative Picard functor* Pic_u as the fppf-sheaf associated to the controvariant Picard functor

$$\operatorname{Pic}_u : \mathrm{Sch}/R_X \to \mathrm{Grps},$$
$$T \mapsto \mathcal{P}ic(\overline{J}_{\mathcal{X}}(\underline{q}) \times_{\operatorname{Spec} \mathrm{R_X}} T),$$

where Sch/R_X is the category of schemes over $\operatorname{Spec} \mathrm{R_X}$, Grps is the category of abelian groups and $\mathcal{P}ic$ denotes the Picard group as defined in 1.8.

Following [FGA05, Chap. 9, Sec. 5, 6] and [BLR90, Sec. 8.4], we consider the two subfunctors of the relative Picard functor

$$\operatorname{Pic}_u^o \subseteq \operatorname{Pic}_u^\tau \subseteq \operatorname{Pic}_u, \tag{5.1}$$

such that Pic_u^o (resp. $\operatorname{Pic}_u^\tau$) consists of the elements of Pic_u whose restriction to all the fibers $\mathcal{X}_s$ for $s \in \operatorname{Spec} \mathrm{R_X}$ belongs to $\operatorname{Pic}^o(\mathcal{X}_s)$ (resp. $\operatorname{Pic}^\tau(\mathcal{X}_s)$), see 1.8.

We summarize the properties of the above functors in the following

Theorem 5.1. *Let X be a curve with locally planar singularities of arithmetic genus $p_a(X)$ and let $\underline{q}$ be a general polarization on X.*

- *(i)* Pic_u *is represented by a group scheme* $\operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))$ *locally of finite presentation over* $\operatorname{Spec} \mathrm{R_X}$.
- *(ii)* $\operatorname{Pic}_u^\tau$ *is represented by an open subgroup scheme* $\operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})) \subseteq \operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))$ *which is of finite type and separated over* $\operatorname{Spec} \mathrm{R_X}$.
- *(iii) If* $h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X)$, *then* $\operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))$ *is smooth over* $\operatorname{Spec} \mathrm{R_X}$.
- *(iv) Assume that* $h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X)$. *Then* Pic_u^o *is represented by an open subgroup scheme* $\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \subseteq \operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))$, *which is smooth, separated and of finite type over* $\operatorname{Spec} \mathrm{R_X}$.

Proof. Since R_X is strictly henselian (being a complete local ring with algebraically closed residue field), the fibers of u are such that their irreducible components are geometrically irreducible, see [Est01, Lemma 18]. Moreover, the morphism u is projective and flat by Theorem 4.3(ii) and the fibers of u are geometrically connected by Corollary 4.4. Therefore, we can apply Mumford's representability criterion for the Picard scheme (see [BLR90, Sec. 8.2, Thm. 2] or [FGA05, Chap. 9, Thm. 4.18.1]) in order to deduce that the relative Picard functor Pic_u is represented by a group scheme locally of finite type over $\mathrm{Spec}\,\mathrm{R_X}$, which proves part (i).

Since u is proper and Pic_u is represented by a scheme, a result of Kleimann [Kle71, Thm. 5.7] gives that Pic_u^τ is represented by an open subgroup scheme $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})) \subseteq \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))$ which is moreover of finite type over $\mathrm{Spec}\,\mathrm{R_X}$. In order to prove that $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))$ is separated over $\mathrm{Spec}\,\mathrm{R_X}$, it is enough to prove, using the valuative criterion of separatedness (see [EGAII, Prop. 7.2.3]), that for any map $\mathrm{Spec\,R} \to \mathrm{Spec}\,\mathrm{R_X}$, where R is a discrete valuation ring, the base change map

$$f_R : \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})) \times_{\mathrm{Spec}\,\mathrm{R_X}} \mathrm{Spec\,R} \cong \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{\mathrm{q}}) \times_{\mathrm{Spec}\,\mathrm{R_X}} \mathrm{Spec\,R}) \to \mathrm{Spec\,R}$$

is separated. Since the fibers of $\overline{J}_{\mathcal{X}}(\underline{q}) \times_{\mathrm{Spec}\,\mathrm{R_X}} \mathrm{Spec\,R} \to \mathrm{Spec\,R}$ are geometrically reduced by Corollary 2.12 and R is a discrete valuation ring, a result of Raynaud [Ray70, Cor. 6.4.5] guarantees that the map f_R is separated. Part (ii) is now proved.

Before proving the remaining assertions, we prove the following

Claim: If $h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X)$, then the fibers of $f : \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q})) \to \mathrm{Spec}\,\mathrm{R_X}$ are smooth of dimension $p_a(X)$.

Indeed, if s is a general point of $\mathrm{Spec}\,\mathrm{R_X}$, then $\mathcal{X}_s$ is a smooth projective curve of genus equal to $p_a(X)$ by Fact (3.1)(iii) and therefore $\overline{J}_{\mathcal{X}_s}(\underline{q})$ is an abelian variety of dimension equal to $p_a(X)$. This implies that $\mathrm{Pic}(\overline{J}_{\mathcal{X}_s}(\underline{q}))$ is a smooth group scheme of dimension $p_a(X)$ over the residue field $k(s)$ of s. By the upper semicontinuity of the dimension of the fibers of f (see [EGAIV3, (13.1.3)]), we deduce that

$$\dim \mathrm{Pic}(\overline{J}_{\mathcal{X}_s}(\underline{q})) \geq p_a(X) \text{ for any } s \in S. \tag{5.2}$$

On the other hand, if $h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X)$ then by upper semicontinuity of the cohomology groups, we get that

$$h^1(\overline{J}_{\mathcal{X}_s}(\underline{q}), \mathcal{O}_{\overline{J}_{\mathcal{X}_s}(\underline{q})}) \leq p_a(X) \text{ for any } s \in S. \tag{5.3}$$

Combining (5.2) and (5.3), we infer that $\dim \mathrm{Pic}(\overline{J}_{\mathcal{X}_s}(\underline{q})) \geq h^1(\overline{J}_{\mathcal{X}_s}(\underline{q}), \mathcal{O}_{\overline{J}_{\mathcal{X}_s}(\underline{q})})$ for any $s \in S$. This implies that, for every $s \in S$, $\mathrm{Pic}(\overline{J}_{\mathcal{X}_s}(\underline{q}))$ is smooth of dimension equal to $p_a(X)$ (see [FGA05, Chap. 9, Cor. 5.13]) which proves the claim.

Let us now prove part (iii). Since f is locally of finite type by part (i), equidimensional and the codomain $\mathrm{Spec}\,\mathrm{R_X}$ of f is geometrically unibranch (being regular), then f is universally open by Chevalley's criterion ([EGAIV3, (14.4.3)]). Moreover, since f is universally open and locally of finite type, the fibers of f are geometrically reduced (being smooth) and the codomain $\mathrm{Spec}\,\mathrm{R_X}$ of f is locally Noetherian and reduced (being Noetherian and regular), then we conclude that f is flat by [EGAIV3, (15.2.3)]. Finally, since f is flat and locally of finite presentation (being locally of finite type over a Noetherian codomain) and the fibers of f are smooth, we conclude that f is smooth by [BLR90, Sec. 2.4, Prop. 8], q.e.d.

Part (iv): since the fibers of f are smooth of the same dimension, the representability of Pic_u^o by an open subgroup scheme $\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \subseteq \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))$ of finite type over $\mathrm{Spec}\,\mathrm{R_X}$ follows from [FGA05, Chap. 9, Prop. 5.20]. The morphism $\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to \mathrm{Spec}\,\mathrm{R_X}$ is separated by (ii) and smooth by (iii). □

6. ABEL MAPS

The aim of this section is to define Abel maps for singular (reduced) curves. Although in the following sections we will only use Abel maps for curves with locally planar singularities, the results of this section are valid for a broader class of singular curves, namely those for which every separating point is a node (see Condition (6.3)), which includes for example all Gorenstein curves.

6.1. Abel maps without separating points.

The aim of this subsection is to define the Abel maps for a reduced curve X without separating points (in the sense of 1.7).

For every (geometric) point p on the curve X, its sheaf of ideals $\mathfrak{m}_p$ is a torsion-free rank-1 sheaf of degree -1 on X. Also, if p is not a separating point of X, then $\mathfrak{m}_p$ is simple (see [Est01, Example 38]). Therefore, if X does not have separating points (which is clearly equivalent to the fact that $\delta_Y \geq 2$ for every proper subcurve Y of X) then $\mathfrak{m}_p$ is torsion-free rank-1 and simple for any $p \in X$.

Let $\mathcal{I}_\Delta$ be the ideal of the diagonal Δ of $X \times X$. For any line bundle $L \in \mathrm{Pic}(X)$, consider the sheaf $\mathcal{I}_\Delta \otimes p_1^* L$, where $p_1 : X \times X \to X$ denotes the projection onto the first factor. The sheaf $\mathcal{I}_\Delta \otimes p_1^* L$ is flat over X (with respect to the second projection $p_2 : X \times X \to X$) and for any point p of X it holds

$$\mathcal{I}_\Delta \otimes p_1^* L_{|X \times \{p\}} = \mathfrak{m}_p \otimes L.$$

Therefore, if X does not have separating points, then $\mathcal{I}_\Delta \otimes p_1^* L \in \overline{\mathbb{J}}_X^*(X)$ where $\overline{\mathbb{J}}_X^*$ is the functor defined by (2.1). Using the universal property of $\overline{\mathbb{J}}_X$ (see Fact 2.2(iii)), the sheaf $\mathcal{I}_\Delta \otimes p_1^* L$ induces a morphism

$$\begin{aligned} A_L : X &\to \overline{\mathbb{J}}_X \\ p &\mapsto \mathfrak{m}_p \otimes L. \end{aligned} \tag{6.1}$$

We call the map A_L the *(L-twisted) Abel map* of X.

From the definition (2.4), it follows that a priori the Abel map A_L takes values in the big compactified Jacobian $\overline{\mathbb{J}}_X^{\deg L-1}$. Actually, we can prove that the Abel map A_L takes always values in a fine compactified Jacobian contained in $\overline{\mathbb{J}}_X^{\deg L-1}$.

Lemma 6.1. *For every $L \in \mathrm{Pic}(X)$ there exists a general polarization $\underline{q}$ on X of total degree $|\underline{q}| = \deg L - 1$ such that* $\mathrm{Im(A_L)} \subseteq \overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$.

Proof. Consider the polarization $\underline{q}'$ on X defined as (for every irreducible component C_i of X)

$$\underline{q}'_{C_i} = \deg_{C_i} L - \frac{1}{\gamma(X)},$$

where $\gamma(X)$ denotes, as usual, the number of irreducible components of X. Note that for any subcurve $Y \subseteq X$ we have that $\underline{q}'_Y = \deg_Y(L) - \frac{\gamma(Y)}{\gamma(X)}$ and in particular $|\underline{q}'| = \deg L - 1$.

We claim that every sheaf in the image of A_L is $\underline{q}'$-stable. Indeed, for any proper subcurve $\emptyset \neq Y \subsetneq X$ and any point $p \in X$, we have that

$$\deg_Y(\mathfrak{m}_p \otimes L) - \underline{q}'_Y = \deg_Y(L) + \deg_Y(\mathfrak{m}_p) - \underline{q}'_Y \geq \deg_Y(L) - 1 - \underline{q}'_Y = -1 + \frac{\gamma(Y)}{\gamma(X)} > -1 \geq \frac{\delta_Y}{2}, \tag{6.2}$$

where we have used that $\gamma(Y) > 0$ since Y is not the empty subcurve and that $\delta_Y \geq 2$ since X does not have separating points by assumption. Since the inequality in (6.2) is strict, we can deform slightly $\underline{q}'$ in order to obtain a general polarization $\underline{q}$ such that $|\underline{q}| = |\underline{q}'|$ and for which (6.2) is still satisfied. This is equivalent to say that $\mathrm{ImA_L} \subseteq \overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$, q.e.d. ☐

Those fine compactified Jacobians for which there exists an Abel map as in the above Lemma 6.1 are quite special, hence they deserve a special name.

Definition 6.2. Let X be a curve without separating points. A fine compactified Jacobian $\overline{J}_X(\underline{q})$ is said to *admit an Abel map* if there exists a line bundle $L \in \mathrm{Pic}^{|\underline{q}|+1}(X)$ such that $\mathrm{ImA_L} \subseteq \overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$.

Observe that clearly the property of admitting an Abel map is invariant under equivalence by translation (in the sense of Definition 2.13).

Remark 6.3. It is possible to prove (we leave the easy proof to the reader) that a curve X without separating points and having at most two irreducible components is such that any fine compactified Jacobian of X admits an Abel map.

However, in Section 7 we are going to show examples of curves with more than two components and having a fine compactified Jacobian which does not admit an Abel map (see Proposition 7.4 and Proposition 7.5). In particular, Proposition 7.4 shows that, as the number of irreducible components of X increases, fine compactified Jacobians of X that admit an Abel map became rare and rare.

6.2. **Abel maps with separating points.**

The aim of this subsection is to define Abel maps for (reduced) curves X having separating points (in the sense of (1.7)) but satisfying the following

(6.3) $\qquad$ $\underline{\text{Condition } (\dagger)}$: Every separating point is a node.

Indeed, there are plenty of curves that satisfy condition (†) due to the following result of Catanese (see [Cat82, Prop. 1.10]).

Fact 6.4 (Catanese)**.** *A (reduced) Gorenstein curve X satisfies condition* (†).

Let us give an example of a curve that do not satisfy condition (†).

Example 6.5. Consider a curve X made of three irreducible smooth components Y_1, Y_2 and Y_3 glued at one point $p \in Y_1 \cap Y_2 \cap Y_3$ with linearly independent tangent directions, i.e. in a such a way that, locally at p, the three components Y_1, Y_2 and Y_3 look like the three coordinate axes in $\mathbb{A}^3$. A straightforward local computation gives that $\delta_{Y_1} = |Y_1 \cap (Y_2 \cup Y_3)| = 1$, so that p is a separating point of X (in the sense of 1.7). However p is clearly not a node of X. Combined with Fact 6.4, this shows that X is not Gorenstein at $p \in X$.

Throughout this section, we fix a connected (reduced) curve X satisfying condition (†) and let $\{n_1, \ldots, n_{r-1}\}$ be its separating points. Since X satisfies condition (†), each n_i is a node. Denote by $\widetilde{X}$ the partial normalization of X at the set $\{n_1, \ldots, n_{r-1}\}$. Since each n_i is a node, the curve $\widetilde{X}$ is a disjoint union of r connected reduced curves $\{Y_1, \ldots, Y_r\}$ such that each Y_i does not have separating points. Note also that X has locally planar singularities if and only if each Y_i has locally planar singularities. We have a natural morphism

$$\tau : \widetilde{X} = \coprod_i Y_i \to X. \tag{6.4}$$

We can naturally identify each Y_i with a subcurve of X in such a way that their union is X and that they do not have common irreducible components. In particular, the irreducible components of X are the union of the irreducible components of the curves Y_i. We call the components Y_i (or their image in X) the *separating blocks* of X.

Let us first show how the study of fine compactified Jacobians of X can be reduced to the study of fine compactified Jacobians of Y_i. Observe that, given a polarization $\underline{q}^i$ on each curve Y_i, we get a polarization $\underline{q} := (\underline{q}^1, \ldots, \underline{q}^r)$ on X defined by $\underline{q}_C = \underline{q}^i_C$ if C is an irreducible component of X contained in Y_i. We say that $\underline{q}$ is the polarization *induced* by the polarizations $(\underline{q}^1, \ldots, \underline{q}^r)$. Note that $|\underline{q}| = \sum_i |\underline{q}^i|$.

Proposition 6.6. *Let X be a connected curve satisfying condition (†).*

(i) The pull-back map

$$\begin{aligned} \tau^* : \overline{\mathbb{J}}_X &\longrightarrow \prod_{i=1}^r \overline{\mathbb{J}}_{Y_i} \\ I &\mapsto (I_{|Y_1}, \ldots, I_{|Y_r}), \end{aligned}$$

is an isomorphism. Moreover $\tau^*(\mathbb{J}_X) = \prod_i \mathbb{J}_{Y_i}$.

(ii) Given a polarization $\underline{q}^i$ on each curve Y_i, consider the induced polarization $\underline{q} := (\underline{q}^1, \ldots, \underline{q}^r)$ on X as above. Then $\underline{q}$ is general if and only if each $\underline{q}^i$ is general and in this case the morphism τ^* induces an isomorphism

$$\tau^* : \overline{J}_X(\underline{q}) \xrightarrow{\cong} \prod_i \overline{J}_{Y_i}(\underline{q}^i). \tag{6.5}$$

(iii) If $\underline{q}$ is a general polarization on X then there exists a general polarization $\underline{q}'$ with $|\underline{q}'| = |\underline{q}|$ on X which is induced by some polarizations $\underline{q}^i$ on Y_i and such that

$$\overline{J}_X(\underline{q}) = \overline{J}_X(\underline{q}').$$

Proof. By induction on the number of separating points, it is enough to consider the case where there is only one separating point $n_1 = n$, i.e. $r = 2$. Therefore the normalization $\widetilde{X}$ of X at n is the disjoint union of two connected curves Y_1 and Y_2, which we also identify with two subcurves of X meeting at the node n. Denote by C_1 (resp. C_2) the irreducible component of Y_1 (resp. Y_2) that contains the separating point n. A warning about the notation: given a subcurve $Z \subset X$, we will denote by Z^c the complementary subcurve of Z inside X, i.e. $\overline{X \setminus Z}$, and we set $\delta_Z := |Z \cap Z^c|$. In the case where $Z \subset Y_i \subset X$ for some $i = 1, 2$ we will write $\overline{Y_i \setminus Z}$ for the complementary subcurve of Z inside Y_i and we denote by $|Z \cap \overline{Y_i \setminus Z}|$ the intersection of Z and $\overline{Y_i \setminus Z}$.

Part (i) is well-known, see [Est01, Example 37]. The crucial fact is that if I is simple then I must be locally free at the separating point n; hence $\tau^*(I)$ is still torsion-free, rank-1 and its restrictions $\tau^*(I)_{|Y_i} = I_{|Y_i}$ are torsion-free, rank-1 and simple. Moreover, since n is a separating point, the sheaf I is completely determined by its pull-back $\tau^*(I)$, i.e. there are no gluing conditions. Finally, I is a line bundle if and only if its pull-back $\tau^*(I)$ is a line bundle.

Part (ii). Observe first that, by the definition of $\underline{q}$, for any subcurve $Z \subset X$ we have

$$\underline{q}_Z = \underline{q}_{Z \cap Y_1} + \underline{q}_{Z \cap Y_2} = \underline{q}^1_{Z \cap Y_1} + \underline{q}^2_{Z \cap Y_2}. \tag{6.6}$$

The subcurves of Z (resp. of Y_i for $i=1,2$) such that Z and Z^c (resp. Z and $\overline{Y_i \setminus Z}$) are connected will play a special role in the sequel in virtue of Remark 2.7 and 2.9; let us examine them carefully. The subcurves $Z \subset X$ such that Z and Z^c are connected can be of three types:

$$\begin{cases} \underline{\text{Case I:}}\ C_1, C_2 \subset Z^c \Longrightarrow Z \subset Y_i \text{ and } \overline{Y_i \setminus Z} \text{ is connected (for some } i=1,2), \\ \underline{\text{Case II:}}\ C_1, C_2 \subset Z \Longrightarrow Z^c \subset Y_i \text{ and } \overline{Y_i \setminus Z^c} \text{ is connected (for some } i=1,2), \\ \underline{\text{Case III:}}\ C_i \subset Z \text{ and } C_{3-i} \subset Z^c \Longrightarrow Z = Y_i \text{ and } Z^c = Y_{3-i} \text{ (for some } i=1,2). \end{cases} \tag{6.7}$$

The number δ_Z is determined in each of the above cases as it follows:

$$\delta_Z = \begin{cases} |Z \cap \overline{Y_i \setminus Z}| & \text{in Case I,} \\ |Z^c \cap \overline{Y_i \setminus Z^c}| & \text{in Case II,} \\ 1 & \text{in Case III,} \end{cases} \tag{6.8}$$

On the other hand, the subcurves $Z \subset Y_i$ such that Z and $\overline{Y_i \setminus Z}$ are connected (for some $i=1,2$) can be of two types:

$$\begin{cases} \underline{\text{Case A:}}\ C_i \not\subset Z \Longrightarrow Z^c \text{ is connected,} \\ \underline{\text{Case B:}}\ C_i \subset Z \Longrightarrow (\overline{Y_i \setminus Z})^c \text{ is connected.} \end{cases} \tag{6.9}$$

The intersection number $|Z \cap \overline{Y_i \setminus Z}|$ is given in each of the above cases by:

$$|Z \cap \overline{Y_i \setminus Z}| = \begin{cases} \delta_Z & \text{in Case A,} \\ \delta_{(\overline{Y_i \setminus Z})^c} & \text{in Case B,} \end{cases} \tag{6.10}$$

After these preliminaries, we can now prove that $\underline{q}$ is general if and only if each $\underline{q}^i$ is general. Assume first that each $\underline{q}^i$ is a general polarization on Y_i for $i=1,2$. Consider a proper subcurve $Z \subset X$ such that Z and Z^c are connected. Using (6.6), (6.7) and (6.8), we compute

$$\underline{q}_Z - \frac{\delta_Z}{2} = \begin{cases} \underline{q}^i_Z - \dfrac{|Z \cap \overline{Y_i \setminus Z}|}{2} & \text{in case I,} \\ |\underline{q}| - \underline{q}^i_{Z^c} - \dfrac{|Z^c \cap \overline{Y_i \setminus Z^c}|}{2} & \text{in case II,} \\ |\underline{q}^i| - \dfrac{1}{2} & \text{in case III.} \end{cases}$$

In each of the cases I, II, III we conclude that $\underline{q}_Z - \frac{\delta_Z}{2} \notin \mathbb{Z}$ using that $\underline{q}^i$ is general and $|\underline{q}|, |\underline{q}^i| \in \mathbb{Z}$. Therefore $\underline{q}$ is general by Remark 2.7.

Conversely, assume that $\underline{q}$ is general and let us show that $\underline{q}^i$ is general for $i=1,2$. Consider a proper subcurve $Z \subset Y_i$ such that Z and $\overline{Y_i \setminus Z}$ is connected. Using (6.6), (6.9) and (6.10), we compute

$$\underline{q}^i_Z - \frac{|Z \cap \overline{Y_i \setminus Z}|}{2} = \begin{cases} \underline{q}_Z - \dfrac{\delta_Z}{2} & \text{in case A,} \\ |\underline{q}^i| - \underline{q}_{(\overline{Y^i \setminus Z})^c} - \dfrac{\delta_{(\overline{Y_i \setminus Z})^c}}{2} & \text{in case B.} \end{cases}$$

In each of the cases A, B we conclude that $\underline{q}^i_Z - \frac{|Z \cap \overline{Y_i \setminus Z}|}{2} \notin \mathbb{Z}$ using that $\underline{q}$ is general and $|\underline{q}^i| \in \mathbb{Z}$. Therefore $\underline{q}^i$ is general by Remark 2.7.

Finally, in order to prove (6.5), it is enough, using part (i), to show that a simple torsion-free rank-1 sheaf I on X has degree $|\underline{q}|$ and is $\underline{q}$-semistable if and only if $I_{|Y_i}$ has degree $|\underline{q}^i|$ and is $\underline{q}^i$-semistable for $i=1,2$. Observe first that, since I is locally free at the node n (see the proof of part (i)), we have that for any subcurve $Z \subset X$ it holds

$$\deg_Z(I) = \deg_{Z \cap Y_1}(I) + \deg_{Z \cap Y_2}(I) = \deg_{Z \cap Y_1}(I_{|Y_1}) + \deg_{Z \cap Y_2}(I_{|Y_2}). \tag{6.11}$$

Assume first that $I_{|Y_i}$ has degree $|\underline{q}^i|$ and is $\underline{q}^i$-semistable for $i=1,2$. Using (6.11), we get

$$\deg(I) = \deg_{Y_1}(I) + \deg_{Y_2}(I) = |\underline{q}^1| + |\underline{q}^2| = |\underline{q}|.$$

Consider a proper subcurve $Z \subset X$ such that Z and Z^c are connected. Using (6.6), (6.7), (6.8) and (6.11) we compute

$$\deg_Z(I)-\underline{q}_Z+\frac{\delta_Z}{2}=\begin{cases} \deg_Z(I_{|Y_i})-\underline{q}^i_Z+\dfrac{|Z\cap\overline{Y_i\setminus Z}|}{2} & \text{in case I,}\\ \deg(I_{|Y_{3-i}})+\deg_{\overline{Y_i\setminus Z^c}}(I_{|Y_i})-|\underline{q}^{3-i}|-\underline{q}^i_{\overline{Y_i\setminus Z^c}}+\dfrac{|Z^c\cap\overline{Y_i\setminus Z^c}|}{2}= & \\ =\deg_{\overline{Y_i\setminus Z^c}}(I_{|Y_i})-\underline{q}^i_{\overline{Y_i\setminus Z^c}}+\dfrac{|Z^c\cap \overline{Y_i\setminus Z^c}|}{2} & \text{in case II,}\\ \deg(I_{|Y_i})-|\underline{q}^i|+\dfrac12=\dfrac12 & \text{in case III.}\end{cases}$$

In each of the cases I, II, III we conclude that $\deg_Z(I)-\underline{q}_Z+\frac{\delta_Z}{2}\geq 0$ using that $I_{|Y_i}$ is $\underline{q}^i$-semistable. Therefore I is $\underline{q}$-semistable by Remark 2.9.

Conversely, assume that I has degree $|\underline{q}|$ and is $\underline{q}$-semistable. Using (6.11), we get that

$$|\underline{q}^1|+|\underline{q}^2|=|\underline{q}|=\deg(I)=\deg_X(I)=\deg_{Y_1}(I)+\deg_{Y_2}(I)=\deg(I_{|Y_1})+\deg(I_{|Y_2}). \tag{6.12}$$

Since I is $\underline{q}$-semistable, inequalities (2.10) applied to Y_i for $i=1,2$ gives that

$$\deg(I_{|Y_i})=\deg_{Y_i}(I)\geq \underline{q}_{Y_i}-\frac{\delta_{Y_i}}{2}=|\underline{q}^i|-\frac12. \tag{6.13}$$

Since $\deg(I_{|Y_i})$ and $|\underline{q}^i|$ are integral numbers, from (6.13) we get that $\deg(I_{|Y_i})\geq|\underline{q}^i|$ which combined with (6.12) gives that $\deg(I_{|Y_i})=|\underline{q}^i|$. Consider now a subcurve $Z\subset Y_i$ (for some $i=1,2$) such that Z and $\overline{Y_i\setminus Z}$ is connected. Using (6.6), (6.9), (6.10) and (6.11) we compute

$$\deg_Z(I_{|Y_i})-\underline{q}^i_Z+\frac{|Z\cap\overline{Y_i\setminus Z}|}{2}=\begin{cases}\deg_Z(I)-\underline{q}_Z+\dfrac{\delta_Z}{2} & \text{in case A,}\\ -\deg(I_{|Y_{3-i}})+\deg_{(\overline{Y_i\setminus Z})^c}(I)+|\underline{q}^{3-i}|-\underline{q}_{(\overline{Y_i\setminus Z})^c}+\dfrac{\delta_{(\overline{Y_i\setminus Z})^c}}{2}= & \\ \deg_{(\overline{Y_i\setminus Z})^c}(I)-\underline{q}_{(\overline{Y_i\setminus Z})^c}+\dfrac{\delta_{(\overline{Y_i\setminus Z})^c}}{2} & \text{in case B.}\end{cases}$$

In each of the cases A, B we conclude that $\deg_Z(I_{|Y_i})-\underline{q}^i_Z+\frac{|Z\cap\overline{Y_i\setminus Z}|}{2}\geq 0$ using that I is $\underline{q}$-semistable. Therefore $I_{|Y_i}$ is $\underline{q}^i$-semistable by Remark 2.9.

Part (iii): note that a polarization $\underline{q}'$ on X is induced by some polarizations $\underline{q}^i$ on Y_i if and only if $\underline{q}'_{Y_i}\in\mathbb{Z}$ for $i=1,2$. For a general polarization $\underline{q}$ on Y, we have that

$$\begin{cases}|\underline{q}|=\underline{q}_{Y_1}+\underline{q}_{Y_2}\in\mathbb{Z},\\ \underline{q}_{Y_i}-\dfrac{Y_1\cap Y_2}{2}=\underline{q}_{Y_i}-\dfrac12\notin\mathbb{Z}.\end{cases}$$

Therefore, we can find unique integral numbers $m_1,m_2\in\mathbb{Z}$ and a unique rational number $r\in\mathbb{Q}$ with $-\frac12<r<\frac12$ such that

$$\begin{cases}\underline{q}_{Y_1}=m_1+r,\\ \underline{q}_{Y_2}=m_2-r.\end{cases} \tag{6.14}$$

Define now the polarization $\underline{q}'$ on X in such a way that for an irreducible component C of X, we have that

$$\underline{q}'_C:=\begin{cases}\underline{q}_C & \text{if } C\neq C_1,C_2,\\ \underline{q}_{C_1}-r & \text{if } C=C_1,\\ \underline{q}_{C_2}+r & \text{if } C=C_2.\end{cases}$$

In particular for any subcurve $Z\subset X$, the polarization $\underline{q}'$ is such that

$$\underline{q}'_Z:=\begin{cases}\underline{q}_Z & \text{if either } C_1,C_2\subset Z \text{ or } C_1,C_2\subset Z^c,\\ \underline{q}_Z-r & \text{if } C_1\subset Z \text{ and } C_2\subset Z^c,\\ \underline{q}_Z+r & \text{if } C_2\subset Z \text{ and } C_1\subset Z^c.\end{cases} \tag{6.15}$$

Specializing to the case $Z = Y_1, Y_2$ and using (6.14), we get that

$$
\begin{cases}
\underline{q}'_{Y_1} = \underline{q}_{Y_1} - r = m_1 \in \mathbb{Z}, \\
\underline{q}'_{Y_2} = \underline{q}_{Y_2} + r = m_2 \in \mathbb{Z}, \\
|\underline{q}'| = \underline{q}'_{Y_1} + \underline{q}'_{Y_2} = m_1 + m_2 = \underline{q}_{Y_1} + \underline{q}_{Y_2} = |\underline{q}|.
\end{cases}
\tag{6.16}
$$

As observed before, this implies that $\underline{q}'$ is induced by two (uniquely determined) polarizations $\underline{q}^1$ and $\underline{q}^2$ on Y_1 and Y_2, respectively, and moreover that $|\underline{q}'| = |\underline{q}|$.

Let us check that $\underline{q}'$ is general. Consider a proper subcurve $Z \subset X$ such that Z and Z^c are connected. Using (6.7), (6.15) and (6.16), we compute that

$$
\underline{q}'_Z - \frac{\delta_Z}{2} =
\begin{cases}
\underline{q}_Z - \dfrac{\delta_Z}{2} & \text{in case I and II}, \\
\underline{q}'_{Y_i} - \dfrac{\delta_{Y_i}}{2} = m_i - \dfrac{1}{2} & \text{in case III}.
\end{cases}
\tag{6.17}
$$

In each of the above cases I, II, III we get that $\underline{q}'_Z - \frac{\delta_Z}{2} \notin \mathbb{Z}$ using that $\underline{q}$ is general and that $m_i \in \mathbb{Z}$. Therefore $\underline{q}'$ is general by Remark 2.7.

Finally, in order to check that $\overline{J}_X(\underline{q}) = \overline{J}_X(\underline{q}')$ we must show that a simple rank-1 torsion-free sheaf I of degree equal to $|\underline{q}| = |\underline{q}'|$ on X is $\underline{q}$-semistable if and only if it is $\underline{q}'$-semistable. Using Remark 2.9, it is sufficient (and necessary) to check that for, any proper subcurve $Z \subset X$ such that Z and Z^c are connected, I satisfies (2.10) with respect to $\underline{q}_Z$ if and only if it satisfies (2.10) with respect to $\underline{q}'_Z$. If Z belong to case I or II (according to the classification (6.7)), this is clear by (6.17). If Z belongs to case III, i.e. if $Z = Y_i$ for some $i = 1, 2$, then, using (6.14) together with the fact that $-\frac{1}{2} < r < \frac{1}{2}$ and $m_i, \deg_{Y_i}(I) \in \mathbb{Z}$, we get that

$$
\deg_{Y_i}(I) \geq \underline{q}_{Y_i} - \frac{\delta_{Y_i}}{2} = m_i + (-1)^{i+1} r - \frac{1}{2} > m_i - 1 \Leftrightarrow \deg_{Y_i}(I) \geq m_i.
$$

Similarly using (6.16), we get that

$$
\deg_{Y_i}(I) \geq \underline{q}'_{Y_i} - \frac{\delta_{Y_i}}{2} = m_i - \frac{1}{2} \Leftrightarrow \deg_{Y_i}(I) \geq m_i.
$$

This show that I satisfies (2.10) with respect to $\underline{q}_{Y_1}$ if and only if it satisfies (2.10) with respect to $\underline{q}'_{Y_1}$, which concludes our proof.

□

We can now define the Abel maps for X.

Proposition 6.7. *Let X be a connected curve satisfying condition (†) as above.*

(i) For any line bundle $L \in \operatorname{Pic}(X)$, there exists a unique morphism $A_L : X \to \overline{\mathbb{J}}_X$ such that for any $i = 1, \ldots, r$ it holds:

(a) the following diagram is commutative

$$
\begin{tikzcd}
Y_i \arrow[r, "A_{L_i}"] \arrow[d, hook] & \overline{\mathbb{J}}_{Y_i} & \\
X \arrow[r, "A_L"] & \overline{\mathbb{J}}_X \arrow[r, "\tau^*", "\cong"'] & \prod_j \overline{\mathbb{J}}_{Y_j} \arrow[lu, "\pi_i"']
\end{tikzcd}
$$

where π_i denotes the projection onto the i-th factor, τ^* is the isomorphism of Proposition 6.6(i) *and A_{L_i} is the L_i-twisted map of (6.4) for $L_i := L_{|Y_i}$.*

(b) The composition

$$
Y_i \hookrightarrow X \xrightarrow{A_L} \overline{\mathbb{J}}_X \underset{\cong}{\xrightarrow{\tau^*}} \prod_j \overline{\mathbb{J}}_{Y_j} \xrightarrow{\prod_{j \neq i} \pi_j} \prod_{j \neq i} \overline{\mathbb{J}}_{Y_j}
$$

is a constant map.

(ii) Let $\underline{q}^i$ be a general polarization on Y_i for any $1 \leq i \leq r$ and denote by $\underline{q}$ the induced (general) polarization on X. Then

$$
A_L(X) \subset \overline{J}_X(\underline{q}) \Leftrightarrow A_{L_i}(Y_i) \subset \overline{J}_{Y_i}(\underline{q}^i) \text{ for any } 1 \leq i \leq r.
$$

Proof. Part (i): assume that such a map A_L exists and let us prove its uniqueness. From (ia) and (ib) it follows that the composition

$$\widetilde{A_L}:\widetilde{X}=\coprod_i Y_i \xrightarrow{\tau} X \xrightarrow{A_L} \overline{\mathbb{J}}_X \xrightarrow{\tau^*} \prod_i \overline{\mathbb{J}}_{Y_i}$$

is such that for every $1\leq i\leq r$ and every $p\in Y_i$ it holds

$$(\widetilde{A_L})_{|Y_i}(p)=(M_1^i,\ldots,M_{i-1}^i,A_{L_i}(p),M_{i+1}^i,\ldots,M_r^i) \tag{6.18}$$

for some elements $M_j^i\in\overline{\mathbb{J}}_{Y_j}$ for $j\neq i$. Moreover, if we set $\tau^{-1}(n_k)=\{n_k^1,n_k^2\}$ then we must have that

$$\widetilde{A_L}(n_k^1)=\widetilde{A_L}(n_k^2) \text{ for any } 1\leq k\leq r-1. \tag{6.19}$$

Claim: There exist uniquely determined elements $M_j^i\in\overline{\mathbb{J}}_{Y_j}$ (for any $i\neq j$) such that the map $\widetilde{A_L}$ in (6.18) satisfies the conditions in (6.19).

The claim clearly implies the uniqueness of the map $\widetilde{A_L}$, hence the uniqueness of the map A_L. Moreover, the same claim also shows the existence of the map A_L with the desired properties: it is enough to define $\widetilde{A_L}$ via the formula (6.18) and notice that, since the conditions (6.19) are satisfied, then the map $\widetilde{A_L}$ descends to a map $A_L:X\to\overline{\mathbb{J}}_X$.

It remains therefore to prove the Claim. Our proof is by induction on r, the case $r=1$ being trivial. In order to avoid confusions, we will now denote by $Y_i'\subset X$ the image of the component $Y_i\subset\widetilde{X}$ via the morphism τ. Certainly one among the subcurves $Y_i'\subset X$, call it Y_1', is such that $X':=\bigcup_{i\geq 2}Y_i'\subset X$ is connected. This is equivalent to say that there is only one separating point, call it n_1, belonging to Y_1. Up to renaming the subcurves Y_i', we can assume that $n_1=Y_1'\cap Y_2'$. The connected curve X' certainly satisfies condition (†) and it has $r-2$ separating points. Therefore, by induction, there exist uniquely determined elements M_i^j with $i\neq j\geq 2$ such that the map defined by (6.18) with respect to the curve X' satisfies the conditions in (6.19) for $k\geq 2$. Now, from condition (6.19) applied to n_1 we deduce that

$$\begin{cases} M_1^2=\mathfrak{m}_{n_1^1}\otimes L_1,\\ M_2^1=\mathfrak{m}_{n_1^2}\otimes L_2,\\ M_k^1=M_k^2 \text{ for any } k\geq 3.\end{cases} \tag{*}$$

Moreover, by taking the first component of the equality that we get from condition (6.19) applied to the other nodes n_h with $h\geq 2$, we deduce that

$$M_1^h=M_1^2 \text{ for any } h\geq 3. \tag{**}$$

By combining (*) and (**), we see that the elements M_i^j with either $i=1$ or $j=1$ are uniquely determined by the elements M_h^k with $h,k\geq 2$ and this concludes the proof of the claim.

Part (ii) follows easily from the diagram in (ia) and the isomorphism (6.5). □

We call the map A_L of Proposition 6.7(i) the *(L-twisted) Abel map* of X. Note that special cases of the Abel map A_L in the presence of separating points have been considered before by Caporaso-Esteves in [CE07, Sec. 4] for nodal curves and Caporaso-Coelho-Esteves in [CCE08, Sec. 4 and 5] for Gorenstein curves.

We can extend Definition 6.2 to the case of curves satisfying condition (†) from (6.4).

Definition 6.8. Let X be a curve satisfying condition (†). We say that a fine compactified Jacobian $\overline{J}_X(\underline{q})$ of X *admits an Abel map* if there exists $L\in\operatorname{Pic}^{|\underline{q}|+1}(X)$ such that $\mathrm{Im A_L}\subseteq\overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$.

By combining Propositions 6.6 and 6.7, we can easily reduce the problem of the existence of an Abel map for a fine compactified Jacobian of X to the analogous question on the separating blocks of X.

Corollary 6.9. *Let X be a curve satisfying condition (†) with separating blocks $Y_1,\cdots Y_r$.*

- *(i) Let $\underline{q}$ be a general polarization of X and assume (without loss of generality by Proposition 6.6*(iii)*) which is induced by some general polarizations $\underline{q}^i$ on Y_i. Then $\overline{J}_X(\underline{q})$ admits an Abel map if and only if each $\overline{J}_{Y_i}(\underline{q}^i)$ admits an Abel map.*
- *(ii) For any $L\in\operatorname{Pic}^d(X)$ there exists a general polarization $\underline{q}$ on X of total degree $|\underline{q}|=d-1$ such that* $\mathrm{Im A_L}\subseteq\overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$.

Proof. Part (i) follows from Proposition 6.7(ii). Part (ii) follows from Proposition 6.7(ii) together with Lemma 6.1. □

Note that if $\overline{J}_X(\underline{q})$ is a fine compactified Jacobian of X and $L \in \mathrm{Pic}^{|\underline{q}|+1}(X)$ is such that $\mathrm{ImA_L} \subseteq \overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$, then the Abel map $A_L : X \to \overline{J}_X(\underline{q}) \subseteq \overline{\mathbb{J}}_X$ induces via pull-back a homomorphism

$$A_L^* : \mathrm{Pic}(\overline{J}_X(\underline{q})) \to \mathrm{Pic}(X) = \mathbb{J}_X, \tag{6.20}$$

which clearly sends $\mathrm{Pic}^o(\overline{J}_X(\underline{q}))$ into $\mathrm{Pic}^o(X) = J(X)$.

When is the Abel map A_L an embedding? The answer is provided by the following result, whose proof is identical to the proof of [CCE08, Thm. 6.3].

Fact 6.10 (Caporaso-Coelho-Esteves)**.** *Let X be a curve satisfying condition (†) and $L \in \mathrm{Pic}(X)$. The Abel map A_L is an embedding if and only if X does not have separating blocks isomorphic to $\mathbb{P}^1$.*

In general, the Abel map A_L contracts the separating blocks isomorphic to $\mathbb{P}^1$ into ordinary singularities with linearly independent tangent directions.

7. EXAMPLES: LOCALLY PLANAR CURVES OF ARITHMETIC GENUS 1

In this section, we are going to study fine compactified Jacobians and Abel maps for singular curves of arithmetic genus 1 with locally planar singularities. According to Fact 6.4, such a curve X satisfies the condition (†) and therefore, using Proposition 6.6 and Proposition 6.7, we can reduce the study of fine compactified Jacobians and Abel maps to the case where X does not have separating points (or equivalently separating nodes). Under this additional assumption, a classification is possible.

Fact 7.1. *Let X be a (reduced) connected singular curve without separating points, with locally planar singularities and $p_a(X) = 1$. Then X is one of the curves depicted in Figure 1 and called Kodaira curves.*

Proof. Since X has non separating points and $p_a(X) = 1$ then X has trivial canonical sheaf by [Est01, Example 39]. These curves were classified by Catanese in [Cat82, Prop. 1.18]. An inspection of the classification in loc. cit. reveals that the only such singular curves that have locally planar singularities are the ones depicted in Figure 1, i.e. the Kodaira curves. ☐

Note that the curves in Figure 1 are exactly the reduced fibers appearing in the well-known Kodaira's classification of singular fibers of minimal elliptic fibrations (see [BPV84, Chap. V, Sec. 7]). This explains why they are called Kodaira curves.



Type I

Type $I_n, n \geq 2$

Type II

Type III

Type IV

FIGURE 1. Kodaira curves.

Abel maps for Kodaira curves behave particularly well, due to the following result proved in [Est01, Example 39].

Fact 7.2 (Esteves)**.** *Let X be a connected curve without separating points and such that $p_a(X) = 1$. Then for any $L \in \mathrm{Pic}(X)$ the image $A_L(X) \subseteq \overline{\mathbb{J}}_X$ of X via the L-twisted Abel map is equal to a fine compactified Jacobian $\overline{J}_X(\underline{q})$ of X and A_L induces an isomorphism*

$$A_L : X \xrightarrow{\cong} A_L(X) = \overline{J}_X(\underline{q}).$$

From the above Fact 7.2, we deduce that, up to equivalence by translation (in the sense of Definition 2.13), there is exactly one fine compactified Jacobian that admits an Abel map and this fine compactified Jacobian is isomorphic to the curve itself. This last property is indeed true for any fine compactified Jacobian of a Kodaira curve, as shown in the following

Proposition 7.3. *Let X be a Kodaira curve. Then every fine compactified Jacobian of X is isomorphic to X.*

Proof. Let $\overline{J}_X(\underline{q})$ be a fine compactified Jacobian of X. By Lemma 4.8, we can find a 1-parameter regular smoothing $f : \mathcal{S} \to B = \operatorname{Spec} R$ of X (in the sense of Definition 4.7), where R is complete discrete valuation ring with quotient field K. Note that the generic fiber $\mathcal{S}_K$ of f is an elliptic curve. Following the notation of §4.1, we can form the f-relative fine compactified Jacobian $\pi : \overline{J}_f(\underline{q}) \to B$ with respect to the polarization $\underline{q}$. Recall that π is a projective and flat morphism whose generic fiber is $\operatorname{Pic}^{|\underline{q}|}(\mathcal{S}_K)$ and whose special fiber is $\overline{J}_X(\underline{q})$. Using Theorem 4.3, it is easy to show that if we choose a generic 1-parameter smoothing $f : \mathcal{S} \to B$ of X, then the surface $\overline{J}_f(\underline{q})$ is regular. Moreover, Fact 4.9 implies that the smooth locus $J_f(\underline{q}) \to B$ of π is isomorphic to the Néron model of the generic fiber $\operatorname{Pic}^{|\underline{q}|}(\mathcal{S}_K)$. Therefore, using the well-known relation between the Néron model and the regular minimal model of the elliptic curve $\operatorname{Pic}^{|\underline{q}|}(\mathcal{S}_K) \cong \mathcal{S}_K$ over K (see [BLR90, Chap. 1.5, Prop. 1]), we deduce that $\pi : \overline{J}_f(\underline{q}) \to B$ is the regular minimal model of $\operatorname{Pic}^{|\underline{q}|}(\mathcal{S}_K)$. In particular, π is a minimal elliptic fibration with reduced fibers and therefore, according to Kodaira's classification (see see [BPV84, Chap. V, Sec. 7]), the special fiber $\overline{J}_X(\underline{q})$ of π must be a smooth elliptic curve or a Kodaira curve.

According to Corollary 4.12, the number of irreducible components of $\overline{J}_X(\underline{q})$ is equal to the complexity $c(X)$ of X. However, it is very easy to see that for a Kodaira curve X the complexity number $c(X)$ is equal to the number of irreducible components of X. Therefore if $c(X) \geq 4$, i.e. if X is of Type I_n with $n \geq 4$, then necessarily $\overline{J}_X(\underline{q})$ is of type I_n, hence it is isomorphic to X.

In the case $n \leq 3$, the required isomorphism $\overline{J}_X(\underline{q}) \cong X$ follows from the fact that the smooth locus of $\overline{J}_X(\underline{q})$ is isomorphic to a disjoint union of torsors under $\operatorname{Pic}^{\underline{0}}(X)$ (see Corollary 2.12) and that

$$\operatorname{Pic}^{\underline{0}}(X) = \begin{cases} \mathbb{G}_m & \text{if } X \text{ is of Type } I \text{ or } I_n \ (n \geq 2), \\ \mathbb{G}_a & \text{if } X \text{ is of Type } II, III \text{ or } IV. \end{cases}$$

□

Let us now classify the fine compactified Jacobians for a Kodaira curve X, up to equivalence by translation, and indicate which of them admits an Abel map.

$\boxed{X \text{ is of Type } I \text{ or Type } II}$

Since the curve X is irreducible, we have that the fine compactified Jacobians of X are of the form $\overline{\mathbb{J}}_X^d$ for some $d \in \mathbb{Z}$. Hence they are all equivalent by translation and each of them admits an Abel map.

$\boxed{X \text{ is of Type } I_n, \text{ with } n \geq 2}$

The fine compactified Jacobians of X up to equivalence by translation and their behavior with respect to the Abel map are described in the following proposition.

Proposition 7.4. *Let X be a Kodaira curve of type I_n (with $n \geq 2$) and let $\{C_1, \ldots, C_n\}$ be the irreducible components of X, ordered in such a way that, for any $1 \leq i \leq n$, C_i intersects C_{i-1} and C_{i+1}, with the cyclic convention that $C_{n+1} := C_1$.*

(i) Any fine compactified Jacobian is equivalent by translation to a unique fine compactified Jacobian of the form $\overline{J}_X(\underline{q})$ for a polarization $\underline{q}$ that satisfies

$$(*) \qquad \underline{q} = \left(q_1, \ldots, q_{n-1}, -\sum_{i=1}^{n-1} q_i\right) \text{ with } 0 \leq q_i < 1,$$

$$(**) \qquad \sum_{i=r}^{s} q_i \notin \mathbb{Z} \text{ for any } 1 \leq r \leq s \leq n-1,$$

$$(***) \qquad q_i = \frac{k_i}{n} \text{ with } k_i = 1, \ldots, n-1, \text{ for any } 1 \leq i \leq n-1.$$

In particular, there are exactly $(n-1)!$ fine compactified Jacobians of X up to equivalence by translation.

(ii) The unique fine compactified Jacobian, up to equivalence by translation, that admits an Abel map is
$$\overline{J}_X\left(\frac{n-1}{n},\dots,\frac{n-1}{n},-\frac{(n-1)^2}{n}\right).$$

Proof. Part (i): given any polarization $\underline{q}'$, there exists a unique polarization $\underline{q}$ that satisfies conditions (*) and such that $\underline{q}-\underline{q}'\in\mathbb{Z}^n$. Since any connected subcurve $Y\subset X$ is such that Y or Y^c is equal to $C_r\cup\ldots\cup C_s$ (for some $1\le r\le s\le n-1$) and $\delta_Y=2$, we deduce that a polarization $\underline{q}$ that satisfies (*) is general if and only if it satisfies (**). Hence any fine compactified Jacobian is equivalent by translation to a unique $\overline{J}_X(\underline{q})$, for a polarization $\underline{q}$ that satisfies (*) and (**). Consider now the arrangement of hyperplanes in $\mathbb{R}^{n-1}$ given by
$$\left\{\sum_{i=r}^{s} q_i = n\right\}$$
for all $1\le r\le s\le n-1$ and all $n\in\mathbb{Z}$. This arrangement of hyperplanes cuts the hypercube $[0,1]^{n-1}$ into finitely many chambers. Notice that a polarization $\underline{q}$ satisfies (*) and (**) if and only if it belongs to the interior of one of these chambers. Arguing as in the proof of Lemma 2.14, it is easy to see that two polarizations $\underline{q}$ and $\underline{q}'$ satisfying (*) and (**) belong to the same chamber if and only if $\overline{J}_X(\underline{q})=\overline{J}_X(\underline{q}')$. Now it is an entertaining exercise (that we leave to the reader) to check that any chamber contains exactly one polarization $\underline{q}$ that satisfies (***). This proves the first claim of part (i). The second claim in part (i) is an easy counting argument that we again leave to the reader.

Part (ii): if we take a line bundle L of multidegree $\underline{\deg}L=(1,\dots,1,-(n-2))$ then from the proof of Lemma 6.1 it follows that $\mathrm{ImA_L}\subseteq\overline{\mathrm{J}}:=\overline{\mathrm{J}}_{\mathrm{X}}(\frac{\mathrm{n}-1}{\mathrm{n}},\dots,\frac{\mathrm{n}-1}{\mathrm{n}},-\frac{(\mathrm{n}-1)^2}{\mathrm{n}})$. Therefore, from Fact 7.2, it follows that $\overline{J}$ is the unique fine compactified Jacobian, up to equivalence by translation, that admits an Abel map.

□

$\boxed{X \text{ is of Type } III}$

Since X has two irreducible components, then every fine compactified Jacobian of X admits an Abel map by Remark 6.3. By Fact 7.2, all fine compactified Jacobians of X are therefore equivalent by translation.

$\boxed{X \text{ is of Type } IV}$

The fine compactified Jacobians of X up to equivalence by translation and their behavior with respect to the Abel map are described in the following proposition.

Proposition 7.5. *Let X be a Kodaira curve of type IV.*

(i) Any fine compactified Jacobian of X is equivalent by translation to either $\overline{J}_1:=\overline{J}_X\left(\frac{2}{3},\frac{2}{3},-\frac{4}{3}\right)$ or $\overline{J}_2:=\overline{J}_X\left(\frac{1}{3},\frac{1}{3},-\frac{2}{3}\right)$.

(ii) $\overline{J}_1$ admits an Abel map while $\overline{J}_2$ does not admit an Abel map.

Proof. Part (i) is proved exactly as in the case of the Kodaira curve of type I_3 (see Proposition 7.4(i)).

Part (ii): if we take a line bundle L of multidegree $\underline{\deg}L=(1,1,-1)$ then $\mathrm{ImA_L}\subseteq\overline{\mathrm{J}}_1$ as it follows from the proof of Lemma 6.1. Therefore $\overline{J}_1$ admits an Abel map.

Let us now show that $\overline{J}_2$ does not admit an Abel map. Suppose by contradiction that there exists a line bundle L of multidegree $\deg L=(d_1,d_2,d_3)$ such that $A_L(p)=\mathfrak{m}_p\otimes L\in\overline{J}_2$ where p denotes the unique singular point of X. The stability of $\mathfrak{m}_p\otimes L$ with respect to the polarization $(q_1,q_2,q_3)=(\frac{1}{3},\frac{1}{3},-\frac{2}{3})$ gives for any irreducible component C_i of X:
$$-1+d_1=\deg_{C_i}(\mathfrak{m}_p\otimes L)>q_i+\frac{\delta_{C_i}}{2}=q_i-1.$$
We deduce that $d_1\ge1$, $d_2\ge1$ and $d_3\ge0$. However if $\mathrm{ImA_L}\subset\overline{\mathrm{J}}_2$ then the total degree of L must be one, which contradicts the previous conditions.

□

Remark 7.6. Realize a Kodaira curve X of Type IV as the plane cubic of equation $y(x+y)(x-y)=0$. One can show that the singular point in $\overline{J}_1$ and in $\overline{J}_2$ correspond to two sheaves that are not locally isomorphic: the singular point of $\overline{J}_1$ is the sheaf $\mathcal{I}_1 := \mathfrak{m}_p \otimes L$ where $\mathfrak{m}_p$ is the ideal sheaf of the point p defined by (x, y) and L is any line bundle on X of multidegree $(1, 1, , -1)$; the singular point of $\overline{J}_2$ is the sheaf $\mathcal{I}_2 := \mathcal{I}_Z \otimes M$ where $\mathcal{I}_Z$ is the ideal sheaf of the length 2 subscheme defined by (x, y^2) and M is any line bundle on X of multidegree $(1, 1, 0)$.

Moreover, denote by $\widetilde{X}$ the seminormalization of X (explicitly $\widetilde{X}$ can be realized as the union of three lines in the projective space meeting in one point with linearly independent direction) and $\pi : \widetilde{X} \to X$ is the natural map. Using the Table 2 of [Kas12(b)] (where the unique singularity of $\widetilde{X}$ is called $\widetilde{D}_4$ and the unique singular point of X is classically called D_4), then it can be shown that, up to the tensorization with a suitable line bundle on X, $\mathcal{I}_2$ is the pushforward of the trivial line bundle on $\widetilde{X}$ while $\mathcal{I}_1$ is the pushforward of the canonical sheaf on $\widetilde{X}$, which is not a line bundle since $\widetilde{X}$ is not Gorenstein (see Example 6.5).

Remark 7.7. Simpson's compactified Jacobians (which are not necessarily fine) of Kodaira curves are studied by A. C. López Martín in [LM05, Sec. 5].

8. The Poincaré bundle

The aim of this section is to introduce the Poincaré line bundle for fine compactified Jacobians and to study its properties.

With this in mind, consider the triple product $X \times \overline{\mathbb{J}}_X \times \mathbb{J}_X$ and, for any $1 \le i < j \le 3$, denote by p_{ij} the projection onto the product of the i-th and j-th factors. Choose a universal sheaf $\mathcal{I}$ on $X \times \overline{\mathbb{J}}_X$ as in Fact 2.2(iii) and denote by $\mathcal{I}^0$ its restriction to $X \times \mathbb{J}_X \subseteq X \times \overline{\mathbb{J}}_X$. Consider the trivial family of curves

$$p_{23} : X \times \overline{\mathbb{J}}_X \times \mathbb{J}_X \to \overline{\mathbb{J}}_X \times \mathbb{J}_X$$

and form the line bundle on $\overline{\mathbb{J}}_X \times \mathbb{J}_X$, called the *Poincaré bundle*:

$$\mathcal{P} := \mathcal{D}_{p_{23}}(p_{12}^*\mathcal{I} \otimes p_{13}^*\mathcal{I}^0)^{-1} \otimes \mathcal{D}_{p_{23}}(p_{13}^*\mathcal{I}^0) \otimes \mathcal{D}_{p_{23}}(p_{12}^*\mathcal{I}) \tag{8.1}$$

where $\mathcal{D}_{p_{23}}$ denotes the determinant of cohomology with respect to the morphism p_{23}. For the basic properties of the determinant of cohomology, we refer to [KM76] (see also [Est01, Sec. 6.1] for a summary).

Let us now examine how $\mathcal{P}$ depends on the choice of the universal sheaf $\mathcal{I}$. According to Fact 2.2(iii), any universal sheaf $\widetilde{\mathcal{I}}$ on $X \times \overline{\mathbb{J}}_X$ is related to $\mathcal{I}$ via

$$\widetilde{\mathcal{I}} = \mathcal{I} \otimes \pi_2^*(N),$$

for some $N \in \operatorname{Pic}(\overline{\mathbb{J}}_X)$, where $\pi_2 : X \times \overline{\mathbb{J}}_X \to \overline{\mathbb{J}}_X$ denotes the projection onto the second factor. Accordingly, the restriction $\widetilde{\mathcal{I}}^0$ of $\widetilde{\mathcal{I}}$ to $X \times \mathbb{J}_X$ is given by $\widetilde{\mathcal{I}}^0 = \mathcal{I}^0 \otimes \pi_2^*(N^0)$, where $N^0 := N_{|\mathbb{J}_X}$ and $\pi_2 : X \times \mathbb{J}_X \to \mathbb{J}_X$ denotes still the second projection (with a slight abuse of notation). If we denote by p_i $(i = 1, 2)$ the projection of $\overline{\mathbb{J}}_X \times \mathbb{J}_X$ onto the i-th factor, then, using the standard properties of the determinant of cohomology (see [KM76]), we get that

$$\begin{cases} \mathcal{D}_{p_{23}}(p_{12}^*\widetilde{\mathcal{I}} \otimes p_{13}^*\widetilde{\mathcal{I}}^0) = \mathcal{D}_{p_{23}}(p_{12}^*\mathcal{I} \otimes p_{13}^*\mathcal{I}^0 \otimes p_{23}^*(p_1^*N \otimes p_2^*N^0)) \\ \qquad\qquad = \mathcal{D}_{p_{23}}(p_{12}^*\mathcal{I} \otimes p_{13}^*\mathcal{I}^0) \otimes (p_1^*N \otimes p_2^*N^0)^{\chi(p_{12}^*\mathcal{I} \otimes p_{13}^*\mathcal{I}^0)}, \\ \mathcal{D}_{p_{23}}(p_{13}^*\widetilde{\mathcal{I}}^0) = \mathcal{D}_{p_{23}}(p_{13}^*\mathcal{I}^0 \otimes p_{23}^*p_2^*N^0) = \mathcal{D}_{p_{23}}(p_{13}^*\mathcal{I}^0) \otimes (p_2^*N^0)^{\chi(p_{13}^*\mathcal{I}^0)}, \\ \mathcal{D}_{p_{23}}(p_{12}^*\widetilde{\mathcal{I}}) = \mathcal{D}_{p_{23}}(p_{12}^*\mathcal{I} \otimes p_{23}^*p_1^*N) = \mathcal{D}_{p_{23}}(p_{12}^*\mathcal{I}) \otimes (p_1^*N)^{\chi(p_{12}^*\mathcal{I})}, \end{cases} \tag{8.2}$$

where χ denotes the Euler characteristic on the fibers of p_{23}. If we denote by deg the degree on the fibers of p_{23}, then, since $\mathcal{I}^0$ is a line bundle, we get that

$$\chi(p_{12}^*\mathcal{I} \otimes p_{13}^*\mathcal{I}^0) = \chi(p_{12}^*\mathcal{I}) + \deg(p_{13}^*\mathcal{I}^0) = \chi(p_{13}^*\mathcal{I}^0) + \deg(p_{12}^*\mathcal{I}). \tag{8.3}$$

Denote now by $\widetilde{\mathcal{P}}$ the sheaf on $\overline{\mathbb{J}}_X \times \mathbb{J}_X$ obtained from formula (8.1) by replacing $\mathcal{I}$ with $\widetilde{\mathcal{I}}$. Combining (8.2) and (8.3), we get that

$$\widetilde{\mathcal{P}} = \mathcal{P} \otimes (p_1^*N)^{-\deg p_{13}^*(\mathcal{I}^0)} \otimes (p_2^*N^0)^{-\deg p_{12}^*(\mathcal{I})}. \tag{8.4}$$

In particular, the restriction of $\mathcal{P}$ to $\overline{\mathbb{J}}_X \times \mathbb{J}_X^0$ is uniquely determined up to pull-back of a line bundle on $\mathbb{J}_X^0$ via the map p_2. Therefore, for any $M \in \operatorname{Pic}^0(X)$, we get a well defined line bundle

$$\mathcal{P}_M := \mathcal{P}_{|\overline{\mathbb{J}}_X \times \{M\}} \in \mathcal{P}ic(\overline{\mathbb{J}}_X).$$

Explicitly, since the determinant of cohomology commutes with base change, using (8.1) we get that the line bundle $\mathcal{P}_M$ on $\overline{\mathbb{J}}_X$ is given by

$$\mathcal{P}_M = \mathcal{D}_{\pi_2}(\mathcal{I}\otimes \pi_1^*M)^{-1}\otimes \mathcal{D}_{\pi_2}(\pi_1^*M)\otimes \mathcal{D}_{\pi_2}(\mathcal{I}), \tag{8.5}$$

where as usual π_i denotes the projection of $X\times\overline{\mathbb{J}}_X$ onto the i-th factor (for $i=1,2$) and $\mathcal{I}$ is a universal sheaf on $X\times\overline{\mathbb{J}}_X$ as in Fact 2.2(iii).

For any general polarization $\underline{q}$ on X, using the universal property of $\operatorname{Pic}(\overline{J}_X(\underline{q}))$, we get an algebraic morphism

$$\begin{aligned} \widetilde{\beta}_{\underline{q}}: \mathbb{J}_X^0 = \operatorname{Pic}^0(X) &\longrightarrow \operatorname{Pic}(\overline{J}_X(\underline{q})),\\ M &\mapsto (\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}. \end{aligned} \tag{8.6}$$

Note that, from (8.5), it follows that $\widetilde{\beta}_{\underline{q}}(\mathcal{O}_X) = (\mathcal{P}_{\mathcal{O}_X})_{|\overline{J}_X(\underline{q})} = \mathcal{O}_{\overline{J}_X(\underline{q})}$. Therefore the morphism $\widetilde{\beta}_{\underline{q}}$ restricts to a morphism

$$\begin{aligned} \beta_{\underline{q}}: \operatorname{Pic}^o(X) = \operatorname{Pic}^{\underline{0}}(X) &\to \operatorname{Pic}^o(\overline{J}_X(\underline{q})),\\ M &\mapsto (\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}. \end{aligned} \tag{8.7}$$

Proposition 8.1. *For any general polarization $\underline{q}$ on X, the maps $\widetilde{\beta}_{\underline{q}}$ and $\beta_{\underline{q}}$ are homomorphisms of group schemes.*

Proof. Since we have already observed that the maps in question are algebraic morphisms, it remains to prove that

$$\mathcal{P}_{M_1\otimes M_2} = \mathcal{P}_{M_1}\otimes \mathcal{P}_{M_2}. \tag{8.8}$$

for any $M_1, M_2\in \operatorname{Pic}^0(X)$.

In order to prove this, observe that we can write (for $i=1,2$) $M_i=\mathcal{O}_X(-\gamma_i+\delta_i)$, where γ_i and δ_i are effective divisors contained in the smooth locus of X. Moreover, we can clearly assume that δ_1 and δ_2 (resp. γ_1 and γ_2) have disjoint support.

Consider the following two exact sequences that are derived from the two exact sequences defining $\mathcal{O}_{\gamma_i}$ and $\mathcal{O}_{\delta_i}$:

$$\begin{aligned} &0\to \mathcal{O}_X(-\gamma_i)\to \mathcal{O}_X\to \mathcal{O}_{\gamma_i}\to 0,\\ &0\to \mathcal{O}_X(-\gamma_i)\to M_i\to M_{i|\delta_i}=\mathcal{O}_{\delta_i}\to 0. \end{aligned} \tag{8.9}$$

Pulling back (8.9) via π_1 and using the additivity of the determinant of cohomology, we get

$$\mathcal{D}_{\pi_2}(\pi_1^*M_i)\cong \mathcal{D}_{\pi_2}(\pi_1^*M_i)\otimes\mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_X)^{-1}\cong \mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_{\delta_i})\otimes \mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_{\gamma_i})^{-1}. \tag{8.10}$$

Similarly, by tensoring the pull-back via π_1 of two exact sequences (8.9) with $\mathcal{I}$ and using the additivity of determinant of cohomology, we get

$$\mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*M_i)^{-1}\otimes \mathcal{D}_{\pi_2}(\mathcal{I})\cong \mathcal{D}_{\pi_2}(\mathcal{I}\otimes \pi_1^*\mathcal{O}_{\delta_i})^{-1}\otimes\mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*\mathcal{O}_{\gamma_i}). \tag{8.11}$$

By plugging (8.10) and (8.11) into the (8.5), we get

$$\mathcal{P}_{M_i} = \mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*\mathcal{O}_{\delta_i})^{-1}\otimes \mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*\mathcal{O}_{\gamma_i})\otimes \mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_{\delta_i})\otimes \mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_{\gamma_i})^{-1}. \tag{8.12}$$

Since $M_1\otimes M_2 = \mathcal{O}_X(-\gamma_1-\gamma_2+\delta_1+\delta_2)$, we get in a similar way that

$$\mathcal{P}_{M_1\otimes M_2} = \mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*\mathcal{O}_{\delta_1\cup\delta_2})^{-1}\otimes \mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*\mathcal{O}_{\gamma_1\cup\gamma_2})\otimes\mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_{\delta_1\cup\delta_2})\otimes \mathcal{D}_{\pi_2}(\pi_1^*\mathcal{O}_{\gamma_1\cup\gamma_2})^{-1}. \tag{8.13}$$

Since δ_1 and δ_2 (reps. γ_1 and γ_2) are zero-dimensional subschemes of X with disjoint support, for any coherent sheaf $\mathcal{F}$ on $X\times\overline{\mathbb{J}}_X$ we have that

$$\begin{aligned} \mathcal{D}_{\pi_2}(\mathcal{F}\otimes\pi_1^*\mathcal{O}_{\delta_1\cup\delta_2}) &= \mathcal{D}_{\pi_2}(\mathcal{F}\otimes\pi_1^*\mathcal{O}_{\delta_1})\otimes\mathcal{D}_{\pi_2}(\mathcal{F}\otimes\pi_1^*\mathcal{O}_{\delta_2}),\\ \mathcal{D}_{\pi_2}(\mathcal{F}\otimes\pi_1^*\mathcal{O}_{\gamma_1\cup\gamma_2}) &= \mathcal{D}_{\pi_2}(\mathcal{F}\otimes\pi_1^*\mathcal{O}_{\gamma_1})\otimes\mathcal{D}_{\pi_2}(\mathcal{F}\otimes\pi_1^*\mathcal{O}_{\gamma_2}). \end{aligned} \tag{8.14}$$

Comparing formulas (8.12) and (8.13) and using (8.14), we get the required formula (8.8). □

Remark 8.2. With an argument similar to the one in the proof of the above Proposition 8.1, it is possible to get a description of the fibers of the Poincaré bundle. More precisely, given $(I,M)\in\overline{\mathbb{J}}_X\times\mathbb{J}_X$ with $M\cong\mathcal{O}_X(\sum a_iQ_i)$ for a divisor $\sum a_iQ_i$ supported on the smooth locus of X, then the fiber of $\mathcal{P}$ over (I,M) is canonically isomorphic to

$$\mathcal{P}_{(I,M)}\cong\bigotimes (I_{Q_i}^{-1}\otimes\mathcal{O}_{Q_i})^{\otimes a_i}. \tag{8.15}$$

An important property of the line bundles $\mathcal{P}_M$ on $\overline{\mathbb{J}}_X$ is the fact that they are invariant under pull-back for the multiplication map by an element $N \in \mathrm{Pic}(X)$:

$$\begin{aligned} -\otimes N : \overline{\mathbb{J}}_X &\to \overline{\mathbb{J}}_X \\ I &\mapsto I \otimes N. \end{aligned} \tag{8.16}$$

Lemma 8.3. *For any $N \in \mathrm{Pic}(X)$ and $M \in \mathrm{Pic}^0(X)$, we have that*

$$(-\otimes N)^* \mathcal{P}_M = \mathcal{P}_M.$$

Proof. Consider the following commutative diagram

$$\begin{tikzcd} X \times \overline{\mathbb{J}}_X \arrow[r, "(\mathrm{id}, -\otimes \mathrm{N})"] \arrow[d, "\pi_2"'] & X \times \overline{\mathbb{J}}_X \arrow[d, "\pi_2"] \\ \overline{\mathbb{J}}_X \arrow[r, "-\otimes N"'] & \overline{\mathbb{J}}_X \end{tikzcd} \tag{8.17}$$

By definition of the multiplication map $-\otimes N$, it follows that

$$(\mathrm{id}, -\otimes \mathrm{N})^* \mathcal{I} = \mathcal{I} \otimes \pi_1^* \mathrm{N}. \tag{8.18}$$

Using (8.5) and (8.18), together with the fact that the determinant of cohomology commutes with pull-back, we get

$$(-\otimes N)^* \mathcal{P}_M = \mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* N \otimes \pi_1^* M)^{-1} \otimes \mathcal{D}_{\pi_2}(\pi_1^* M) \otimes \mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* N). \tag{8.19}$$

By comparing (8.5) with (8.19), we deduce that the statement of the Lemma is equivalent to

$$\mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* M)^{-1} \otimes \mathcal{D}_{\pi_2}(\mathcal{I}) \cong \mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* N \otimes \pi_1^* M)^{-1} \otimes \mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* N). \tag{8.20}$$

In order to prove (8.20), write $M \cong \mathcal{O}_X(-\gamma + \delta)$ where γ and δ are two effective divisors on X_{sm}. By pulling back via π_1 the two exact sequences associated to the two effective divisors $\gamma, \delta \subset X$ and tensoring with a suitable line bundle on $X \times \overline{\mathbb{J}}_X$, we get the two exact sequences on $X \times \overline{\mathbb{J}}_X$:

$$\begin{cases} 0 \to \mathcal{I} \otimes \pi_1^* \mathcal{O}(-\gamma) \to \mathcal{I} \to \mathcal{I}_{|\pi_1^{-1}(\gamma)} \to 0, \\ 0 \to \mathcal{I} \otimes \pi_1^* \mathcal{O}(-\gamma) \to \mathcal{I} \otimes \pi_1^* M \to \mathcal{I}_{|\pi_1^{-1}(\delta)} \to 0, \end{cases} \tag{8.21}$$

From (8.21) and the additivity of the determinant of cohomology, we get that

$$\mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* M)^{-1} \otimes \mathcal{D}_{\pi_2}(\mathcal{I}) \cong \mathcal{D}_{\pi_2}(\mathcal{I}_{|\pi_1^{-1}(\delta)})^{-1} \otimes \mathcal{D}_{\pi_2}(\mathcal{I}_{|\pi_1^{-1}(\gamma)}). \tag{8.22}$$

Similarly, by tensoring the two exact sequences (8.21) with $\pi_1^*(N)$ and using the additivity of the determinant of cohomology together with the fact that $(\mathcal{I} \otimes \pi_1^* N)_{|\pi_1^{-1}(\delta)} \cong \mathcal{I}_{|\pi_1^{-1}(\delta)}$ and similarly with δ replaced by γ, we get that

$$\mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* M \otimes \pi_1^* N)^{-1} \otimes \mathcal{D}_{\pi_2}(\mathcal{I} \otimes \pi_1^* N) \cong \mathcal{D}_{\pi_2}(\mathcal{I}_{|\pi_1^{-1}(\delta)})^{-1} \otimes \mathcal{D}_{\pi_2}(\mathcal{I}_{|\pi_1^{-1}(\gamma)}). \tag{8.23}$$

By putting together (8.22) and (8.23), we get the isomorphism in (8.20), q.e.d. □

The Poincaré bundle behaves well with respect to the decomposition of a curve into its separating blocks in the sense of §6.2.

Lemma 8.4. *Let X be a curve satisfying condition (†) from (6.3) and denote by $Y_1, \ldots, Y_r$ its separating blocks. Let $M \in J(X) = \mathrm{Pic}^{\underline{0}}(X)$ and set $M_j := M_{|Y_j} \in J(Y_j) = \mathrm{Pic}^{\underline{0}}(Y_j)$. Denote by $\mathcal{P}_M \in \mathrm{Pic}^o(\overline{\mathbb{J}}_X)$ and $\mathcal{P}_{M_j} \in \mathrm{Pic}^o(\overline{\mathbb{J}}_{Y_j})$ the corresponding fibers of the Poincaré bundles for the curves X and Y_j, respectively. Then the push-forward of $\mathcal{P}_M$ via the isomorphism $\tau^* : \overline{\mathbb{J}}_X \xrightarrow{\cong} \prod_j \overline{\mathbb{J}}_{Y_j}$ of Proposition 6.6*(i) *is equal to*

$$\mathcal{P}_{M_1} \boxtimes \ldots \boxtimes \mathcal{P}_{M_r} := p_1^*(\mathcal{P}_{M_1}) \otimes \ldots \otimes p_r^*(\mathcal{P}_{M_r}),$$

where $p_j : \prod_i \overline{\mathbb{J}}_{Y_i} \to \overline{\mathbb{J}}_{Y_j}$ is the projection onto the j-th factor.

Proof. Consider the following commutative diagram

$$\begin{tikzcd}
\mathcal{I} \arrow[d, dash] & \widetilde{\mathcal{I}} := (\tau\times(\tau^*)^{-1})^*(\mathcal{I}) \arrow[d, dash] & \widetilde{\mathcal{I}_j} := (\mathrm{id}\times \mathrm{p_j})^*(\mathcal{I}_\mathrm{j}) \arrow[d, dash] & \mathcal{I}_j \arrow[d, dash] \\
X\times\overline{\mathbb{J}}_X \arrow[d, "\pi_2"] & \widetilde{X}\times\prod_i\overline{\mathbb{J}}_{Y_i} \arrow[l, "\tau\times(\tau^*)^{-1}"'] \arrow[d, "\widetilde{\pi_2}"] & Y_j\times\prod_i\overline{\mathbb{J}}_{Y_i} \arrow[l, hook', "\eta_j\times\mathrm{id}"'] \arrow[r, "\mathrm{id}\times\mathrm{p_j}"] \arrow[d, "\widetilde{\pi_2}^j"] \arrow[dr, phantom, "\square"] & Y_j\times\overline{\mathbb{J}}_{Y_j} \arrow[d, "\pi_2^j"] \\
\overline{\mathbb{J}}_X & \prod_i\overline{\mathbb{J}}_{Y_i} \arrow[l, "\cong"', "(\tau^*)^{-1}"] & \prod_i\overline{\mathbb{J}}_{Y_i} \arrow[l, equal] \arrow[r, "p_j"'] & \overline{\mathbb{J}}_{Y_j}
\end{tikzcd} \tag{8.24}$$

where $\mathcal{I}$ (resp. $\mathcal{I}_j$) is a universal sheaf for $\overline{\mathbb{J}}_X$ (resp. $\overline{\mathbb{J}}_{Y_j}$) as in Fact 2.2(iii), $\tau : \widetilde{X} = \coprod_i Y_i \to X$ is the normalization of X at the separating nodes of X (see (6.4)), $\eta_j : Y_j \hookrightarrow \widetilde{X} = \coprod_i Y_i$ is the natural inclusion. Denote by π_1, $\widetilde{\pi_1}$, $\widetilde{\pi_1}^j$, π_1^j the projections onto the first factors of the products appearing in the middle row of diagram (8.24).

Since a torsion-free rank-1 sheaf on X is completely determined by its pull-back to $\widetilde{X}$ by Proposition 6.6(i), we have that the pull-back of $\mathcal{P}_M$ to $\prod_i\overline{\mathbb{J}}_{Y_i}$ via the isomorphism $(\tau^*)^{-1}$ is equal to

$$\begin{aligned}((\tau^*)^{-1})^*(\mathcal{P}_M) &= ((\tau^*)^{-1})^*\left(\mathcal{D}_{\pi_2}(\mathcal{I}\otimes\pi_1^*M_j)^{-1}\otimes\mathcal{D}_{\pi_2}(\pi_1^*M_j)\otimes\mathcal{D}_{\pi_2}(\mathcal{I})\right) = \\ &= \mathcal{D}_{\widetilde{\pi_2}}(\widetilde{\mathcal{I}}\otimes\widetilde{\pi_1}^*M)^{-1}\otimes\mathcal{D}_{\widetilde{\pi_2}}(\widetilde{\pi_1}^*M)\otimes\mathcal{D}_{\widetilde{\pi_2}}(\widetilde{\mathcal{I}}).\end{aligned} \tag{*}$$

Since X is the disjoint union of the subcurves Y_i and $(\eta_j\times\mathrm{id})^*(\widetilde{\mathcal{I}}) = \widetilde{\mathcal{I}}_\mathrm{j}$, we have that

$$\begin{aligned}&\mathcal{D}_{\widetilde{\pi_2}}(\widetilde{\mathcal{I}}\otimes\widetilde{\pi_1}^*M)^{-1}\otimes\mathcal{D}_{\widetilde{\pi_2}}(\widetilde{\pi_1}^*M)\otimes\mathcal{D}_{\widetilde{\pi_2}}(\widetilde{\mathcal{I}}) = \\ &\bigotimes_{j=1}^r\left[\mathcal{D}_{\widetilde{\pi_2}^j}(\widetilde{\mathcal{I}_j}\otimes(\widetilde{\pi_1}^j)^*M_j)^{-1}\otimes\mathcal{D}_{\widetilde{\pi_2}^j}((\widetilde{\pi_1}^j)^*M_j)\otimes\mathcal{D}_{\widetilde{\pi_2}^j}(\widetilde{\mathcal{I}_j})\right].\end{aligned} \tag{**}$$

Finally, since the square in the right of diagram (8.24) is cartesian, applying the base change properties of the determinant of cohomology, we get

$$\begin{aligned}p_j^*(\mathcal{P}_{M_j}) &= p_j^*\left(\mathcal{D}_{\pi_2{}^j}(\mathcal{I}_j\otimes(\pi_1{}^j)^*M_j)^{-1}\otimes\mathcal{D}_{\pi_2{}^j}((\pi_1{}^j)^*M_j)\otimes\mathcal{D}_{\pi_2{}^j}(\mathcal{I}_j)\right) = \\ &= \mathcal{D}_{\widetilde{\pi_2}^j}(\widetilde{\mathcal{I}_j}\otimes(\widetilde{\pi_1}^j)^*M_j)^{-1}\otimes\mathcal{D}_{\widetilde{\pi_2}^j}((\widetilde{\pi_1}^j)^*M_j)\otimes\mathcal{D}_{\widetilde{\pi_2}^j}(\widetilde{\mathcal{I}_j}).\end{aligned} \tag{***}$$

By combining (*), (**) and (***) we get the equality

$$((\tau^*)^{-1})^*(\mathcal{P}_M) = \bigotimes_{j=1}^r p_j^*(\mathcal{P}_{M_j}),$$

which concludes the proof. ☐

Given any Abel map A_L and choosing a fine compactified Jacobian $\overline{J}_X(\underline{q})$ such that $\mathrm{ImA_L} \subseteq \overline{\mathrm{J}}_\mathrm{X}(\underline{\mathrm{q}})$ (which is always possible due to Corollary 6.9(ii)), the morphism $\beta_{\underline{q}}$ of (8.7) provides a right inverse for the morphism A_L^* of (6.20). This is originally due to Esteves-Gagné-Kleiman in the case where X is integral (see [EGK02, Prop. 2.2]).

Proposition 8.5. *Let X be a curve satisfying condition (†). Then, for every $L \in \mathrm{Pic}(X)$ and any general polarization $\underline{q}$ such that $\mathrm{ImA_L} \subseteq \overline{\mathrm{J}}_\mathrm{X}(\underline{\mathrm{q}})$, we have that*

$$A_L^*\circ\beta_{\underline{q}} = \mathrm{id}_{J(X)}.$$

In other words, for every $M \in J(X) = \mathrm{Pic}^{\underline{0}}(X)$ we have that $A_L^*((\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}) = M$.

Proof. We will first prove the Proposition in the case where X does not have separating points and then in the general case.

Case I: X does not have separating points.

The proof in this case is an easy adaptation of [EGK02, Prop. 2.2]. We include the proof for the reader's convenience.

Clearly, the composition $A_L^*\circ\beta_{\underline{q}} : J(X) \to J(X)$ is the morphism induced by the line bundle $(A_L\times\mathrm{id})^*\mathcal{P}$ on $X\times J(X)$. Since the determinant of cohomology commutes with base change, we get from (8.1) that

$$(A_L\times\mathrm{id})^*\mathcal{P} = \mathcal{D}_{p_{23}}(p_{12}^*(\mathrm{id}\times A_L)^*\mathcal{I}\otimes p_{13}^*\mathcal{I}^0)^{-1}\otimes\mathcal{D}_{p_{23}}(p_{13}^*\mathcal{I}^0)\otimes\mathcal{D}_{p_{23}}(p_{12}^*(\mathrm{id}\times A_L)^*\mathcal{I}), \tag{8.25}$$

where we denote, as usual, by p_{ij} the projection of $X \times X \times J(X)$ onto the product of the i-th and j-th factor.

Note that the Abel map $A_L : X \to \overline{\mathbb{J}}_X$ is defined by the sheaf $(\mathrm{id} \times \mathrm{A_L})^*(\mathcal{I})$ on $X \times X$ because of the universal property of $\mathcal{I}$ (see Fact 2.2(iii)) as well as by the sheaf $\mathcal{I}_\Delta \otimes p_1^* L$ by the definition of A_L (see (6.1)). Therefore, the two sheaves $(\mathrm{id} \times \mathrm{A_L})^*(\mathcal{I})$ and $\mathcal{I}_\Delta \otimes p_1^* L$ differ by the pull-back of a line bundle on X via the second projection $p_2 : X \times X \to X$. With the same computations that lead to (8.4), it is easy to prove that, since we are restricting ourselves to $\mathrm{Pic}^{\underline{0}}(X) = J(X)$, we can replace in (8.25) the sheaf $(\mathrm{id} \times \mathrm{A_L})^*(\mathcal{I})$ with the sheaf $\mathcal{I}_\Delta \otimes p_1^* L$; hence we get

$$(A_L \times \mathrm{id})^* \mathcal{P} = \mathcal{D}_{p_{23}}(p_{12}^* \mathcal{I}_\Delta \otimes p_1^* L \otimes p_{13}^* \mathcal{I}^0)^{-1} \otimes \mathcal{D}_{p_{23}}(p_{13}^* \mathcal{I}^0) \otimes \mathcal{D}_{p_{23}}(p_{12}^* \mathcal{I}_\Delta \otimes p_1^* L), \tag{8.26}$$

In order to evaluate the right hand side of (8.26), consider the exact sequence defining the diagonal $\Delta \subset X \times X$:

$$0 \to \mathcal{I}_\Delta \to \mathcal{O}_{X \times X} \to \mathcal{O}_\Delta \to 0. \tag{8.27}$$

Pulling back the sequence (8.27) to $X \times X \times J(X)$, tensoring it either with $p_1^* L \otimes p_{13}^* \mathcal{I}^0$ or with $p_1^* L$ and using the additivity of the determinant of cohomology, we get

$$\begin{cases} \mathcal{D}_{p_{23}}(p_{12}^* \mathcal{I}_\Delta \otimes p_1^* L \otimes p_{13}^* \mathcal{I}^0) = \mathcal{D}_{p_{23}}(p_1^* L \otimes p_{13}^* \mathcal{I}^0) \otimes (p_1^* L \otimes \mathcal{I}^0)^{-1}, \\ \mathcal{D}_{p_{23}}(p_{12}^* \mathcal{I}_\Delta \otimes p_1^* L) = \mathcal{D}_{p_{23}}(p_1^* L) \otimes (p_1^* L)^{-1}. \end{cases} \tag{8.28}$$

Consider the following Cartesian diagram

$$\begin{tikzcd}
X \times X \times J(X) \arrow[r, "p_{13}"] \arrow[d, "p_{23}"'] \arrow[dr, phantom, "\square"] & X \times J(X) \arrow[d, "p_2"] \\
X \times J(X) \arrow[r, "p_2"'] & J(X).
\end{tikzcd}$$

Since forming the determinant commutes with base change, we get that

$$\begin{cases} \mathcal{D}_{p_{23}}(p_1^* L \otimes p_{13}^* \mathcal{I}^0) = p_2^* \mathcal{D}_{p_2}(p_1^* L \otimes \mathcal{I}^0), \\ \mathcal{D}_{p_{23}}(p_1^* L) = p_2^* \mathcal{D}_{p_2}(p_1^* L), \\ \mathcal{D}_{p_{23}}(p_{13}^* \mathcal{I}^0) = p_2^* \mathcal{D}_{p_2}(\mathcal{I}^0). \end{cases} \tag{8.29}$$

Substituting (8.28) and (8.29) into (8.26), we get that $(A_L \times \mathrm{id})^* \mathcal{P}$ differs from $\mathcal{I}^0$ by tensor product with the pull-back of a line bundle on $J(X)$ via the projection map p_2. Since $\mathcal{I}^0$ is the universal sheaf above $X \times J(X)$, the map $J(X) = \mathrm{Pic}^{\underline{0}}(X) \to \mathrm{Pic}(X)$ induced by $\mathcal{I}^0$ is the identity. The same is therefore true for the map induced by $(A_L \times \mathrm{id})^* \mathcal{P}$, which is the map $A_L^* \circ \beta_{\underline{q}}$ as observed earlier, q.e.d.

Case II: X satisfies condition (†).

Let Y_i for $1 \le i \le r$ be the separating blocks of X as in §6.2 and set $L_i := L_{|Y_i} \in \mathrm{Pic}(Y_i)$. According to Proposition 6.6, we can assume that the general polarization $\underline{q}$ is induced by the general polarizations $\underline{q}^i$ on Y_i, for $1 \le i \le r$. Proposition 6.7(ii), together with our hypothesis on $\underline{q}$, ensures that $\mathrm{ImA_{L_i}} \subseteq \overline{\mathrm{J}}_{\mathrm{Y_i}}(\underline{\mathrm{q}}^{\mathrm{i}}) \subseteq \overline{\mathbb{J}}_{\mathrm{Y_i}}$ for every $1 \le i \le r$. Using Proposition 6.6(ii), we get the following diagram

$$\begin{tikzcd}
& \prod_i \mathrm{Pic}^o(\overline{J}_{Y_i}(\underline{q}^i)) \arrow[dl, bend right, "\prod_i A^*_{L_i}"'] \arrow[dr, "\otimes_i p_i^*(-)"] & \\
\prod_i J(Y_i) \arrow[ur, bend right, "\prod_i \beta_{\underline{q}_i}"'] & & \mathrm{Pic}^o(\prod_i \overline{J}_{Y_i}(\underline{q}^i)) \arrow[d, "\cong"', "\widehat{\tau^*}"] \\
J(X) \arrow[u, "\tau^*", "\cong"'] \arrow[rr, bend right, "\beta_{\underline{q}}"'] & & \mathrm{Pic}^o(\overline{J}_X(\underline{q})), \arrow[ll, bend right, "A_L^*"']
\end{tikzcd} \tag{8.30}$$

where $\widehat{\tau^*}$ is the map induced on Pic^o by the map τ^* of Proposition 6.6(ii) and $\beta_{\underline{q}_i}$ is the map (8.7) with respect to the general polarization $\underline{q}^i$ on the curve Y_i. Proposition 6.7(i) implies that the two maps from $\prod_i \mathrm{Pic}^o(\overline{\mathbb{J}}_{Y_i})$ to $\prod_i J(Y_i)$ that arise from diagram (8.30) are equal. Lemma 8.4 can be re-interpreted as saying that the two maps from $J(X)$ to $\mathrm{Pic}^o(\overline{\mathbb{J}}_X)$ that arise from diagram (8.30) are equal.

Since each Y_i does not have separating points, then we have that $A_{L_i}^* \circ \beta_{\underline{q}_i} = \mathrm{id_{J(Y_i)}}$ by Case I. This implies that $A_L^* \circ \beta_{\underline{q}} = \mathrm{id_{J(X)}}$ by an easy diagram chase in (8.30). ☐

An immediate consequence of the above result is the following:

Corollary 8.6. *Let X be a curve satisfying condition (†) and let $\underline{q}$ be a general polarization on X. If $\overline{J}_X(\underline{q})$ admits an Abel map (in the sense of Definition 6.8), then the homomorphism $\beta_{\underline{q}} : \mathrm{Pic}^{\underline{0}}(X) \to \mathrm{Pic}^o(\overline{J}_X(\underline{q}))$ is injective.*

9. Cohomology of restricted Poincaré bundles

The aim of this section is to prove some results about the cohomology of restricted Poincaré bundles $\mathcal{P}_M := \mathcal{P}_{|\mathbb{J}_X \times \{M\}}$ (with $M \in \mathrm{Pic}^{\underline{0}}(X)$) to the fine compactified Jacobians of X.

The first result is a generalization [Ari11, Prop. 1], which deals with X integral.

Proposition 9.1. *Let $M \in \mathrm{Pic}^{\underline{0}}(X) = J(X)$ and let $\underline{q}$ be a general polarization on X. If $H^i(\overline{J}_X(\underline{q}), \mathcal{P}_M) \neq 0$ for some i, then $\mathcal{P}_{M|\mathbb{J}_X} = \mathcal{O}_{\mathbb{J}_X}$.*

Proof. The proof is an adaptation of the proof of [Ari11, Prop. 1]. However, for the benefit of the reader, we chose to give some more details than in loc. cit.

Note that the generalized Jacobian $J(X) = \mathrm{Pic}^{\underline{0}}(X) = \mathbb{J}_X^{\underline{0}}$ acts on $\overline{J}_X(\underline{q})$. Denote by $T \to J(X)$ the $\mathbb{G}_m$-torsor corresponding to the line bundle $(\mathcal{P}_M)_{|J(X)}$. Then we have that:

Claim 1: T is a commutative algebraic group that is an extension of $J(X)$ by $\mathbb{G}_m$, i.e. there is a sequence of commutative algebraic groups

$$0 \to \mathbb{G}_m \to T \to J(X) \to 0. \tag{9.1}$$

Let $p \in X$ and let $\mathcal{I}^{\underline{0}}$ be a universal sheaf on $X \times J(X)$ such that its restriction at $p \times J(X)$ is trivial. Let $m : J(X) \times J(X) \to J(X)$ be the multiplication map and $\pi_{i,j} : X \times J(X) \times J(X) \to X \times J(X)$ be the projection maps. By the see-saw principle, the line bundles $\pi_{1,2}^* \mathcal{I}^{\underline{0}} \otimes \pi_{1,3}^* \mathcal{I}^{\underline{0}}$ and $(\mathrm{id}_X \times m)^* \mathcal{I}^{\underline{0}}$ on $X \times J(X) \times J(X)$ are isomorphic. Let σ be a nowhere vanishing section of $(\mathcal{I}^{\underline{0}})_{|p \times J(X)}$. The section σ induces nowhere vanishing sections $\overline{\sigma}$ of $(\pi_{1,2}^* \mathcal{I}^{\underline{0}} \otimes \pi_{1,3}^* \mathcal{I}^{\underline{0}})_{|p \times J(X) \times J(X)}$ and $\hat{\sigma}$ of $((\mathrm{id}_X \times m)^* \mathcal{I}^{\underline{0}})_{|p \times J(X) \times J(X)}$. Let $\phi : \pi_{1,2}^* \mathcal{I}^{\underline{0}} \otimes \pi_{1,3}^* \mathcal{I}^{\underline{0}} \to (\mathrm{id}_X \times m)^* \mathcal{I}^{\underline{0}}$ on $X \times J(X) \times J(X)$ be an isomorphism sending $\overline{\sigma}$ to $\hat{\sigma}$. A straightforward computation shows that ϕ makes the complement of the zero section in $\mathcal{I}^{\underline{0}}$ into a group scheme over X. As a consequence, for any $s \in X$ the isomorphism ϕ induces a group structure on the complement T_s of the zero section in $\mathcal{I}^{\underline{0}}_{|s \times J(X)}$.

Let p_i be smooth points of X such that $M = \mathcal{O}(\sum a_i p_i)$ and set $\delta := \sum_{a_i > 0} a_i p_i$ and $\gamma := -\sum_{a_i < 0} a_i p_i$. By equation (8.15), we get a canonical isomorphism

$$\mathcal{P}_{M|J(X)} \cong \bigotimes_i (\mathcal{I}^{\underline{0}}_{|p_i \times J(X)})^{-a_i}.$$

Hence T is the complement of the zero section in $\bigotimes_i (\mathcal{I}^{\underline{0}}_{|p_i \times J(X)})^{-a_i}$ and it carries a group structure which is induced by the group structures on the T_{p_i}.

This group structure makes T an abelian group and its natural group morphism onto $J(X)$ produces the exact sequence (9.1).

Claim 2: The action of $J(X)$ on $\overline{J}_X(\underline{q})$ lifts to an action of T on $(\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}$. Moreover, $\mathbb{G}_m \subset T$ acts on $(\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}$ fiberwise in the standard way by multiplication.

As for the previous claim, let $p \in X$ and let $\mathcal{I}^{\underline{q}}$ be a universal sheaf on $X \times \overline{J}_X(\underline{q})$ such that its restriction at $p \times \overline{J}_X(\underline{q})$ is trivial. Denote by $p_{1,2} : X \times J(X) \times \overline{J}_X(\underline{q}) \to X \times J(X)$ and by $p_{1,3} : X \times J(X) \times \overline{J}_X(\underline{q}) \to X \times \overline{J}_X(\underline{q})$ the projection maps and let $a : J(X) \times \overline{J}_X(\underline{q}) \to \overline{J}_X(\underline{q})$ be the action of $J(X)$ on $\overline{J}_X(\underline{q})$. In this case the see-saw principle gives an isomorphism $\psi : p_{1,2}^* \mathcal{I}^{\underline{0}} \otimes p_{1,3}^* \mathcal{I}^{\underline{q}} \to (\mathrm{id}_X \times a)^* \mathcal{I}^{\underline{q}}$. Moreover a suitable choice of ψ (analogous to the choise of ϕ in the previous claim) gives an action over X of the complement of the zero section in $\mathcal{I}^{\underline{0}}$ on $\mathcal{I}^{\underline{q}}$. Hence, for every $s \in X$, the isomorphism ψ induces an action of T_s on $\mathcal{I}^{\underline{q}}_{|s \times \overline{J}_X(\underline{q})}$. Since equation (8.15) gives the equality

$$\mathcal{P}_{M|\overline{J}_X(\underline{q})} = \bigotimes_i (\mathcal{I}^{\underline{q}}_{|p_i \times \overline{J}_X(\underline{q})})^{-a_i},$$

we finally get that T acts on $(\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}$ lifting the action of $J(X)$ on $\overline{J}_X(\underline{q})$. The second part of the claim follows from our description of the action.

According to Claim 2, the algebraic group T acts on any cohomology group $H^i(\overline{J}_X(\underline{q}), (\mathcal{P}_M)_{|\overline{J}_X(\underline{q})})$. Suppose that $H^i(\overline{J}_X(\underline{q}), (\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}) \neq 0$ for some index i. Consider a T-irreducible non-trivial submodule $0 \neq V \subseteq H^i(\overline{J}_X(\underline{q}), (\mathcal{P}_M)_{|\overline{J}_X(\underline{q})})$. Since T is commutative, V is a one-dimensional representation of T. Therefore the action of T on V is given by a character $\chi : T \to \mathbb{G}_m$ and, since $\mathbb{G}_m \subset T$ acts on $(\mathcal{P}_M)_{|\overline{J}_X(\underline{q})}$ fiberwise in the standard way by multiplication, it follows that $\chi_{|\mathbb{G}_m} = \mathrm{id}$. As a consequence, the character χ gives a splitting of the exact sequence (9.1), from which we deduce that $T \cong J(X) \times \mathbb{G}_m$. This is indeed equivalent to the fact that $(\mathcal{P}_M)_{|J(X)} = \mathcal{O}_{J(X)}$.

We conclude now by using the Lemma 8.3. Indeed, $\mathbb{J}_X$ is the disjoint union of $\mathbb{J}_X^{\underline{d}}$, as $\underline{d}$ varies among all multidegrees (see (2.3)). Therefore, it is enough to show that $(\mathcal{P}_M)_{|\mathbb{J}_X^{\underline{d}}} = \mathcal{O}_{\mathbb{J}_X^{\underline{d}}}$ for each multidegree $\underline{d}$. Fix such multidegree $\underline{d}$ and take a line bundle N of multidegree $\underline{d}$. The multiplication by N^{-1} induces an isomorphism $- \otimes N^{-1} : \mathbb{J}_X^{\underline{d}} \xrightarrow{\cong} J(X)$. Using Lemma 8.3, we now get

$$(\mathcal{P}_M)_{|\mathbb{J}_X^{\underline{d}}} = (- \otimes N^{-1})^*(\mathcal{P}_M)_{|\mathbb{J}_X^{\underline{d}}} = (- \otimes N^{-1})^*((\mathcal{P}_M)_{|J(X)}) = (- \otimes N^{-1})^*(\mathcal{O}_{J(X)}) = \mathcal{O}_{\mathbb{J}_X^{\underline{d}}}.$$

□

The previous Proposition implies the following two Corollaries, that generalize [Ari11, Cor. 2] and [Ari11, Cor. 3] to the case where X is not integral.

Corollary 9.2. *Assume that X satisfies condition (†) from (6.3). Let $M \in \mathrm{Pic}^{\underline{0}}(X) = J(X)$ and let $\underline{q}$ be a general polarization on X. If $H^i(\overline{J}_X(\underline{q}), \mathcal{P}_M) \neq 0$ for some i then $M_{|X_{\mathrm{sm}}} = \mathcal{O}_{X_{\mathrm{sm}}}$.*

Proof. Consider the Abel map $A_L : X \to \overline{\mathbb{J}}_X$ for some $L \in \mathrm{Pic}(X)$. Clearly we have that $A_L(X_{\mathrm{sm}}) \subseteq \mathbb{J}_X$. Using Proposition 9.1 and Proposition 8.5, we get

$$M_{|X_{\mathrm{sm}}} = (A_L^*\mathcal{P}_M)_{|X_{\mathrm{sm}}} = (A_L|_{X_{sm}})^*((\mathcal{P}_M)_{|\mathbb{J}_X}) = (A_L|_{X_{sm}})^*(\mathcal{O}_{\mathbb{J}_X}) = \mathcal{O}_{X_{\mathrm{sm}}}.$$

□

For any general polarization $\underline{q}$ on X, consider the locus

$$\mathcal{N}(\underline{q}) := \{M \in J(X) : H^i(\overline{J}_X(\underline{q}), \mathcal{P}_M) \neq 0 \text{ for some } i\} \subseteq J(X).$$

Notice that, by semicontinuity, $\mathcal{N}(\underline{q})$ is a closed subset of $J(X)$ and that

$$\mathcal{N}(\underline{q}) = \mathrm{supp}(Rp_{2*}(\mathcal{P}_{|\overline{J}_X(\underline{q})})),$$

where $p_2 : \overline{J}_X(\underline{q}) \times J(X) \to J(X)$ is the second projection.

Corollary 9.3. *Assume that X satisfies condition (†) from (6.3) and let $\underline{q}$ be a general polarization on X. Then*

$$\dim \mathcal{N}(\underline{q}) \leq p_a(X) - g^\nu(X).$$

Proof. The normalization morphism $\nu : X^\nu \to X$ induces by pull-back a smooth and surjective morphism $\nu^* : J(X) \to J(X^\nu)$ with fibers of dimension equal to $p_a(X) - g^\nu(X)$. Denote by $\widetilde{N} \subseteq J(X^\nu)$ the locus of line bundles on X^ν that are trivial on $\nu^{-1}(X_{\mathrm{sm}}) \subseteq X^\nu$.

Claim: $\widetilde{N}$ is a countable set.

Set $F := X^\nu \setminus \nu^{-1}(X_{\mathrm{sm}})$. We have an exact sequence

$$\mathbb{Z}^F \xrightarrow{\alpha} \mathrm{Pic}(X^\nu) \to \mathrm{Pic}(\nu^{-1}(X_{\mathrm{sm}})),$$

where the last map is the restriction map and α sends $\{m_P\}_{P \in F} \in \mathbb{Z}^F$ into $\mathcal{O}_X(\sum_{P \in F} m_P \cdot P)$. The claim follows since $\widetilde{N}$ is equal to $\mathrm{Im}(\alpha) \cap \mathrm{J(X^\nu)}$.

The result now follows since Corollary 9.2 implies that $\nu^*(\mathcal{N}(\underline{q})) \subseteq \widetilde{N}$. □

Proposition 9.1 can be strengthened for $i = 0$ if the curve X has locally planar singularities.

Proposition 9.4. *Assume that X has locally planar singularities and let $\underline{q}$ be a general polarization on X. If $M \in \mathrm{Pic}^{\underline{0}}(X) = J(X)$ is such that $H^0(\overline{J}_X(\underline{q}), \mathcal{P}_M) \neq 0$, then $\mathcal{P}_{M|\overline{J}_X(\underline{q})} = \mathcal{O}_{\overline{J}_X(\underline{q})}$.*

Proof. We know that $\overline{J}_X(\underline{q})$ is a connected reduced projective scheme over k by Fact 2.11, Corollary 2.12 and Corollary 4.4. As already observed in (8.7), we have that $\mathcal{P}_{M|\overline{J}_X(\underline{q})} \in \mathrm{Pic}^o(\overline{J}_X(\underline{q}))$. We can apply Lemma 9.5 below in order to conclude that $\mathcal{P}_{M|\overline{J}_X(\underline{q})} = \mathcal{O}_{\overline{J}_X(\underline{q})}$. □

Lemma 9.5. *Let V be a connected reduced projective scheme over an algebraically closed field k. Let $\mathcal{L}$ be a line bundle belonging to $\mathrm{Pic}^o(V)$, i.e. the connected component of $\mathrm{Pic}(V)$ containing the identity. If $H^0(V,\mathcal{L})\neq 0$ then $\mathcal{L}=\mathcal{O}_V$.*

Proof. Assume first that V is irreducible. Let s be a non-zero section of $\mathcal{L}$ and consider it as a map $\widetilde{s}:\mathcal{O}_V\to\mathcal{L}$. Since V is irreducible and $s\not\equiv 0$, the map $\widetilde{s}$ is generically injective. This implies that the kernel of $\widetilde{s}$ is a torsion sheaf of $\mathcal{O}_V$, hence it is zero since $\mathcal{O}_V$ does not contain torsion sheaves. In other words, $\widetilde{s}$ realizes $\mathcal{O}_V$ as a subsheaf of $\mathcal{L}$. However, since $\mathcal{L}\in\mathrm{Pic}^o(V)$ by assumption, $\mathcal{L}$ and $\mathcal{O}_V$ are algebraically equivalent and therefore they have the same Hilbert polynomial with respect to any ample line bundle on V. This can only happen if $\widetilde{s}$ is an isomorphism, q.e.d.

Let us now prove the general case. Let $V_1,\ldots,V_r$ be the irreducible components of V. Take a non-zero section $s\in H^0(V,L)$ and consider its zero set:

$$Z(s):=\{P\in V: s(P)=0\}\subsetneq V.$$

For each irreducible component V_i, consider the restriction $s_i:=s_{|V_i}\in H^0(V_i,\mathcal{L}_{|V_i})$. If $s_i\equiv 0$ then clearly $V_i\subseteq Z(s)$. Otherwise s_i defines a non-zero section of the line bundle $\mathcal{L}_{|V_i}$ which clearly belongs to $\mathrm{Pic}^o(V_i)$. By what proved above, we must have that $\mathcal{L}_{|V_i}=\mathcal{O}_{V_i}$. Since V_i is projective and reduced, then the space $H^0(V_i,L_{|V_i})=H^0(V_i,\mathcal{O}_{V_i})=k$ can be identified with the constant functions on V_i. In particular, s_i is given by a constant non-zero function on V_i and therefore s_i is nowhere vanishing on V_i. This implies that $V_i\cap Z(s)=\emptyset$.

Repeating the same argument for each irreducible component V_i of X, we deduce that either $Z(s)$ is empty or it is union of connected components of X, hence it is equal to X since X is assumed to be connected. However, s was assumed to be non-zero, hence the second possibility cannot occur. Therefore $Z(s)=\emptyset$, or in other words s is a nowhere section of $\mathcal{L}$, which clearly implies that $\mathcal{L}=\mathcal{O}_V$. □

10. PROOF OF THEOREM C FOR NODAL CURVES

The aim of this section is to prove Theorem C from the introduction for nodal curves. The key fact about fine compactified Jacobians of nodal curves that we are going to use is the following result of Alexeev-Nakamura [AN99] and Alexeev [Ale04].

Fact 10.1 (Alexeev-Nakamura, Alexeev). *Let X be a nodal curve and let $\underline{q}$ be a general polarization on X. Then we have that*

$$h^1(\overline{J}_X(\underline{q}),\mathcal{O}_{\overline{J}_X(\underline{q})})=p_a(X). \tag{10.1}$$

Proof. Every fine compactified Jacobian of a nodal curve X is a stable quasiabelian variety of dimension $p_a(X)$ in the sense of [AN99], as proved by Alexeev in [Ale04, Sec. 5.5]. Therefore, the conclusion now follows from [AN99, Thm. 4.3], which says that for a stable quasiabelian variety of dimension g the structure sheaf has the same cohomology groups of the structure sheaf of an abelian variety of dimension g. □

We will now prove Theorem C from the introduction for fine compactified Jacobians $\overline{J}_X(\underline{q})$ that satisfy condition (10.1), and in particular for all fine compactified Jacobians of nodal curves by the above Fact 10.1. Note that, a posteriori, it will follow from Corollary B that every fine compactified Jacobian $\overline{J}_X(\underline{q})$ of any curve X with locally planar singularities satisfies condition (10.1). However, we do not know a direct proof of this fact avoiding the use of the Fourier-Mukai transform.

The special case of Theorem C that we are going to prove will follow from a more general result involving the semiuniversal deformation family of X. Let us fix the set-up. Consider the semiuniversal deformation family $\pi:\mathcal{X}\to\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ for X as in §3.3. The generalized Jacobian $J(X)$ and the fine compactified Jacobian $\overline{J}_X(\underline{q})$ deform over $\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ to, respectively, the universal generalized Jacobian $v:J(\mathcal{X})\to\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ (see Fact 4.6) and the universal fine compactified Jacobian $u:\overline{J}_{\mathcal{X}}(\underline{q})\to\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$ with respect to the polarization $\underline{q}$ (see Theorem 4.2). By using the universal sheaf $\widehat{\mathcal{I}}$ on $\mathcal{X}\times_{\mathrm{Spec}\,\mathrm{R}_\mathrm{X}}\overline{\mathbb{J}}_{\mathcal{X}}$ (see Fact 3.6(i)), it is possible to define a universal Poincaré line bundle $\mathcal{P}^{\mathrm{un}}$ on the fiber product $\overline{J}_{\mathcal{X}}(\underline{q})\times_{\mathrm{Spec}\,\mathrm{R}_\mathrm{X}}J(\mathcal{X})$, similarly to the definition (8.1) of the Poincaré line bundle $\mathcal{P}$ on $\overline{J}_X(\underline{q})\times J(X)$ via the determinant of cohomology. Indeed, since the determinant of cohomology commutes with base change, it follows that $\mathcal{P}^{\mathrm{un}}$ restricts to $\mathcal{P}$ on the central fiber of $\overline{J}_{\mathcal{X}}(\underline{q})\times_{\mathrm{Spec}\,\mathrm{R}_\mathrm{X}}J(\mathcal{X})\to\mathrm{Spec}\,\mathrm{R}_\mathrm{X}$.

Assume now that X has locally planar singularities and that $\overline{J}_X(\underline{q})$ satisfies condition (10.1). In analogy with the definition (8.7) of the map $\beta_{\underline{q}}$, the existence of a universal Poincaré line bundle $\mathcal{P}^{\mathrm{un}}$ on the fiber product $\overline{J}_{\mathcal{X}}(\underline{q}) \times_{\operatorname{Spec} \mathrm{R_X}} J(\mathcal{X})$ defines a homomorphism

$$\beta_{\underline{q}}^{\mathrm{un}} : J(\mathcal{X}) \to \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})), \tag{10.2}$$

between two group schemes which are smooth, separated and of finite type over $\operatorname{Spec} \mathrm{R_X}$ (see Fact 4.6 and Theorem 5.1(iv)). Moreover, by construction, the fibers of $J(\mathcal{X}) \to \operatorname{Spec} \mathrm{R_X}$ and of $\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to \operatorname{Spec} \mathrm{R_X}$ are non empty and geometrically connected. Therefore, using that $\operatorname{Spec} \mathrm{R_X}$ is smooth over k by Theorem 3.7(ii), we get that the schemes $J(\mathcal{X})$ and $\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$ are smooth over k and connected, hence irreducible. Since $\mathcal{P}^{\mathrm{un}}$ restricts to $\mathcal{P}$ on the central fiber of $\overline{J}_{\mathcal{X}}(\underline{q}) \times_{\operatorname{Spec} \mathrm{R_X}} J(\mathcal{X}) \to \operatorname{Spec} \mathrm{R_X}$, the homomorphism $\beta_{\underline{q}}^{\mathrm{un}}$ restricts to the homomorphism $\beta_{\underline{q}} : J(X) \to \operatorname{Pic}^o(\overline{J}_X(\underline{q}))$ on the central fiber.

The main result of this Section is the following

Theorem 10.2. *Let X be a reduced curve with locally planar singularities and let $\overline{J}_X(\underline{q})$ be a fine compactified Jacobian of X having the property that $h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X)$. Then the group homomorphism*

$$\beta_{\underline{q}}^{\mathrm{un}} : J(\mathcal{X}) \to \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$$

is an isomorphism.

Proof. Consider the open subset $U \subseteq \operatorname{Spec} \mathrm{R_X}$ consisting of all the points s such that the fiber $\mathcal{X}_s$ of the universal family $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R_X}$ over s is smooth or has a unique singular point which is a node. By Lemma 3.5, the complement of U inside $\operatorname{Spec} \mathrm{R_X}$ has codimension at least two.

CLAIM 1: The restriction of $\beta_{\underline{q}}^{\mathrm{un}}$ to U

$$(\beta_{\underline{q}}^{\mathrm{un}})_{|U} : J(\mathcal{X})_{|U} \to \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))_{|U}$$

is an isomorphism. In particular, $\beta_{\underline{q}}^{\mathrm{un}}$ is an isomorphism in codimension one.

Indeed, since the map $\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))_{|U} \to U$ is flat, using [EGAIV4, (17.9.5)] it is enough to prove that the restriction of $\beta_{\underline{q}}^{\mathrm{un}}$ to the fibers over s

$$(\beta_{\underline{q}}^{\mathrm{un}})_s : J(\mathcal{X})_s = J(\mathcal{X}_s) \to \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))_s = \operatorname{Pic}^o(\overline{J}_{\mathcal{X}_s}(\underline{q}^s)) \tag{10.3}$$

is an isomorphism for every $s \in U$. By the definition of U, the fiber $\mathcal{X}_s$ can be of three types:

(i) $\mathcal{X}_s$ is smooth;
(ii) $\mathcal{X}_s$ is an irreducible curve having a unique singular point which is a node;
(iii) $\mathcal{X}_s$ has two smooth irreducible components $\mathcal{X}_s^1$ and $\mathcal{X}_s^2$ which meet in a separating node.

In case (i), the morphism $(\beta_{\underline{q}}^{\mathrm{un}})_s$ is well-known to be an isomorphism (see e.g. [GIT94, Prop. 6.9]). In case (ii), the fact that the morphism $(\beta_{\underline{q}}^{\mathrm{un}})_s$ is an isomorphism is a particular case of the main result of Esteves-Gagné-Kleiman in [EGK00], who proved that the same conclusion is true for any integral curve with double points singularities. Finally, in case (iii), using Proposition 6.6, we get that $\overline{J}_{\mathcal{X}_s}(\underline{q}^s) \cong \overline{J}_{\mathcal{X}_s^1}(\underline{q}^1) \times \overline{J}_{\mathcal{X}_s^2}(\underline{q}^2)$ for some general polarizations $\underline{q}^i$ on $\mathcal{X}_s^i$ (for $i = 1, 2$). The diagram (8.30) in Section 8 translates into the following commutative diagram

$$\begin{tikzcd}
& \operatorname{Pic}^o(\overline{J}_{\mathcal{X}_s^1}(\underline{q}^1)) \times \operatorname{Pic}^o(\overline{J}_{\mathcal{X}_s^2}(\underline{q}^2)) \arrow[dr, "p_1^*(-)\otimes p_2^*(-)"] & \\
J(\mathcal{X}_s^1) \times J(\mathcal{X}_s^2) \arrow[ur, "\beta_{\underline{q}^1}\times\beta_{\underline{q}^2}"] & & \operatorname{Pic}^o(\overline{J}_{\mathcal{X}_s^1}(\underline{q}^1) \times \overline{J}_{\mathcal{X}_s^2}(\underline{q}^2)) \arrow[d, "\cong"'] \\
J(\mathcal{X}_s) \arrow[u, "\cong"'] \arrow[rr, "\beta_{\underline{q}^s}"'] & & \operatorname{Pic}^o(\overline{J}_{\mathcal{X}_s}(\underline{q}^s))
\end{tikzcd} \tag{10.4}$$

The maps $\beta_{\underline{q}^1}$ and $\beta_{\underline{q}^2}$ are isomorphism since $\mathcal{X}_s^1$ and $\mathcal{X}_s^2$ are smooth curves (as in case (i)); hence the fact that $\beta_{\underline{q}^s}$ is an isomorphism follows from the previous diagram together with the Lemma 10.3 below.

CLAIM 2: $\beta_{\underline{q}}^{\mathrm{un}}$ is an open embedding.

Indeed, since $\beta_{\underline{q}}^{\mathrm{un}}$ is a birational map between two integral schemes which is an isomorphism in codimension one (by Claim 1) and the codomain is normal and locally factorial (being smooth), we deduce that $\beta_{\underline{q}}^{\mathrm{un}}$ is a local isomorphism by Van der Waerden's theorem on the purity of the ramification locus (see [EGAIV4, (21.12.12)]). In particular, $\beta_{\underline{q}}^{\mathrm{un}}$ is quasi-finite. Moreover, since $\beta_{\underline{q}}^{\mathrm{un}}$ is birational (by

Claim 1) and separated (which follows from the fact that $J(\mathcal{X}) \to \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$ is separated, see [Har77, Cor. 4.6(e)]) and the codomain is normal (being smooth), we deduce that $\beta_{\underline{q}}^{\mathrm{un}}$ is an open embedding by Zariski's main theorem (see [EGAIII1, (4.4.9)]).

We can now easily conclude the proof of the Theorem. Indeed, for any $s \in \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$, the restriction $(\beta_{\underline{q}}^{\mathrm{un}})_s$ of (10.3) is a group homomorphism between two connected and smooth algebraic groups over $k(s)$ of the same dimension, $p_a(X)$, which is moreover an open embedding by Claim 2. This forces $(\beta_{\underline{q}}^{\mathrm{un}})_s$ to be surjective (see e.g. [BLR90, Sec. 7.3, Lemma 1]), hence an isomorphism. Since the map $\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$ is flat, using again [EGAIV4, (17.9.5)], it follows that $\beta_{\underline{q}}^{\mathrm{un}}$ is an isomorphism, q.e.d.

□

Lemma 10.3. *Let V_1 and V_2 be two schemes proper over a field k. Assume that $\operatorname{Pic}^o(V_1)$ and $\operatorname{Pic}^o(V_2)$ are smooth. Then the natural pull-back map*

$$\begin{aligned} \boxtimes : \operatorname{Pic}^o(V_1) \times \operatorname{Pic}^o(V_2) &\longrightarrow \operatorname{Pic}^o(V_1 \times V_2), \\ (L, M) &\mapsto p_1^*(L) \otimes p_2^*(M), \end{aligned}$$

where p_i is the projection onto V_i (for $i = 1, 2$), is an isomorphism.

Proof. Clearly, the map $\boxtimes$ is an injective group homomorphism. Since $\operatorname{Pic}^o(V_i)$ is smooth, then its dimension is equal to $h^1(V_i, \mathcal{O}_{V_i})$ by [FGA05, Cor. 9.5.13]. Since $\dim \operatorname{Pic}^o(V_1 \times V_2) \leq h^1(V_1 \times V_2, \mathcal{O}_{V_1 \times V_2})$ by loc. cit. and $\boxtimes$ is injective, we get that

$$\begin{aligned} h^1(V_1, \mathcal{O}_{V_1}) + h^1(V_2, \mathcal{O}_{V_2}) = \dim \operatorname{Pic}^o(V_1) + \dim \operatorname{Pic}^o(V_2) \leq \operatorname{Pic}^o(V_1 \times V_2) \leq \\ \leq h^1(V_1 \times V_2, \mathcal{O}_{V_1 \times V_2}) = h^1(V_1, \mathcal{O}_{V_1}) + h^1(V_2, \mathcal{O}_{V_2}). \end{aligned}$$

This implies that $\operatorname{Pic}^o(V_1 \times V_2)$ is smooth of the same dimension as $\operatorname{Pic}^o(V_1) \times \operatorname{Pic}^o(V_2)$ by loc. cit. Since $\operatorname{Pic}^o(V_1 \times V_2)$ is connected, this easily implies that $\boxtimes$ is surjective (see e.g. [BLR90, Sec. 7.3, Lemma 1]; hence $\boxtimes$ is an isomorphism, q.e.d. □

Corollary 10.4. *Let X be a reduced curve with locally planar singularities and let $\overline{J}_X(\underline{q})$ be a fine compactified Jacobian of X having the property that $h^1(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = p_a(X)$. Then the group homomorphism*

$$\beta_{\underline{q}} : J(X) \to \operatorname{Pic}^o(\overline{J}_X(\underline{q}))$$

is an isomorphism.

Proof. This follows directly from Theorem 10.2 by restricting to the central fiber. □

11. PROOF OF THEOREM A, COROLLARY B AND THEOREM C

The aim of this section is to prove the first three results that were stated in the introduction, namely Theorem A, Corollary B and Theorem C.

A key role will be played by the semiuniversal deformation family $\pi : \mathcal{X} \to \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$ for X as in §3.3. More precisely, we will be looking at the following Cartesian diagram

(11.1)
$$\begin{tikzcd}
& \overline{J}_{\mathcal{X}}(\underline{q}) \times_{\operatorname{Spec} \mathrm{R}_{\mathrm{X}}} J(\mathcal{X}) \arrow[dl, "\widetilde{u}"'] \arrow[dr, "\widetilde{v}"] & \\
J(\mathcal{X}) \arrow[dr, "v"] & \square & \overline{J}_{\mathcal{X}}(\underline{q}) \arrow[dl, "u"'] \arrow[ul, bend right, "\widetilde{\zeta}"'] \\
& \operatorname{Spec} \mathrm{R}_{\mathrm{X}} \arrow[ul, bend left, "\zeta"] &
\end{tikzcd}$$

where $v : J(\mathcal{X}) \to \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$ is the universal generalized Jacobian (see Fact 4.6), $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$ is the universal fine compactified Jacobian with respect to the polarization $\underline{q}$ (see Theorem 4.2), ζ is the zero section of v and $\widetilde{\zeta} := \mathrm{id} \times \zeta$. As observed already in Section 10, there exists a universal Poincaré line bundle $\mathcal{P}^{\mathrm{un}}$ on the fiber product $\overline{J}_{\mathcal{X}}(\underline{q}) \times_{\operatorname{Spec} \mathrm{R}_{\mathrm{X}}} J(\mathcal{X})$ which restricts to $\mathcal{P}$ on the central fiber $\overline{J}_X(\underline{q}) \times J(X)$.

Assuming that X has locally planar singularities and setting $g := p_a(X)$, the morphisms appearing in the above diagram satisfy the following properties: the morphism v (hence also $\widetilde{v}$) is smooth of relative dimension g (see Fact 3.6(i)); the morphism u (and hence also $\widetilde{u}$) is projective, flat of relative dimension g with trivial relative dualizing sheaf and geometrically connected fibers (see Theorem 4.3, Corollary 4.4 and Corollary 4.5). Moreover all the schemes appearing in diagram (11.1) are smooth over $\overline{k}=k$: $\mathrm{Spec}\mathrm{R_X}$ is smooth by Theorem 3.7(ii); $\overline{J}_{\mathcal{X}}(\underline{q})$ is smooth by Theorem 4.3; $J(\mathcal{X})$ (resp. $\overline{J}_{\mathcal{X}}(\underline{q})\times_{\mathrm{Spec}\,\mathrm{R_X}} J(\mathcal{X})$) is smooth because the morphism v (resp. $\widetilde{v}$) is smooth over a smooth codomain.

The following result is a generalization of a well-known result of Mumford for abelian varieties (see [Mum70, Sec. III.13]) and it is the key for the proof of our main theorems.

Theorem 11.1. *Let X be a reduced curve with locally planar singularities of arithmetic genus $g:=p_a(X)$ and let $\underline{q}$ be a general polarization. There is a natural isomorphism of complexes of coherent sheaves on $J(\mathcal{X})$:*

$$\Phi : R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) \to \zeta_*(\mathcal{O}_{\mathrm{Spec}\,\mathrm{R_X}})[-g]. \tag{11.2}$$

In particular, we get that

$$Rp_{2*}\mathcal{P} = \mathbf{k}(0)[-g] \tag{11.3}$$

where $\mathbf{k}(0)$ denotes the skyscraper sheaf supported at the origin $0=[\mathcal{O}_X]\in J(X)$, and $p_2 : \overline{J}_X(\underline{q})\times J(X)\to J(X)$ is the projection onto the second factor.

Proof. Clearly, the last assertion follows from the first one by base change to the central fiber of $v : J(\mathcal{X})\to \mathrm{Spec}\,\mathrm{R_X}$; hence it is enough to prove (11.2).

We will first explain how the morphism Φ is defined. By applying base change (see e. g. [Huy06, Rmk. 3.33]) to the diagram (11.1), we get a morphism

$$\zeta^*(R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) \to Ru_*(\widetilde{\zeta}^*(\mathcal{P}^{\mathrm{un}})). \tag{11.4}$$

It follows from Proposition 8.1 that $\widetilde{\zeta}^*(\mathcal{P}^{\mathrm{un}})\cong \mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}$. Since the complex of sheaves $Ru_*(\mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})})$ is concentrated in cohomological degrees from 0 to g, we get a morphism of complexes of sheaves

$$Ru_*(\widetilde{\zeta}^*(\mathcal{P}^{\mathrm{un}})) \cong Ru_*(\mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}) \to R^g u_*(\mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})})[-g]. \tag{11.5}$$

Moreover, since the morphism u is proper of relative dimension g, with trivial relative dualizing sheaf and geometrically connected fibers, then the relative duality applied to u gives that (see [Har77, Cor. 11.2(g)]) :

$$R^g u_*(\mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}) \cong \mathcal{O}_{\mathrm{Spec}\,\mathrm{R_X}}. \tag{11.6}$$

By composing the morphisms (11.4), (11.5) and using the isomorphism (11.6), we get a morphism

$$\zeta^*(R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) \to \mathcal{O}_{\mathrm{Spec}\,\mathrm{R_X}}[-g]. \tag{11.7}$$

Since ζ_* is right adjoint to ζ^*, the morphism (11.7) gives rise to the morphism Φ.

The remaining part of the proof will be devoted to showing that the morphism Φ is an isomorphism of complexes of sheaves. We need three preliminary results that we collect under the name of Claims. The first result says that Φ is generically an isomorphism. More precisely, let $(\mathrm{Spec}\,\mathrm{R_X})_{\mathrm{sm}}$ be the open subset of $\mathrm{Spec}\,\mathrm{R_X}$ consisting of the points $s\in \mathrm{Spec}\,\mathrm{R_X}$ such that $\mathcal{X}_s$ is smooth. Then we have:

CLAIM 1: The morphism Φ is an isomorphism over the open subset $v^{-1}((\mathrm{Spec}\,\mathrm{R_X})_{\mathrm{sm}})$.

Indeed, $v^{-1}((\mathrm{Spec}\mathrm{R_X})_{\mathrm{sm}})$ is an abelian group scheme over $(\mathrm{Spec}\mathrm{R_X})_{\mathrm{sm}}$ via the map v. Claim 1 follows then from a well-known theorem of Mumford (see [Mum70, Sec. III.13]).

CLAIM 2: $R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})\cong R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})[g]$ is a Cohen-Macaulay sheaf of codimension g.

Let us first prove that

$$\mathrm{codim}(\mathrm{supp}(R\widetilde{u}_*\mathcal{P}^{\mathrm{un}})) = g. \tag{11.8}$$

First of all, Claim 1 gives that $\mathrm{codim}(\mathrm{supp}(R\widetilde{u}_*\mathcal{P}^{\mathrm{un}}))\leq g$. In order to prove the reverse inequality, we stratify the scheme $\mathrm{Spec}\,\mathrm{R_X}$ into locally closed subsets according to the geometric genus of the fibers of the universal family $\mathcal{X}\to\mathrm{Spec}\,\mathrm{R_X}$:

$$(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^\nu=\mathrm{k}} := \{\mathrm{s}\in\mathrm{Spec}\,\mathrm{R_X}\ :\ \mathrm{g}^\nu(\mathcal{X}_\mathrm{s})=\mathrm{k}\},$$

for any $g^\nu(X)\leq k\leq p_a(X)=g$. Corollary 3.4 gives that $\mathrm{codim}(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^\nu=\mathrm{k}}\geq \mathrm{g}-\mathrm{k}$. On the other hand, on the fibers of v over $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^\nu=\mathrm{k}}$, the sheaf $R\widetilde{u}_*\mathcal{P}^{\mathrm{un}}$ has support of codimension at least k by Corollary 9.3. Therefore, we get

$$\mathrm{codim}(\mathrm{supp}(R\widetilde{u}_*\mathcal{P}^{\mathrm{un}})\cap v^{-1}((\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^\nu=\mathrm{k}}))\geq \mathrm{g}\text{ for any }\mathrm{g}^\nu(\mathrm{X})\leq\mathrm{k}\leq\mathrm{g}. \tag{11.9}$$

Since the locally closed subsets $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^{\nu}=\mathrm{k}}$ form a stratification of $\mathrm{Spec}\,\mathrm{R_X}$, we deduce that $g \leq \mathrm{codim}(\mathrm{supp}(R\widetilde{u}_*\mathcal{P}^{\mathrm{un}}))$, which concludes the proof of (11.8).

We now apply the relative duality (see e.g. [Har77, Chap. VII.3]) to the proper morphism $\widetilde{u}$. Since $\widetilde{u}$ is flat of relative dimension g and it has trivial relative dualizing sheaf, we get that

$$R\mathcal{H}om(R\widetilde{u}_*\,(\mathcal{P}^{\mathrm{un}})^{-1}, \mathcal{O}_{J(\mathcal{X})}) \cong R\widetilde{u}_*\mathcal{P}^{\mathrm{un}}[g], \tag{11.10}$$

where $(\mathcal{P}^{\mathrm{un}})^{-1}$ is the inverse of $\mathcal{P}^{\mathrm{un}}$, i.e. $(\mathcal{P}^{\mathrm{un}})^{-1} := \mathcal{H}om(\mathcal{P}^{\mathrm{un}}, \mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})\times_{\mathrm{Spec}\,\mathrm{R_X}} J(\mathcal{X})})$. The right hand side of (11.10) can be computed using the following spectral sequence (see [Huy06, Chap. 3, Formula (3.2)]):

$$E_2^{p,q} = \mathcal{E}xt^p(R^{-q}\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}, \mathcal{O}_{J(\mathcal{X})}) \Rightarrow \mathcal{E}xt^{p+q}(R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}, \mathcal{O}_{J(\mathcal{X})}), \tag{11.11}$$

where clearly $E_2^{p,q} = 0$ unless $0 \leq -q \leq g$ and $p \geq 0$. If we denote by i the involution of the group scheme $v : J(\mathcal{X}) \to \mathrm{Spec}\,\mathrm{R_X}$ that sends $\mathcal{M} \in J(\mathcal{X})$ into $\mathcal{M}^{-1} \in J(\mathcal{X})$, then Proposition 8.1 gives that $(\mathcal{P}^{\mathrm{un}})^{-1} = (\mathrm{id} \times \mathrm{i})^*(\mathcal{P}^{\mathrm{un}})$; hence

$$R\widetilde{u}_*\,(\mathcal{P}^{\mathrm{un}})^{-1} = i^*(R\widetilde{u}_*\mathcal{P}^{\mathrm{un}}). \tag{11.12}$$

In particular, the complex $R\widetilde{u}_*\,(\mathcal{P}^{\mathrm{un}})^{-1}$ has codimension g by (11.8). This implies that for any $0 \leq -q \leq g$, the sheaf $R^{-q}\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}$ has codimension at least g; hence, since the dualizing sheaf of $J(\mathcal{X})$ is trivial, we get that (see [HL97, Prop. 1.1.6]):

$$E_2^{p,q} = \mathcal{E}xt^p(R^{-q}\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}, \mathcal{O}_{J(\mathcal{X})}) = 0 \quad \text{for every } p < g \text{ and every } q. \tag{11.13}$$

Using the spectral sequence (11.11) and the vanishing (11.13), it is easily seen that the complex in the right hand side of (11.10) can have non-vanishing cohomology only in non-negative degrees. On the other hand, since $\widetilde{u}$ has fibers of dimension g, the complex in the left hand side of (11.10) can have non-vanishing cohomology only in degrees belonging to the interval $[-g, 0]$. We deduce that $R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})$ is supported in degree g, i.e. that

$$R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) \cong R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})[-g]. \tag{11.14}$$

Similarly, also the complex $R\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}$ is supported in cohomological degree g by (11.12). Hence the spectral sequence (11.11) together with the isomorphism (11.10) give that

$$\mathcal{E}xt^p(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}, \mathcal{O}_{J(\mathcal{X})}) = \begin{cases} R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) & \text{if } p = g, \\ 0 & \text{if } p \neq g. \end{cases} \tag{11.15}$$

This implies that $R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})^{-1}$ is Cohen-Macaulay of codimension g by [BH98, Cor. 3.5.11]. Using (11.12), we get that also $R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})$ is Cohen-Macaulay of codimension g, and the proof of Claim 2 is now complete.

CLAIM 3: We have a set-theoretic equality $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) = \mathrm{Im}(\zeta)$.

Observe that the restriction of $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ to the fiber $v^{-1}(s) = J(\mathcal{X}_s)$ over the point $s \in \mathrm{Spec}\mathrm{R_X}$ is equal to the locus of all elements $M \in J(\mathcal{X}_s)$ such that $H^g(\overline{J}_{\mathcal{X}_s}(\underline{q}^s), \mathcal{P}^s_M) \neq 0$, where we have set $\mathcal{P}^s_M := (\mathcal{P}^{\mathrm{un}})_{|\overline{J}_{\mathcal{X}_s}(\underline{q}^s)\times\{M\}}$. Since $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ has trivial dualizing sheaf by Corollary 4.5, Serre's duality gives that

$$H^g(\overline{J}_{\mathcal{X}_s}(\underline{q}^s), \mathcal{P}^s_M) \cong H^0(\overline{J}_{\mathcal{X}_s}(\underline{q}^s), (\mathcal{P}^s_M)^{-1})^\vee = H^0(\overline{J}_{\mathcal{X}_s}(\underline{q}^s), \mathcal{P}^s_{M^{-1}})^\vee.$$

Applying now Proposition 9.4, we get the set-theoretic equality

$$\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) \cap J(\mathcal{X}_s) = \{M \in J(\mathcal{X}_s) \,:\, \mathcal{P}^s_{M^{-1}} = \mathcal{O}_{\overline{J}_{\mathcal{X}_s}(\underline{q}^s)}\} \text{ for every } s \in \mathrm{Spec}\,\mathrm{R_X}. \tag{11.16}$$

If moreover the curve $\mathcal{X}_s$ is nodal then, combining (11.16) with Corollary 10.4 and Fact 10.1, we get

$$\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) \cap J(\mathcal{X}_s) = \{\mathcal{O}_{\mathcal{X}_s}\} \text{ for every } s \in \mathrm{Spec}\,\mathrm{R_X} \text{ such that } \mathcal{X}_{\mathrm{s}} \text{ is nodal.} \tag{11.17}$$

The above formula (11.17) allows us to improve the estimate (11.9) on the codimension of the intersection of $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ with the locally closed subset $v^{-1}((\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^{\nu}=\mathrm{k}})$. Indeed, by looking at the proof of (11.9), we see that can have an equality in (11.9) for some $g^\nu(X) \leq k \leq g$ only if:

- v maps $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ to a generic point η of $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^{\nu}=\mathrm{k}}$ of codimension $g-k$ in $\mathrm{Spec}\,\mathrm{R_X}$,
- the codimension of $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) \cap v^{-1}(\eta)$ in $J(\mathcal{X}_\eta)$ is equal to k.

However, since a generic point η of the stratum $(\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^{\nu}=\mathrm{k}}$ of $\mathrm{Spec}\,\mathrm{R_X}$ is such that $\mathcal{X}_\eta$ is nodal by Fact 3.3(ii), formula (11.17) tells us that equality in (11.9) is only possible for $k = g$; in other words we have that

$$\mathrm{codim}(\mathrm{supp}(R\widetilde{u}_*\mathcal{P}^{\mathrm{un}}) \cap v^{-1}((\mathrm{Spec}\,\mathrm{R_X})^{\mathrm{g}^{\nu}=\mathrm{k}})) \geq \mathrm{g}+1 \text{ for any } \mathrm{g}^{\nu}(\mathrm{X}) \leq \mathrm{k} < \mathrm{g}. \tag{11.18}$$

After these preliminaries, we can now finish the proof of Claim 3. Since $R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})$ is a Cohen-Macaulay sheaf of codimension g, then all the irreducible components of $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ have codimension g by [Mat89, Thm. 6.5(iii), Thm. 17.3(i)]. Clearly $\mathrm{Im}(\zeta)$ is one of the irreducible components of $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$. If Z is another irreducible component of $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ then, using (11.18) and the fact that Z has codimension g, we get that $v(Z)$ must contain the generic point η of $\mathrm{Spec}\,\mathrm{R}_{\mathrm{X}}$. Then Claim 1 implies that necessarily we must have $Z = \mathrm{Im}(\zeta)$, q.e.d.

CLAIM 4: We have a scheme-theoretic equality $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) = \mathrm{Im}(\zeta)$.

Since the subscheme $\mathrm{Im}(\zeta)$ is reduced, Claim 3 gives the inclusion of subschemes $\mathrm{Im}(\zeta) \subseteq \mathrm{supp}(\mathrm{R}^{\mathrm{g}}\widetilde{\mathrm{u}}_*(\mathcal{P}^{\mathrm{un}}))$. Moreover, Claim 1 says that this inclusion is generically an equality; in particular $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ is generically reduced. Moreover, since $R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})$ is a Cohen-Macaulay sheaf by Claim 2, Lemma 11.2 below implies that $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}))$ is reduced. Therefore, we must have the equality of subschemes $\mathrm{supp}(R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}})) = \mathrm{Im}(\zeta)$.

We can now finish the proof of the fact that Φ is an isomorphism. According to Claim 4, we get that Φ is an isomorphism if and only if its shifted pullback $\zeta^*\Phi[g]$ is an isomorphism. By definition of Φ and using Claim 2, the shifted pull-back

$$\zeta^*\Phi[g] : \zeta^* R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) \to \zeta^*\zeta_*(\mathcal{O}_{\mathrm{Spec}\,\mathrm{R}_{\mathrm{X}}}) = \mathcal{O}_{\mathrm{Spec}\,\mathrm{R}_{\mathrm{X}}} \tag{11.19}$$

coincides, up to the shift, with the morphism (11.7). By tracing back the definition of the morphism (11.7), we get that $\zeta^*\Phi[g]$ is the composition of the top degree base change morphism

$$\zeta^* R^g\widetilde{u}_*(\mathcal{P}^{\mathrm{un}}) \to R^g u_*(\widetilde{\zeta}^*(\mathcal{P}^{\mathrm{un}})) = R^g u_*(\mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}) \tag{11.20}$$

with the isomorphism (11.6). However, since u has fibers of dimension g, the top degree base change (11.20) is an isomorphism, hence we are done.

□

Lemma 11.2. *Let Y be a Noetherian scheme and $\mathcal{F}$ a coherent sheaf on Y. Assume that $\mathcal{F}$ is Cohen-Macaulay and that the scheme-theoretic support* $\mathrm{supp}(\mathcal{F})$ *of $\mathcal{F}$ is generically reduced. Then* $\mathrm{supp}(\mathcal{F})$ *is reduced.*

Proof. The statement is clearly local; hence we may assume that $Y = \mathrm{Spec}\,\mathrm{R}$ with R a Noetherian ring and that $\mathcal{F}$ is equal to the sheafification of a finitely generated module M over R. Therefore, $\mathrm{supp}(\mathcal{F})$ is the closed subscheme $V(\mathrm{ann}(M))$ of $\mathrm{Spec}\,\mathrm{R}$ defined by the annihilator ideal $\mathrm{ann}(M)$ of M. Consider the set $\mathrm{Ass}(M) := \{P_1, \cdots, P_r\}$ of associated primes of M. Since M is a Cohen-Macaulay module, all its associated primes are minimal by [Mat89, Thm. 17.3(i)]; therefore $\{P_1, \ldots, P_r\}$ are exactly the associated minimal primes of $R/\mathrm{ann}(M)$ by [Mat89, Thm. 6.5(iii)].

Consider now a finite set of generators $\{m_1, \cdots, m_s\}$ of the R-module M. Clearly, we have that

$$\mathrm{ann}(M) = \bigcap_{i=1}^{s} \mathrm{ann}(m_i), \tag{11.21}$$

where $\mathrm{ann}(m_i)$ is the annihilator ideal of the element $m_i \in M$.

Since we have an inclusion $R/\mathrm{ann}(m_i) \hookrightarrow M$ of R-modules obtained by sending the class of 1 to m_i, the set of associated primes of $R/\mathrm{ann}(m_i)$ is a subset of $\mathrm{Ass}(M)$; say $\mathrm{Ass}(R/\mathrm{ann}(m_i)) = \{P_j \,:\, j \in A_i\}$ for some $A_i \subseteq \{1, \cdots, r\}$. In particular, $R/\mathrm{ann}(m_i)$ does not have embedded primes. Moreover, since $V(\mathrm{ann}(m_i)) \subseteq V(\mathrm{ann}(M))$ and $V(\mathrm{ann}(M))$ is generically reduced along the subvarieties $V(P_i)$ by hypothesis, it follows that $V(\mathrm{ann}(m_i))$ is also generically reduced. This implies that $V(\mathrm{ann}(m_i))$ is reduced, or in other words that

$$\mathrm{ann}(m_i) = \bigcap_{j\in A_i} P_j. \tag{11.22}$$

Combining (11.21) and (11.22), together with the fact that $\{P_1, \cdots, P_r\} \subseteq \mathrm{Ass}(R/\mathrm{ann}(M))$, we get that

$$\mathrm{ann}(M) = \bigcap_{i=1}^{r} P_i, \tag{11.23}$$

which shows that $V(\mathrm{ann}(M))$ is a reduced subscheme of $\mathrm{Spec}\,\mathrm{R}$. □

The formula (11.3) established in Theorem 11.1 allows us to prove Theorem A from the introduction using a criterion of fully faithfulness for integral functors established by A. C. López Martín in Appendix B.

Proof of Theorem A. Let

$$\begin{aligned} \Phi^{\mathcal{P}}_{J(X)\to \overline{J}_X(\underline{q})} = \Phi^{\mathcal{P}} : D^b(J(X)) & \longrightarrow & D^b(\overline{J}_X(\underline{q})) \\ \mathcal{E}^\bullet & \longmapsto & \mathbf{R}p_{1*}(p_2^*(\mathcal{E}^\bullet)\otimes \mathcal{P}) \end{aligned}$$

be the Fourier-Mukai functor (or integral functor) with kernel the Poincaré line bundle $\mathcal{P}$ on $\overline{J}_X(\underline{q}) \times J(X)$. In order to show that $\Phi^{\mathcal{P}}$ is fully faithfull, we will use Theorem 14.6 from Appendix B. Observe that $J(X)$ is smooth and connected (see §1.9) and that $\overline{J}_X(\underline{q})$ is connected by Corollary 4.4 and Gorenstein by Corollary 2.12(i). Let us check that the three assumptions of Theorem 14.6 are satisfied.

Observe first that, from the definition of $\Phi^{\mathcal{P}}$, it follows that

$$\Phi^{\mathcal{P}}(\mathbf{k}(M)) = \mathbf{R}p_{1*}(p_2^*(\mathbf{k}(M))\otimes\mathcal{P}) = \mathbf{R}p_{1*}(\mathcal{P}_{|\overline{J}_X(\underline{q})\times\{M\}}) = \mathcal{P}_M, \tag{11.24}$$

where $\mathbf{k}(M)$ denotes the structure sheaf of the point $M \in J(X)$.

Using (11.3), (11.24) and Proposition 8.1, we get for any $M \neq M' \in J(X)$ and any i that:

$$\mathrm{Hom}^i_{D^b(\overline{J}_X(\underline{q}))}(\Phi^{\mathcal{P}}(\mathbf{k}(M)), \Phi^{\mathcal{P}}(\mathbf{k}(M'))) = \mathrm{Ext}^i_{\overline{J}_X(\underline{q})}(\mathcal{P}_M, \mathcal{P}_{M'}) = H^i(\overline{J}_X(\underline{q}), \mathcal{P}_{M'\otimes M^{-1}}) = 0, \tag{11.25}$$

which shows that condition (1) of Theorem 14.6 is satisfied.

Similarly, using moreover that $\overline{J}_X(\underline{q})$ is reduced by Corollary 2.12, we have, for any $M \in J(X)$, that:

$$\mathrm{Hom}^0_{D^b(\overline{J}_X(\underline{q}))}(\Phi^{\mathcal{P}}(\mathbf{k}(M)), \Phi^{\mathcal{P}}(\mathbf{k}(M))) = \mathrm{Hom}_{\overline{J}_X(\underline{q})}(\mathcal{P}_M, \mathcal{P}_M) = H^0(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = k, \tag{11.26}$$

which shows that condition (2) of Theorem 14.6 is satisfied.

In order to show condition (3) of Theorem 14.6, we will prove that

$$\left(\Phi^{\mathcal{P}^{-1}\otimes p_1^*(\omega_{\overline{J}_X(\underline{q})})[g]}_{\overline{J}_X(\underline{q})\to J(X)} \circ \Phi^{\mathcal{P}}_{J(X)\to\overline{J}_X(\underline{q})}\right)(\mathbf{k}(0)) = \mathbf{k}(0), \tag{11.27}$$

where $g := p_a(X)$ is the dimension of $\overline{J}_X(\underline{q})$ (see Corollary 2.12(ii)) and $0 = [\mathcal{O}_X]$ is the origin of $J(X)$. From (11.24) and Proposition 8.1, we get that

$$\Phi^{\mathcal{P}}_{J(X)\to\overline{J}_X(\underline{q})}(\mathbf{k}(0)) = \mathcal{O}_{\overline{J}_X(\underline{q})}. \tag{11.28}$$

Observe now that $\omega_{\overline{J}_X(\underline{q})} = \mathcal{O}_{\overline{J}_X(\underline{q})}$ by Corollary 4.5. Moreover, if we denote by $\imath$ the involution of $J(X)$ that sends M into M^{-1}, then Proposition 8.1 implies that $\mathcal{P}^{-1} = (\mathrm{id}\times \mathrm{i})^*(\mathcal{P})$. Using these two facts and formula (11.3), we now compute

$$\Phi^{\mathcal{P}^{-1}\otimes p_1^*(\omega_{\overline{J}_X(\underline{q})})[g]}_{\overline{J}_X(\underline{q})\to J(X)}(\mathcal{O}_{\overline{J}_X(\underline{q})}) = \mathbf{R}p_{2*}((\mathrm{id}\times\mathrm{i})^*(\mathcal{P})[\mathrm{g}]) = \mathrm{i}^*(\mathbf{R}\mathrm{p}_{2*}(\mathcal{P}))[\mathrm{g}] = \mathrm{i}^*(\mathbf{k}(0)[-\mathrm{g}])[\mathrm{g}] = \mathbf{k}(0). \tag{11.29}$$

Combining (11.28) and (11.29), equality (11.27) follows and therefore condition (3) of Theorem 14.6 holds true.

Therefore, the three conditions of Theorem 14.6 of Appendix B are satisfied and Theorem A follows. ☐

Corollary B follows now quite easily from Theorem 11.1 and Theorem A.

Proof of Corollary B. Let $M \in J(X)$.

If $M \neq [\mathcal{O}_X]$ then the vanishing of $H^i(\overline{J}_X(\underline{q}), \mathcal{P}_M)$ for any i follows by (11.3).

If $M = [\mathcal{O}_X] = 0$ is the identity element of $J(X)$ then $\mathcal{P}_M = \mathcal{O}_{\overline{J}_X(\underline{q})}$ by Proposition 8.1. Using formula (11.24) and the fully faithfulness of the Fourier-Mukai transform $\Phi^{\mathcal{P}}$ (see Theorem A), we get

$$H^i(\overline{J}_X(\underline{q}), \mathcal{O}_{\overline{J}_X(\underline{q})}) = \mathrm{Ext}^i_{\overline{J}_X(\underline{q})}(\mathcal{O}_{\overline{J}_X(\underline{q})}, \mathcal{O}_{\overline{J}_X(\underline{q})}) = \mathrm{Ext}^i_{\overline{J}_X(\underline{q})}(\mathcal{F}_{\mathcal{P}}(\mathbf{k}(0)), \mathcal{F}_{\mathcal{P}}(\mathbf{k}(0))) = \mathrm{Ext}^i_{J(X)}(\mathbf{k}(0), \mathbf{k}(0)),$$

where $\mathbf{k}(0)$ is the skyscraper sheaf supported at $0 = [\mathcal{O}_X] \in J(X)$.

Now we conclude using the well-known facts that $\mathrm{Ext}^i_{J(X)}(\mathbf{k}(0), \mathbf{k}(0)) = \Lambda^i \mathrm{Ext}^1_{J(X)}(\mathbf{k}(0), \mathbf{k}(0))$ (see e.g. the proof of [Mum70, Cor. 2, p. 129]) and that $\mathrm{Ext}^1_{J(X)}(\mathbf{k}(0), \mathbf{k}(0))$ is canonically isomorphic to the tangent space of $J(X)$ at $\mathbf{k}(0)$, which is isomorphic to $H^1(X, \mathcal{O}_X)$ (see e.g. [BLR90, Sec. 8.4]). ☐

We can now prove that autoduality holds for fine compactified Jacobians.

Proof of Theorem C. Theorem C follows from Corollary 10.4, whose hypothesis is satisfied by Corollary B.

☐

12. Proof of Theorem D

The aim of this section is to prove Theorem D from the introduction. We will fist prove the result for fine compactified Jacobians that admit an Abel map, in the sense of Definition 6.8.

Theorem 12.1. *Let X be a reduced curve with locally planar singularities and let $\underline{q}$ be a general polarization on X such that the associated fine compactified Jacobian $\overline{J}_X(\underline{q})$ admits an Abel map, i.e. there exists $L \in \mathrm{Pic}^{|\underline{q}|+1}(X)$ such that $\mathrm{Im}\mathrm{A}_{\mathrm{L}} \subseteq \overline{\mathrm{J}}_{\mathrm{X}}(\underline{\mathrm{q}})$. Then we have that*

$$\mathrm{Pic}^o(\overline{J}_X(\underline{q})) = \mathrm{Pic}^\tau(\overline{J}_X(\underline{q})).$$

Proof. Arguing as in the proof of Theorem 4.2, we can find a line bundle $\mathcal{L}$ on the effective semiuniversal deformation $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$ of X such that $\mathcal{L}$ restricted to the central fiber of π is equal to L. A slight extension of the arguments used in §6.1 and in §6.2 allows us to define a global Abel map $A_{\mathcal{L}} : \mathcal{X} \to \overline{\mathbb{J}}_{\mathcal{X}}$ over $\mathrm{Spec}\, \mathrm{R_X}$ whose restriction to the fiber over $s \in \mathrm{Spec}\, \mathrm{R_X}$ is the Abel map $A_{\mathcal{L}_s} : \mathcal{X}_s \to \overline{\mathbb{J}}_{\mathcal{X}_s}$ associated to $\mathcal{L}_s := \mathcal{L}_{|\mathcal{X}_s}$. In particular, the restriction of $A_{\mathcal{L}}$ to the closed point $[\mathfrak{m}_X] \in \mathrm{Spec}\, \mathrm{R_X}$ is equal to the Abel map A_L. Since A_L takes values in $\overline{J}_X(\underline{q})$ by hypothesis, it is easily checked that $A_{\mathcal{L}}$ takes values in $\overline{J}_{\mathcal{X}}(\underline{q})$, or in other words we get a global Abel map

$$A_{\mathcal{L}} : \mathcal{X} \to \overline{J}_{\mathcal{X}}(\underline{q}). \tag{12.1}$$

The pull-back morphism $A_{\mathcal{L}}^* : \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q})) \to \mathrm{Pic}(\mathcal{X}) = \mathbb{J}_{\mathcal{X}}$ between the two relative Picard schemes (whose existence is guaranteed by Fact 3.6(i) and Theorem 5.1(i)) will clearly sends $\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$, which exists by Theorem 5.1(iv) and Corollary B, into the universal generalized Jacobian $J(\mathcal{X}) = \mathrm{Pic}^o(\mathcal{X})$ of $\mathcal{X}$, which exists by Fact 4.6. Moreover, the proof of Proposition 8.5 easily extends to the global case and gives that the induced homomorphism $A_{\mathcal{L}}^{*,o} : \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to J(\mathcal{X})$ is a right inverse of the homomorphism $\beta_{\underline{q}}^{\mathrm{un}} : J(\mathcal{X}) \to \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$ defined in (10.2). Since $\beta_{\underline{q}}^{\mathrm{un}}$ is an isomorphism by Theorem 10.2 and Corollary B, we get that $A_{\mathcal{L}}^* : \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to J(\mathcal{X})$ is an isomorphism of group schemes over $\mathrm{Spec}\, \mathrm{R_X}$.

The pull-back morphism $A_{\mathcal{L}}^*$ will also sends $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))$, which exists by Theorem 5.1(ii), into the generalized Jacobian $J(\mathcal{X})$ of $\mathcal{X}$ since it is well-known that for the family of curves $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$ we have that $\mathrm{Pic}^\tau(\mathcal{X}) = \mathrm{Pic}^o(\mathcal{X}) = J(\mathcal{X})$ (see e.g. [FGA05, Chap. 9.6, Ex. 9.6.21]). Therefore, we get that the induced homomorphism $A_{\mathcal{L}}^{*,\tau} : \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})) \to J(\mathcal{X})$ is a surjective homomorphism of $\mathrm{Spec}\, \mathrm{R_X}$-group schemes.

Summing up this discussion, we get the following diagram of homomorphisms of group schemes over $\mathrm{Spec}\, \mathrm{R_X}$:

$$\begin{tikzcd}
\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})) \arrow[dr, "A_{\mathcal{L}}^{*,\tau}"] & \\
& J(\mathcal{X}) = \mathrm{Pic}^\tau(\mathcal{X}) = \mathrm{Pic}^o(\mathcal{X}) \\
\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \arrow[uu, hook, "i"] \arrow[ur, "\cong", "A_{\mathcal{L}}^{*,o}"'] &
\end{tikzcd} \tag{12.2}$$

where i is an open embedding between two smooth group schemes over $\mathrm{Spec}\ \mathrm{R_X}$ (as it follows from Theorem 5.1 and Corollary B).

Consider now the open subset $U \subseteq \mathrm{Spec}\, \mathrm{R_X}$ (introduced in Lemma 3.5) consisting of all the points s such that the fiber $\mathcal{X}_s$ of the universal family $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$ over s is smooth or it has a unique singular point which is a node. By Lemma 3.5, the complement of U inside $\mathrm{Spec}\, \mathrm{R_X}$ has codimension at least two.

CLAIM 1: The restriction of $A_{\mathcal{L}}^{*,\tau}$ to U

$$(A_{\mathcal{L}}^{*,\tau})_{|U} : \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))_{|U} \to J(\mathcal{X})_{|U}$$

is an isomorphism. In particular, $A_{\mathcal{L}}^{*,\tau}$ is an isomorphism in codimension one.

Indeed, using the above diagram (12.2), it is enough to prove that the open embedding i is an isomorphism over U or, in other words, that:

$$\mathrm{Pic}^o(\overline{J}_{\mathcal{X}_s}(\underline{q}^s)) = \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}_s}(\underline{q}^s)) \text{ for any } s \in U. \tag{12.3}$$

By the definition of U, the fiber $\mathcal{X}_s$ over a point $s \in U$ can be of three types:

(i) $\mathcal{X}_s$ is smooth;

(ii) $\mathcal{X}_s$ has two smooth irreducible components $\mathcal{X}_s^1$ and $\mathcal{X}_s^2$ which meet in a separating node.

(iii) $\mathcal{X}_s$ is an irreducible curve having a unique singular point which is a node;

In case (i), $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ is an abelian variety and the equality (12.3) is proved for abelian varieties by Mumford in [Mum70, Cor. 2, p. 178]. Also in case (ii), $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ is an abelian variety by Proposition 6.6 and the result follows again from the above mentioned result of Mumford. In case (iii), the equality (12.3) is due to Esteve-Gagné-Kleiman [EGK00], where the same result is proved for integral curves with at worst double points.

CLAIM 2: $A_{\mathcal{L}}^{*,\tau}$ is an isomorphism.

Indeed, since $A_{\mathcal{L}}^{*,\tau}$ is a birational map between two integral schemes which is an isomorphism in codimension one (by Claim 1) and the codomain is normal and locally factorial (being smooth by Theorem 3.7(ii) and Fact 4.6), we deduce that $A_{\mathcal{L}}^{*,\tau}$ is a local isomorphism by Van der Waerden's theorem on the purity of the ramification locus (see [EGAIV4, (21.12.12)]). In particular, $A_{\mathcal{L}}^{*,\tau}$ is quasi-finite. Moreover, since $A_{\mathcal{L}}^{*,\tau}$ is birational (by Claim 1) and separated (which follows from the fact that $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})) \to \mathrm{Spec}\, \mathrm{R_X}$ is separated by Theorem 5.1(ii), see [Har77, Cor. 4.6(e)]) and the codomain is normal (being smooth), we deduce that $A_{\mathcal{L}}^{*,\tau}$ is an open embedding by Zariski's main theorem (see [EGAIII1, (4.4.9)]). Since we know that $A_{\mathcal{L}}^{*,\tau}$ is surjective, we deduce that $A_{\mathcal{L}}^{*,\tau}$ is an isomorphism, q.e.d.

From diagram (12.2) and Claim 2, we deduce that the open embedding i must be an equality. By considering the central fiber of i, the theorem now follows.

□

We can now prove the general case of Theorem D.

Proof of Theorem D. Consider an arbitrary fine compactified Jacobian $\overline{J}_X(\underline{q})$ of the curve X. Since $\mathrm{Pic}^o(\overline{J}_X(\underline{q}))$ is an open subscheme of $\mathrm{Pic}^\tau(\overline{J}_X(\underline{q}))$ and they are both defined over an algebraically closed field k, in order to prove that they are equal, it is sufficient (and necessary) to prove that they have the same k-points, i.e. that

$$\mathrm{Pic}^o(\overline{J}_X(\underline{q}))(k) = \mathrm{Pic}^\tau(\overline{J}_X(\underline{q}))(k). \tag{12.4}$$

Consider now the schemes $\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$ and $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))$, which are smooth over $\mathrm{Spec}\, \mathrm{R_X}$ by Theorem 5.1 and Corollary B. Since $S := \mathrm{Spec}\, \mathrm{R_X}$ is henselian (because R_X is a complete ring), the canonical reduction maps $\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \to \mathrm{Pic}^o(\overline{J}_X(\underline{q}))(k)$ and $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \to \mathrm{Pic}^\tau(\overline{J}_X(\underline{q}))(k)$ from the set of S-valued points to the set of k-valued points are surjective by [BLR90, Sec. 3.2, Prop. 5]. Therefore, in order to show the equality (12.4), it is enough to show that

$$\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S) = \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S). \tag{12.5}$$

Observe that we have the following natural inclusions

$$\mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \subseteq \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \subseteq \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(S). \tag{12.6}$$

Moreover, since $\mathrm{Pic}\, S = \mathrm{Pic}(\mathrm{Spec}\, \mathrm{R_X}) = 0$ because R_X is a power series ring and the morphism $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to S$ admits a section passing through its smooth locus (see Theorem 4.3) $J_{\mathcal{X}}(\underline{q}) \to S$ by [BLR90, Sec. 2.2, Prop. 4], we have a natural identification (see [BLR90, Sec. 8.1, Prop. 4])

$$\mathcal{P}ic(\overline{J}_{\mathcal{X}}(\underline{q})) \xrightarrow{\cong} \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(S), \tag{12.7}$$

where, as usual (see 1.8), we denote by $\mathcal{P}ic(\overline{J}_{\mathcal{X}}(\underline{q}))$ the group of line bundles on $\overline{J}_{\mathcal{X}}(\underline{q})$.

Consider now the open subscheme $\overline{J}_{\mathcal{X}}(\underline{q})_{|U} := u^{-1}(U) \subseteq \overline{J}_{\mathcal{X}}(\underline{q})$, where U is the open subset of $\mathrm{Spec} \mathrm{R_X}$ consisting of those points $s \in \mathrm{Spec}\, \mathrm{R_X}$ such that $\mathcal{X}_s$ has at most a unique singular point which is a node (as in Lemma 3.5). The complement of $\overline{J}_{\mathcal{X}}(\underline{q})_{|U}$ inside $\overline{J}_{\mathcal{X}}(\underline{q})$ has codimension at least two by Lemma 3.5. Since $\overline{J}_{\mathcal{X}}(\underline{q})$ is a smooth, irreducible and separated scheme by Theorem 4.3, we can apply [Har77, Prop. 6.5, Cor. 6.16] in order to conclude that the natural restriction map induces an isomorphism

$$\mathcal{P}ic(\overline{J}_{\mathcal{X}}(\underline{q})) \xrightarrow{\cong} \mathcal{P}ic(\overline{J}_{\mathcal{X}}(\underline{q})_{|U}). \tag{12.8}$$

By combining (12.7) and (12.8), we get that the following natural restriction map of sections is an isomorphism

$$\mathrm{res} : \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \xrightarrow{\cong} \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(U) = \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q})_{|U})(U). \tag{12.9}$$

It is clear that $\operatorname{res}(\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S)) \subseteq \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})_{|U})(U)$ and similarly that $\operatorname{res}(\operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S)) \subseteq \operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q})_{|U})(U)$.

Consider any general polarization $\underline{q}'$ on X such that the associated fine compactified Jacobian $\overline{J}_X(\underline{q}')$ admits an Abel map in the sense of Definition 6.8. Note that there are plenty of such general polarizations due to Corollary 6.9(ii). Then the inclusions (12.6) and the isomorphism (12.9) hold true also for the polarization $\underline{q}'$. Moreover, Theorem 12.1 (see also its proof) implies that

$$\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}')) = \operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}')). \tag{12.10}$$

Lemma 12.2 below implies that $\overline{J}_{\mathcal{X}}(\underline{q})_{|U} \cong \overline{J}_{\mathcal{X}}(\underline{q}')_{|U}$; we can therefore set

$$\overline{J}_U := \overline{J}_{\mathcal{X}}(\underline{q})_{|U} \cong \overline{J}_{\mathcal{X}}(\underline{q}')_{|U}. \tag{12.11}$$

Combining (12.6) and (12.9) (and their analogues for $\underline{q}'$) together with (12.10) and (12.11), we get the following commutative diagram of (abstract) abelian groups

$$\begin{tikzcd}
\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \arrow[r, hook] \arrow[d] & \operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \arrow[r, hook] \arrow[d] & \operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \arrow[d, "\operatorname{res}", "\cong"'] \\
\operatorname{Pic}^o(\overline{J}_U)(U) \arrow[r, equal] & \operatorname{Pic}^\tau(\overline{J}_U)(U) \arrow[r, hook] & \operatorname{Pic}(\overline{J}_U)(U) \\
\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}'))(S) \arrow[r, equal] \arrow[u] & \operatorname{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}'))(S) \arrow[r, hook] \arrow[u] & \operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}'))(S) \arrow[u, "\operatorname{res}'"', "\cong"]
\end{tikzcd} \tag{12.12}$$

CLAIM: For any section $\sigma \in \operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(S)$, if we set $\sigma' := (\operatorname{res}')^{-1}(\operatorname{res}(\sigma)) \in \operatorname{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}'))(S)$, then we have that

$$\sigma \in \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \Longleftrightarrow \sigma' \in \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}'))(S). \tag{12.13}$$

Let us prove the implication $\Rightarrow$ (the other being analogous). We apply Lemma 12.3 below to the group scheme morphism $\pi : \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to S$ (which is smooth by Theorem 5.1(iv) and has geometrically connected fibers by construction), endowed with the sections $\tau_1 := \sigma$ and $\tau_2 := \underline{0}$, where $\underline{0}$ is the zero section of π. Notice that the sections τ_1 and τ_2 are contained in an affine open subset of $\operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))$ since the central fiber $\operatorname{Pic}^o(\overline{J}_X(\underline{q}))$ of π is a k-group scheme, hence quasi-projective (see e.g. [BLR90, Sec. 6.4, Thm. 1]). From Lemma 12.3 together with the fact that every section of $\pi_V : S(B) \times_S \operatorname{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q})) \to S(B)$ (for any affine k-variety $V = \operatorname{Spec} \mathrm{B}$) is represented by a (possibly non unique) line bundle on $S(B) \times_S \overline{J}_{\mathcal{X}}(\underline{q})$ by [BLR90, Sec. 8.1, Prop. 4] (using that π has a section, as observed before), we get the existence of two line bundles L_1 and L_2 on $S(k[t]) \times_S \overline{J}_{\mathcal{X}}(\underline{q})$ and of a line bundle L_Z on $S(R) \times_S \overline{J}_{\mathcal{X}}(\underline{q})$ such that

1. $L = L_{1|f_{\mathbb{A}^1}^{-1}(1) \times_S \overline{J}_{\mathcal{X}}(\underline{q})}$,
2. $L_{1|f_{\mathbb{A}^1}^{-1}(0) \times_S \overline{J}_{\mathcal{X}}(\underline{q})} = L_{Z|f_Z^{-1}(q_1) \times_S \overline{J}_{\mathcal{X}}(\underline{q})}$,
3. $L_{Z|f_Z^{-1}(q_2) \times_S \overline{J}_{\mathcal{X}}(\underline{q})} = L_{2|f_{\mathbb{A}^1}^{-1}(0) \times_S \overline{J}_{\mathcal{X}}(\underline{q})}$,
4. $L_{2|f_{\mathbb{A}^1}^{-1}(1) \times_S \overline{J}_{\mathcal{X}}(\underline{q})} = \mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q})}$,

where L is the line bundle on $\overline{J}_{\mathcal{X}}(\underline{q})$ corresponding to the section σ under the bijection (12.7).

Since $S(k[t])$ and $S(R)$ are smooth and $\overline{J}_{\mathcal{X}}(\underline{q})$ and $\overline{J}_{\mathcal{X}}(\underline{q}')$ are isomorphic in codimension 1 by Lemma 12.2, arguing as before we get two line bundles L_1' and L_2' on $S(k[t]) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')$ and a line bundle L_Z' on $S(R) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')$ such that

(1′) $L' = L'_{1|f_{\mathbb{A}^1}^{-1}(1) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')}$,
(2′) $L'_{1|f_{\mathbb{A}^1}^{-1}(0) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')} = L'_{Z|f_Z^{-1}(q_1) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')}$,
(3′) $L'_{Z|f_Z^{-1}(q_2) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')} = L'_{2|f_{\mathbb{A}^1}^{-1}(0) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')}$,
(4′) $L'_{2|f_{\mathbb{A}^1}^{-1}(1) \times_S \overline{J}_{\mathcal{X}}(\underline{q}')} = \mathcal{O}_{\overline{J}_{\mathcal{X}}(\underline{q}')}$,

where L' is the line bundle on $\overline{J}_{\mathcal{X}}(\underline{q}')$ corresponding to the section σ' under the analogous of the bijection (12.7) for $\overline{J}_{\mathcal{X}}(\underline{q})$. Restricting the four isomorphisms (1′)–(4′) to the fiber $\overline{J}_{\mathcal{X}_s}(\underline{q}'^s)$ of $u : \overline{J}_{\mathcal{X}}(\underline{q}') \to \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$ over the point $s \in \operatorname{Spec} \mathrm{R}_{\mathrm{X}}$, we deduce that $L'_{|\overline{J}_{\mathcal{X}_s}(\underline{q}'^s)}$ is algebraically equivalent to the trivial line bundle $\mathcal{O}_{\overline{J}_{\mathcal{X}_s}(\underline{q}'^s)}$ (because $\mathbb{A}^1$ and Z are connected varieties). This is equivalent to the fact that $\sigma'(s) \in \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}'^s))$ for every $s \in \mathrm{Spec}\, \mathrm{R_X}$, or in other words that $\sigma' \in \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}'))$, which concludes the proof of the Claim.

Let us now conclude the proof of Theorem D. From [Kle71, Thm. 5.1], it follows that there exists a number N such that any section $\sigma \in \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S)$ is such that $\sigma^N \in \mathrm{Pic}^o(\overline{J}_{\mathcal{X}}(\underline{q}))(S)$ and similarly for the sections in $\mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}'))(S)$. Therefore, from the commutative diagram (12.12) and (12.13), we deduce that for a section $\sigma \in \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}))(S)$, if we set $\sigma' := (\mathrm{res}')^{-1}(\mathrm{res}(\sigma)) \in \mathrm{Pic}(\overline{J}_{\mathcal{X}}(\underline{q}'))(S)$ as before, then we have that

$$\sigma \in \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}))(S) \Longleftrightarrow \sigma' \in \mathrm{Pic}^\tau(\overline{J}_{\mathcal{X}}(\underline{q}'))(S). \tag{12.14}$$

An easy chasing in diagram (12.12) together with (12.13) and (12.14) shows that the required equality (12.5) does hold true and this concludes the proof of Theorem D.

□

The following result, which was used in the above proof, allows to compare two different universal fine compactified Jacobians of X over the open subset $U \subset \mathrm{Spec}\, \mathrm{R_X}$ considered in Lemma 3.5.

Lemma 12.2. *Let $\underline{q}$ and $\underline{q}'$ be two general polarizations on a curve X with locally planar singularities. Let U be the open subset of* $\mathrm{Spec}\, \mathrm{R_X}$ *consisting of those points $s \in \mathrm{Spec}\, \mathrm{R_X}$ such that $\mathcal{X}_s$ has at most a unique singular point which is a node (as in Lemma 3.5). Consider the induced universal fine compactified Jacobians $u : \overline{J}_{\mathcal{X}}(\underline{q}) \to \mathrm{Spec}\, \mathrm{R_X}$ and $u' : \overline{J}_{\mathcal{X}}(\underline{q}') \to \mathrm{Spec}\, \mathrm{R_X}$ (as in Theorem 4.2), and set $\overline{J}_{\mathcal{X}}(\underline{q})_{|U} := u^{-1}(U)$ and $\overline{J}_{\mathcal{X}}(\underline{q}')_{|U} := (u')^{-1}(U)$. Then there exists a line bundle $\mathcal{L}$ on $\mathcal{X}$ such that the multiplication by $\mathcal{L}$ induces an isomorphism of schemes over U:*

$$- \otimes \mathcal{L} : \overline{J}_{\mathcal{X}}(\underline{q})_{|U} \xrightarrow{\cong} \overline{J}_{\mathcal{X}}(\underline{q}')_{|U}.$$

Proof. Let $Y_1, \ldots, Y_r$ be the separating blocks of X as in §6.2. By Proposition 6.6(iii), we can assume without loss of generality that $\underline{q}$ and $\underline{q}'$ are induced by general polarizations $\underline{q}^i$ and $\underline{q}'^i$ on Y_i for $i = 1, \ldots, r$. This implies that the multidegree $\underline{d} := \underline{q}' - \underline{q}$ on X is such that $\underline{d}_{Y_i} \in \mathbb{Z}$ for every $i = 1, \ldots, r$. Take now a line bundle $L_i \in \mathrm{Pic}^{\underline{d}_{Y_i}}(Y_i)$ for every $i = 1, \ldots, r$ and consider the unique line bundle L on X such that $L_{|Y_i} = L_i$ (see e.g. Proposition 6.6(i)). With the same argument used in the proof of Theorem 4.2, we can find a line bundle $\mathcal{L}$ on the total space $\mathcal{X}$ of the semiuniversal deformation $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$ such that the restriction of $\mathcal{L}$ to the central fiber X of π is isomorphic to L. Clearly, the multiplication by $\mathcal{L}$ induces an isomorphism

$$- \otimes \mathcal{L} : \overline{\mathbb{J}}_{\mathcal{X}} \xrightarrow{\cong} \overline{\mathbb{J}}_{\mathcal{X}},$$

the inverse being given by the multiplication by $\mathcal{L}^{-1}$. Since any universal fine compactified Jacobian is an open subscheme of $\overline{\mathbb{J}}_X$, in order to conclude the proof it is enough to prove that for any $s \in U$

$$(- \otimes \mathcal{L}_s)(\overline{J}_{\mathcal{X}_s}(\underline{q}^s)) = \overline{J}_{\mathcal{X}_s}(\underline{q}'^s) \tag{12.15}$$

where $\mathcal{L}_s$ denotes the restriction of $\mathcal{L}$ to the fiber $\mathcal{X}_s$ and $\overline{J}_{\mathcal{X}_s}(\underline{q}^s)$ (resp. $\overline{J}_{\mathcal{X}_s}(\underline{q}'^s)$) is the fiber of $\overline{J}_{\mathcal{X}}(\underline{q})$ (resp. $\overline{J}_{\mathcal{X}}(\underline{q}')$) over s (see Theorem 4.2). If $\mathcal{X}_s$ is smooth or irreducible with one node, then (12.15) follows simply from the fact that $|\underline{d}| + |\underline{q}| = |\underline{q}'|$. Suppose now that $\mathcal{X}_s$ has a separating node, i.e. $\mathcal{X}_s$ consists of two smooth curves Z_1 and Z_2 joined in a separating node p. Since p will specialize to a separating node of the central fiber X of $\pi : \mathcal{X} \to \mathrm{Spec}\, \mathrm{R_X}$, there exists a subset $\emptyset \neq J_1 \subsetneq [r] := \{1, \ldots, r\}$ with complementary subset $J_2 := [r] \setminus J_1$ such that, with respect to the specialization map Σ_s of (4.1), we have (for $i = 1, 2$)

$$\Sigma_s(Z_i) = \overline{Z_i} \cap X = \bigcup_{k \in J_i} Y_k.$$

Using Lemma-Definition 4.1, we get for $i = 1, 2$

$$\deg \mathcal{L}_{s|Z_i} = \sum_{k \in J_i} \deg L_{|Y_k} \in \mathbb{Z}, \qquad \underline{q}^s_{Z_i} = \sum_{k \in J_i} \underline{q}_{Y_k} \in \mathbb{Z}, \qquad \underline{q}'^s_{Z_i} = \sum_{k \in J_i} \underline{q}'_{Y_k} \in \mathbb{Z},$$

which implies, by our choice of the multidegree $\underline{d}$, that

$$\deg \mathcal{L}_{s|Z_i} + \underline{q}^s_{Z_i} = \underline{q}'^s_{Z_i}. \tag{12.16}$$

Using Proposition 6.6(iii), we get that the pull-back via the normalization morphism $\tau : Z_1 \coprod Z_2 \to \mathcal{X}_s$ induces the following isomorphisms

$$\overline{J}_{\mathcal{X}_s}(\underline{q}^s) \xrightarrow[\cong]{\tau^*} \prod_{i=1}^{2} \mathrm{Pic}^{\underline{q}^s_{Z_i}}(Z_i) \quad \text{and} \quad \overline{J}_{\mathcal{X}_s}(\underline{q}'^s) \xrightarrow[\cong]{\tau^*} \prod_{i=1}^{2} \mathrm{Pic}^{\underline{q}'^s_{Z_i}}(Z_i). \tag{12.17}$$

Combining (12.16) and (12.17), we get that (12.15) does hold also in the case when $\mathcal{X}_s$ has a separating node, q.e.d. □

The following technical result is a crucial ingredient in the above proof of Theorem D.

Lemma 12.3. *Let* $S = \mathrm{Spec\,k[[x_1, \dots, x_r]]}$, *let* $\overline{0}$ *be the unique closed point of* S *and let* $\pi : \mathcal{Y} \to S$ *be a surjective smooth morphism such that* $Y := \pi^{-1}(\overline{0})$ *is connected. Let* τ_1 *and* τ_2 *be two sections of* π *and assume that there exists an open affine subset* $\mathcal{Z} = \mathrm{Spec\,(A)} \subseteq \mathcal{Y}$ *containing* $q_1 := \tau_1(\overline{0})$ *and* $q_2 := \tau_2(\overline{0})$.

For any affine k*-variety* $V = \mathrm{Spec\,(B)}$, *set* $S(B) := \mathrm{Spec\,B[[x_1, \dots, x_r]]}$ *and let* $f_V : S(B) \to V$ *be the map induced by the natural inclusion* $B \subset B[[x_1, \dots, x_r]]$. *Let* $\pi_V : S(B) \times_S \mathcal{Y} \to S(B)$ *be the base change map of* π *via the natural map* $S(B) \to k[[x_1, \dots, x_r]]$.

Denote by $Z = \mathrm{Spec\,(R)}$ *the smooth affine connected variety* $\mathcal{Z} \cap Y$, *by* $\mathbb{A}^1$ *the affine line* $\mathrm{Spec\,(k[t])}$ *over* k *and let* 0 *(resp.* 1*) be the point of* $\mathbb{A}^1$ *defined by the ideal* (t) *(resp.* $(t-1)$*).*

Then there exist two sections $\tau_{i,\mathbb{A}^1}$ *(for* $i = 1, 2$*) of* $\pi_{\mathbb{A}^1}$ *and a section* τ_Z *of* π_Z *such that*

(1) $\tau_1 = \tau_{1,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(1)}$,
(2) $\tau_{1,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(0)} = \tau_{Z|f_Z^{-1}(q_1)}$,
(3) $\tau_{Z|f_Z^{-1}(q_2)} = \tau_{2,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(0)}$,
(4) $\tau_{2,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(1)} = \tau_2$.

Proof. Let $I \subset A$ be the ideal of $\mathrm{Spec(R)}$ in A (i.e. $A/I = R$). As $S = \mathrm{Speck[[x_1, \dots, x_r]]}$, any section τ of π factors through $\mathrm{Spec(\hat{A})}$ where $\hat{A}$ is the I-adic completion of A. Moreover, as $\mathrm{Spec(R)}$ is a smooth affine variety, it is rigid and its infinitesimal deformations are trivial. As a consequence, $A/I^n = \frac{A/I[x_1,\dots,x_r]}{(x_1,\dots,x_r)^n}$ for any $n \in \mathbb{N}$ and $\hat{A} = A/I[[x_1, \dots, x_r]] = R[[x_1, \dots, x_r]]$. So it will suffice to prove the Lemma assuming that $\mathcal{Y} = \mathrm{Spec\,(R[[x_1, \dots, x_r]])}$ and that $\pi : S(R) = \mathrm{Spec\,(R[[x_1, \dots, x_r]])} \to \mathrm{S} = \mathrm{Spec\,k[[x_1, \dots, x_r]]}$ is the map induced by the natural embedding $k[[x_1, \dots, x_r]] \subset R[[x_1, \dots, x_r]]$. Under these assumptions, we are going to define $\tau_{i,\mathbb{A}^1}$ and τ_Z such that

(1) $\tau_1 = \tau_{1,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(1)}$,
(2) $\tau_{1,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(0)} = \tau_{Z|f_Z^{-1}(q_1)} =$ constant section q_1,
(3) $\tau_{Z|f_Z^{-1}(q_2)} = \tau_{2,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(0)} =$ constant section q_2,
(4) $\tau_{2,\mathbb{A}^1|f_{\mathbb{A}^1}^{-1}(1)} = \tau_2$.

We define $\tau_Z : S(R) \to S(R) \times_S S(R)$ as the diagonal embedding. This definition implies that $\tau_{Z|f_Z^{-1}(q)}$ is the constant section whose value is q for every $q \in Z$.

To define $\tau_{i,\mathbb{A}^1} : S(k[t]) \to S(k[t]) \times_S S(R)$ (for $i = 1, 2$) notice that $\tau_i : S \to S(R)$ is induced by a morphism $g_i : R[[x_1, \dots, x_r]] \to k[[x_1, \dots, x_r]]$ sending the ideal J_i of q_i to the maximal ideal of $k[[x_1, \dots, x_r]]$. Therefore g_i factors through the J_i-adic completion $\widehat{R[[x_1, \dots, x_r]]}$ of $R[[x_1, \dots, x_r]]$. Since Z and π are smooth there exists a $k[[x_1, \dots, x_r]]-$algebra isomorphism $\widehat{R[[x_1, \dots, x_r]]} \simeq k[[x_1, \dots, x_s]]$ for some $s > r$. Summing up, there exist $k[[x_1, \dots, x_r]]-$algebra morphisms $\phi_i : R[[x_1, \dots, x_r]] \to k[[x_1, \dots, x_s]]$ and $\overline{g}_i : k[[x_1, \dots, x_s]] \to k[[x_1, \dots, x_r]]$ such that $g_i = \overline{g}_i \circ \phi_i$. Using these morphisms we define

$$\phi_{i,\mathbb{A}^1} : k[t][[x_1, \dots, x_r]] \otimes_{k[[x_1,\dots,x_r]]} R[[x_1, \dots, x_r]] \to k[t][[x_1, \dots, x_r]] \otimes_{k[[x_1,\dots,x_r]]} k[[x_1, \dots, x_s]]$$

as the unique $k[t][[x_1, \dots, x_r]]-$morphism induced by ϕ_i and

$$\begin{aligned} &\overline{g}_{i,\mathbb{A}^1} : k[t][[x_1, \dots, x_r]] \otimes_{k[[x_1,\dots,x_r]]} k[[x_1, \dots, x_s]] \longrightarrow k[t][[x_1, \dots, x_r]], \\ &a(t, x_1, \dots, x_r) \otimes b(x_1, \dots, x_r, x_{r+1}, \dots, x_s) \mapsto a(t, x_1, \dots, x_r) \otimes \overline{g}_i(b)(x_1, \dots, x_r, tx_{r+1}, \dots, tx_s). \end{aligned}$$

Finally, the desired map $\tau_{i,\mathbb{A}^1}$ is obtained as the map induced on spectra by the composition $\overline{g}_{i,\mathbb{A}^1} \circ \phi_{i,\mathbb{A}^1}$. □

13. Appendix A: Hitchin fibration vs compactified Jacobians of spectral curves

Let C be a fixed connected smooth and projective curve of genus g over an algebraically closed field k and let L be a line bundle on C (often it is convenient to assume that L has high degree, e.g. $\deg L \geq 2g-2$). Fix a natural number $r \geq 1$ and an integral number $d \in \mathbb{Z}$.

An *L-twisted Higgs pair* (or simply a Higgs pair when L is clear from the context) on C is a pair (E, ϕ) consisting of a vector bundle E on C and a homomorphism $\phi : E \to E \otimes L$ (called the Higgs field). The degree (resp. the rank) of a Higgs pair (E, ϕ) is the degree $\deg E$ (resp. the rank $\operatorname{rk} E$) of the underlying vector bundle E. In the important special case when $L = \omega_C$, an ω_C-twisted Higgs pair is simply called a *Higgs bundle.*

The algebraic stack $\mathcal{M} = \mathcal{M}(r, d, L)$ of all L-twisted Higgs pairs (E, ϕ) on C of rank r and degree d is endowed with a morphism (called the *Hitchin morphism*)

$$\begin{aligned} \mathcal{H} : \mathcal{M}(r,d,L) = \mathcal{M} &\longrightarrow \mathcal{A} = \mathcal{A}(r,L) := \oplus_{i=1}^{r} H^0(C, L^i) \\ (E,\phi) &\mapsto \mathcal{H}(E,\phi) := (a_1(E,\phi), \ldots, a_r(E,\phi)), \end{aligned} \tag{13.1}$$

where $a_i(E, \phi) := (-1)^i \mathrm{Tr}(\Lambda^i \phi) \in H^0(C, L^i)$.

The algebraic (Artin) stack $\mathcal{M}$ is not of finite type. In order to obtain a space of finite type (and indeed a variety), one introduces a semistability condition as follows. A Higgs pair (E, ϕ) is called *semistable* (resp. *stable*) if for all non-trivial proper subsheaves $F \subsetneq E$ that are stable with respect to ϕ (i.e. such that $\phi(F) \subseteq F \otimes L$) we have

$$\frac{\deg F}{\operatorname{rk} F} \leq \frac{\deg E}{\operatorname{rk} E} \quad (\text{resp. } <).$$

Observe that, given a Higgs pair (E, ϕ), if E is a semistable (resp. stable) vector bundle then (E, ϕ) is semistable (resp. stable) but the converse is in general false.

The coarse moduli space $M = M(r, d, L)$ of S-equivalence classes of semistable L-twisted Higgs pairs (E, ϕ) of rank r and degree d has been constructed by N. Hitchin [Hit86] using analytic methods (namely gauge theory) and later by N. Nitsure [Nit91] using algebro-geometric methods (namely geometric invariant theory). The Hitchin fibration (13.1) restricts to a projective fibration (called the *Hitchin fibration*):

$$\begin{aligned} H : M(r,d,L) = M &\longrightarrow \mathcal{A} = \mathcal{A}(r,L) := \oplus_{i=1}^{r} H^0(C, L^i) \\ (E,\phi) &\mapsto (a_1(E,\phi), \ldots, a_r(E,\phi)). \end{aligned} \tag{13.2}$$

Remark 13.1. In the special case where $L = \omega_C$, r and d are coprime (so that there are no strictly semistable Higgs pairs) and $k = \mathbb{C}$, it is known that (see [Hit86]):

- $M = M(r, d, \omega_C)$ is an hyperkähler (non compact) manifold containing, as an open subset, the cotangent bundle of the moduli space of stable (= semistable) vector bundles on C of degree d and rank r;
- H is an algebraically completely integrable Hamiltonian system.

The fibers of the Hitchin morphism $\mathcal{H}$ and of the Hitchin fibration H can be described in terms of compactified Jacobians of spectral curves, as we are now going to explain. Consider the $\mathbb{P}^1$-fibration $p : P = \mathbb{P}(\mathcal{O}_C \oplus L^*) \to C$ and let $\mathcal{O}(1)$ be the relatively ample line bundle on P. We will denote by y the section of $\mathcal{O}(1)$ whose pushforward via p corresponds to the constant section $(1, 0)$ of the vector bundle $p_*\mathcal{O}(1) = \mathcal{O}_C \oplus L^*$. Similarly, we will denote by x the section of $\mathcal{O}(1) \otimes p^*(L)$ whose pushforward via p corresponds to the constant section $(0, 1)$ of the vector bundle $p_*(\mathcal{O}(1) \otimes p^*(L)) = (\mathcal{O}_C \oplus L^*) \otimes L = L \oplus \mathcal{O}_C$. Given $\underline{a} = (a_1, \ldots, a_r) \in \oplus_{i=1}^{r} H^0(C, L^i) = \mathcal{A}$, the *spectral curve* $C_{\underline{a}}$ associated to $\underline{a}$ is the projective (but possibly singular) curve inside P given as the zero locus of the following section of $\mathcal{O}(1) \otimes p^*(L)^n$:

$$x^r + p^*(a_1) \cdot x^{r-1} \cdot y + \ldots + p^*(a_r) \cdot y^n.$$

Remark 13.2. The spectral curve $C_{\underline{a}}$ can be very singular (although it has locally planar singularities because it is embedded in the smooth surface P), and in particular it is not necessarily reduced nor irreducible. The base $\mathcal{A}$ of the Hitchin morphism admits two notably open (non-empty if L is sufficiently positive, e.g. if L^r is very ample) subsets $\mathcal{A}^{\mathrm{ell}} \subseteq \mathcal{A}^{\mathrm{reg}} \subseteq \mathcal{A}$, called respectively the elliptic locus and the regular locus, defined as follows:

$$\mathcal{A}^{\mathrm{ell}} := \{\underline{a} \in \mathcal{A} \,:\, C_{\underline{a}} \text{ is integral}\},$$
$$\mathcal{A}^{\mathrm{reg}} := \{\underline{a} \in \mathcal{A} \,:\, C_{\underline{a}} \text{ is reduced}\}.$$

The study of the topology of the Hitchin morphism restricted to the elliptic locus $\mathcal{A}^{\mathrm{ell}}$ has played a crucial role in B. C. Ngô's proof of the fundamental lemma (see [Ngo06] and [Ngo10]) and, more generally, the study of the Hitchin morphism over the regular locus $\mathcal{A}^{\text{reg}}$ was a crucial ingredient in Chaudouard-Laumon's proof of the weighted fundamental lemma (see [CL10] and [CL12]).

The restriction of the morphism $p : P \to C$ to $C_{\underline{a}}$ is a degree-r finite morphism $\pi_{\underline{a}} : C_{\underline{a}} \to C$, called the *spectral cover* associated to $\underline{a} \in \mathcal{A}$. Note that, since the zero sets of x and y are disjoint in P, the restriction of the section y to $C_{\underline{a}}$ is everywhere non-zero, which implies that $\mathcal{O}(1)_{|C_{\underline{a}}} = \mathcal{O}_{C_{\underline{a}}}$. Therefore, the restriction $x_{|C_{\underline{a}}}$ of the section x to $C_{\underline{a}}$ can be considered as a section of $[p^*(L) \otimes \mathcal{O}(1)]_{|C_{\underline{a}}} = p^*(L)_{|C_{\underline{a}}} = \pi_{\underline{a}}^*(L)$.

Fact 13.3 (Beauville-Narashiman-Ramanan [BNR89], Shaub [Sch98]). *Let $\underline{a} \in \mathcal{A} = \oplus_{i=1}^r H^0(C, L^i)$ and consider the associated degree-r spectral cover $\pi_{\underline{a}} : C_{\underline{a}} \to C$.*

(i) There is a bijective correspondence

$$\Pi : \left\{ \begin{matrix} \textit{Torsion-free rank-}1 \textit{ sheaves } I \textit{ on } C_{\underline{a}} \\ \textit{of degree } d + \binom{r}{2} \deg L \end{matrix} \right\} \longrightarrow \left\{ \begin{matrix} \textit{Higgs pairs } (E, \phi) \textit{ on } C \textit{ of degree } d \textit{ and rank } r \\ \textit{such that } \mathcal{H}(E, \phi) = \underline{a} \end{matrix} \right\}$$

obtained by associating to a torsion-free rank-1 sheaf I on $C_{\underline{a}}$ the Higgs pair $\Pi(I) = (E, \phi)$ on C consisting of the vector bundle $E := (\pi_{\underline{a}})_*(I)$ on C together with the Higgs field $\phi : (\pi_{\underline{a}})_*(I) \to (\pi_{\underline{a}})_*(I) \otimes L = (\pi_{\underline{a}})_*(I \otimes \pi_{\underline{a}}^*(L))$ given by the multiplication with $x_{|C_{\underline{a}}} \in H^0(C_{\underline{a}}, \pi_{\underline{a}}^*(L))$.

(ii) Given a torsion-free rank-1 sheaf I on $C_{\underline{a}}$, then $\Pi(I)$ is a semistable (resp. stable) Higgs pair on C if and only if I is semistable (resp. stable) with respect to the polarization $\underline{q}^{\underline{a}}$ on $C_{\underline{a}}$ given by

$$\underline{q}^{\underline{a}}_Z := \frac{\deg I}{r} \deg(\pi_{\underline{a}|Z}) \quad \textit{for any subcurve} \quad Z \subseteq C_{\underline{a}}. \tag{13.3}$$

(iii) Assume that $\underline{a} \in \mathcal{A}^{\text{reg}}$ (i.e. $C_{\underline{a}}$ is a reduced curve) and that $\underline{q}^{\underline{a}}$ is a general polarization in the sense of Definition 2.6 (i.e. every torsion-free rank-1 sheaf on $C_{\underline{a}}$ which is $\underline{q}^{\underline{a}}$-semistable is also $\underline{q}^{\underline{a}}$-stable). Then the bijective correspondence Π gives rise to an isomorphism

$$\overline{J}_{C_{\underline{a}}}(\underline{q}^{\underline{a}}) \xrightarrow[\cong]{\Pi} H^{-1}(\underline{a}), \tag{13.4}$$

where $\overline{J}_{C_{\underline{a}}}(\underline{q}^{\underline{a}})$ is Esteves's fine compactified Jacobian of the reduced curve $C_{\underline{a}}$ with respect to the general polarization $\underline{q}^{\underline{a}}$ (see Section 2.2).

Proof. Part (i) is proved in [BNR89, Prop. 3.6] under the hypothesis that the spectral curve $C_{\underline{a}}$ is integral and in [Sch98, Prop. 2.1] for an arbitrary spectral curve.

Part (ii) is a reformulation in our language of [Sch98, Thm. 3.1].

Part (iii) follows from (ii) and the fact that the bijective correspondence Π of (i) does hold in families as well (see [Sch98, Prop. 5.1]).

□

Remark 13.4. Note that if d and r are coprime, then for every $\underline{a} \in \mathcal{A}$ the polarization $\underline{q}^{\underline{a}}$ on $C_{\underline{a}}$ is general since there are no strictly semistable Higgs pairs, hence no strictly semistable torsion-free rank-1 sheaves on $C_{\underline{a}}$ by Fact 13.3(ii).

In the general case, Chaudouard-Laumon [CL10] have introduced, after taking a suitable cover of $\mathcal{A}^{\text{reg}}$, alternative semistablity conditions on the stack $\mathcal{M}$ of Higgs pairs, for which there are no strictly semistable objects. The moduli spaces defined by these new semistability conditions are such that the fibers of the associated Hitchin fibrations are always isomorphic to Esteves's fine compactified Jacobians.

The Hitchin fibration (13.2) in the case $L = \omega_C$ is conjectured to satisfy the following autoduality property, which can be viewed as a "classical limit" of the (conjectural) geometric Langlands correspondence (see [DP12, Sec. 2] for a discussion of the geometric Langlands correspondence and the passage to the classical limit).

Conjecture 13.5 (Langlands duality for the Hitchin fibration). *Let $M = \mathcal{M}(d, r, \omega_C)$ be the moduli space of S-equivalence classes of Higgs bundles of rank r and degree d and let $H : M \to \mathcal{A}$ be the Hitchin fibration (13.2). Then there exists a universal Poincaré sheaf $\mathcal{P}$ on $M \times_{\mathcal{A}} M$ such that the Fourier-Mukai transform with kernel $\mathcal{P}$*

$$\Phi^{\mathcal{P}} : D^b(M) \to D^b(M)$$

is an auto-equivalence of the bounded derived category $D^b(M)$ of coherent sheaves on M.

More precisely, the above auto-equivalence $\Phi^{\mathcal{P}}$ is expected to intertwine the action of the classical limit tensorization functors with the action of the classical limit Hecke functors (see [DP12, Conj. 2.5] for a precise formulation). In loc. cit., a more general Langlands duality is conjectured to hold between the moduli space of G-Higgs bundles (for G any reductive group) and the moduli space of ${}^LG$-Higgs bundles, where ${}^LG$ is the Langlands dual of G. Conjecture 13.5 is the special case in which $G = \mathrm{GL}_r = {}^LG$. In [DP12, Cor. 5.5], Donagi-Pantev proved Conjecture 13.5 (and its generalized version for any reductive group G) over the open subset of $\mathcal{A}$ where the Hitchin fibration H is smooth, or equivalently over the open subset of $\mathcal{A}$ parametrizing smooth spectral curves.

By restricting to the fibers of the Hicthin fibration (13.2), Conjecture 13.5 should imply the following autoduality result.

Conjecture 13.6 (Fiberwise Langlands duality for the Hitchin fibration)**.** *Same notations as in Conjecture 13.5. For any $\underline{a} \in \mathcal{A}$, the Fourier-Mukai transform with kernel $\mathcal{P}_{\underline{a}} := \mathcal{P}_{|H^{-1}(\underline{a})}$*

$$\Phi^{\mathcal{P}_{\underline{a}}} : D^b(H^{-1}(\underline{a})) \to D^b(H^{-1}(\underline{a}))$$

is an auto-equivalence of the bounded derived category $D^b(H^{-1}(\underline{a}))$ of coherent sheaves on $H^{-1}(\underline{a})$.

Several particular cases of Conjecture 13.6 are known:

- If $\underline{a} \in \mathcal{A}$ is such that $C_{\underline{a}}$ is smooth, then $H^{-1}(\underline{a})$ is isomorphic to the Jacobian of $C_{\underline{a}}$ (by Fact 13.3(iii)) and therefore this boils down to the celebrated result of S. Mukai [Muk81].
- If $\underline{a} \in \mathcal{A}^{\mathrm{ell}}$, i.e. if the associated spectral curve $C_{\underline{a}}$ is integral, then $H^{-1}(\underline{a})$ is isomorphic to the compactified Jacobian of $C_{\underline{a}}$ (by Fact 13.3(iii)) and therefore Conjecture 13.6 follows from the work of A. Arinkin [Ari].
- If $\underline{a} \in \mathcal{A}^{\mathrm{reg}}$, i.e. $C_{\underline{a}}$ is reduced, and $\underline{q}^{\underline{a}}$ is a general polarization on $C^{\underline{a}}$ (for example, if d and r are coprime by Remark 13.4), then $H^{-1}(\underline{a})$ is isomorphic to a fine compactified Jacobian of $C_{\underline{a}}$ (by Fact 13.3(iii)) and therefore Conjecture 13.6 follows from Theorem E of the introduction (whose proof will appear in [MRV2]).

14. Appendix B: Fully faithfulness criteria (by Ana Cristina López Martín)

Given a scheme X quasi-projective over an algebraically closed field k, denote by $D(X)$ the derived category of complexes of $\mathcal{O}_X$-modules with quasi-coherent cohomology sheaves and by $D^b(X) \subset D(X)$ its bounded derived subcategory. We will use the subscript c to refer to the corresponding subcategory of complexes of $\mathcal{O}_X$-modules with coherent cohomology sheaves.

Given another scheme Y quasi-projective over k, consider the following commutative diagram

$$\begin{tikzcd} & X \times Y \arrow[dl, "p_1"'] \arrow[dr, "p_2"] & \\ X \arrow[dr, "p"'] & & Y \arrow[dl, "q"] \\ & \mathrm{Spec}\,\mathrm{k} & \end{tikzcd}$$

If $\mathcal{K}^\bullet$ is an object in $D(X \times Y)$, the *integral functor* with kernel $\mathcal{K}^\bullet$ is the functor

$$\begin{aligned} \Phi^{\mathcal{K}^\bullet}_{X\to Y} \colon D(X) &\to D(Y) \\ \mathcal{E}^\bullet &\mapsto \mathbf{R}p_{2_*}(p_1^*\mathcal{E}^\bullet \overset{\mathbf{L}}{\otimes} \mathcal{K}^\bullet)\,. \end{aligned}$$

In the case that the kernel $\mathcal{K}^\bullet$ is a line bundle $\mathcal{P}$ on $X \times Y$, the functor $\Phi^{\mathcal{P}}_{X\to Y}$ maps $D^b(X)$ to $D^b(Y)$. Observe that, if p_2 is not a proper morphism (as in the case we are interested in), an integral functor might not preserve coherence so that it is important to work with D (or D^b) instead of D_c (or D^b_c).

The aim of this Appendix is to prove two fully faithful criteria for integral functors, one in characteristic zero and a variant of it valid in arbitrary characteristic; the second criterion is used in the proof of Theorem A. Before stating the criteria, let us collect some lemmas that will be used in the proof.

Lemma 14.1. *Let X and Y be quasi-projective k-schemes. Suppose that X is smooth and connected and Y is projective connected and Gorenstein of dimension n and denote by ω_Y its dualizing sheaf. Let $\mathcal{P}$ be a line bundle on $X \times Y$ and denote by $\mathcal{P}^{-1}$ its dual line bundle. Then, the integral functor*

$$\Phi^{\mathcal{P}^{-1}\otimes p_2^*\omega_Y[n]}_{Y\to X} \colon D^b(Y) \to D^b(X)$$

is a left adjoint to the functor $\Phi^{\mathcal{P}}_{X\to Y}\colon D^b(X)\to D^b(Y)$.

Proof. There is a chain of isomorphisms

$$\begin{aligned}\mathrm{Hom}_{\mathrm{D^b(Y)}}(\mathcal{G}^\bullet,\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathcal{F}^\bullet)) &\simeq \mathrm{Hom}_{\mathrm{D^b(X\times Y)}}(\mathbf{L}\mathrm{p}_2^*\mathcal{G}^\bullet,\mathrm{p}_1^*\mathcal{F}^\bullet\otimes\mathcal{P})\simeq\\ &\simeq \mathrm{Hom}_{\mathrm{D^b(X\times Y)}}(\mathbf{L}\mathrm{p}_2^*\mathcal{G}^\bullet\otimes\mathcal{P}^*,\mathrm{p}_1^*\mathcal{F}^\bullet)\simeq\\ &\simeq \mathrm{Hom}_{\mathrm{D^b(X\times Y)}}(\mathbf{L}\mathrm{p}_2^*\mathcal{G}^\bullet\otimes\mathcal{P}^*\otimes\mathrm{p}_2^*\omega_{\mathrm{Y}}[\mathrm{n}],\mathrm{p}_1^!\mathcal{F}^\bullet)\simeq\\ &\simeq \mathrm{Hom}_{\mathrm{D^b(X)}}(\Phi^{\mathcal{P}^{-1}\otimes\mathrm{p}_2^*\omega_{\mathrm{Y}}[\mathrm{n}]}_{\mathrm{Y\to X}}(\mathcal{G}^\bullet),\mathcal{F}^\bullet).\end{aligned}$$

where the first isomorphism is the adjunction formula between the derived inverse and the derived direct images (see for instance [BBH08, Prop A.80]), the second is [Har66, Proposition 5.16], the third follows from (C.2) in [BBH08] taking into account that Y is Gorenstein of dimension n, and the forth is obtained by applying Grothendieck's duality to the projective morphism $p_1\colon X\times Y\to X$. □

For any closed point $x\in X$ denote by $\mathbf{k}(x)$ the skyscraper sheaf supported at x. Notice that it is a perfect complex if X is a smooth scheme.

Lemma 14.2. *Let* $j\colon Y\hookrightarrow X$ *be a closed immersion of smooth irreducible quasi-projective* k*-schemes of dimensions* n *and* m*, respectively, and set* $d:=m-n$*. Let* $0\neq\mathcal{K}^\bullet\in D^b_c(X)$ *be a bounded complex with coherent cohomology sheaves. For any closed point* $x\in X$ *denote by* $j_x\colon\{x\}\hookrightarrow X$ *the natural immersion. Then, if*

$$\mathbf{L}^i j_x^*\mathcal{K}^\bullet\simeq\mathrm{Hom}^{\mathrm{m-i}}_{\mathrm{D(X)}}(\mathbf{k}(\mathrm{x}),\mathcal{K}^\bullet)=0$$

unless $x\in Y$ *and* $i\in[0,d]$*, then* $\mathcal{K}^\bullet\simeq\mathcal{K}$ *is a sheaf on* X *and* $\operatorname{supp}\mathcal{K}=Y$ *(topologically).*

Proof. This follows from Proposition 1.5 of [BO95] (see also Proposition 1.6 in [San09]). □

The following lemma was proved in [Bri99] for projective schemes. The proof here is similar.

Lemma 14.3. *Let* S *and* X *be quasi-projective* k*-schemes of finite type, and let* $\mathcal{F}$ *be a coherent sheaf on* $S\times X$ *flat over* S *and schematically supported on a closed scheme* $Z\hookrightarrow S\times X$*. Suppose that for every closed point* $s\in S$*, the restriction* $\mathcal{F}_s$ *of* $\mathcal{F}$ *to the fiber* X_s *is topologically supported at a single point* $x\in X$ *and that*

$$\mathrm{Hom_X}(\mathcal{F}_{\mathrm{s}},\mathbf{k}(\mathrm{x}))\simeq\mathrm{k}\,.$$

Then $\mathcal{F}\simeq j_*\mathcal{L}$ *for a line bundle* $\mathcal{L}$ *on* Z*.*

Moreover, if $\operatorname{char}(k)=0$ *and for all pairs of distinct points* $s_1,s_2\in S$ *the sheaves* $\mathcal{F}_{s_1}$ *and* $\mathcal{F}_{s_2}$ *are not isomorphic, then the Kodaira-Spencer map of the family* $\mathcal{F}$ *is injective at some point* $s\in S$*.*

Proof. Let us see first that for any closed point $s\in S$ the sheaf $\mathcal{F}_s$ is the structure sheaf of a zero dimensional scheme with support at x. For this it is enough to prove that there is a surjective morphism $\mathcal{O}_X\to\mathcal{F}_s$. Indeed, consider non-zero morphisms $g\colon\mathcal{F}_s\to\mathbf{k}(x)$ and $f\colon\mathcal{O}_X\to\mathcal{F}_s$. If the morphism f is not surjective, its cokernel is a non-zero sheaf topologically supported at x. Then, there is a non-zero morphism $\operatorname{Coker}f\to\mathbf{k}(x)$ that induces another non-zero morphism $h\colon\mathcal{F}_s\to\mathbf{k}(x)$ such that $h\circ f=0$. Since $\mathrm{Hom_X}(\mathcal{F}_{\mathrm{s}},\mathbf{k}(\mathrm{x}))\simeq\mathrm{k}$, one has that $g\circ f=0$ so that any non surjective morphism f takes values on $\operatorname{Ker}g$. However, since we have the following exact sequence of sheaves supported at x

$$0\to\operatorname{Ker}g\to\mathcal{F}_s\xrightarrow{g}\mathbf{k}(x)\to 0$$

there is some $f\in H^0(\mathcal{F}_s)$ such that $f\notin H^0(\operatorname{Ker}g)$. The corresponding morphism $f\colon\mathcal{O}_X\to\mathcal{F}_s$ is such that $g\circ f\neq 0$, and thus it is surjective.

Let $s\in S$ be a closed point. Since $\mathcal{F}_s$ is topologically supported at x, we may assume, up to shrinking S, that S is affine and that the support of $\mathcal{F}$ is included in $V\times S$ for a suitable affine neighborhood V of $x\in X$. Since $V\times S$ is affine and the natural map of sheaves $\mathcal{F}\to\mathcal{F}_s$ is surjective, then the restriction morphism $H^0(X\times S,\mathcal{F})=H^0(V\times S,\mathcal{F})\to H^0(V\times S,\mathcal{F}_s)=H^0(X_s,\mathcal{F}_s)$ is also surjective. Therefore, there exists a global section σ of $\mathcal{F}$ mapping to f. For a suitable open neighborhood U of s, $\sigma_U\colon\mathcal{O}_U\to\mathcal{F}_U$ is still surjective, that is $\mathcal{F}_U\simeq\mathcal{O}_{Z_U}$, where $Z_U=Z\cap(U\times X)$, which proves that $\mathcal{F}$ is a line bundle on its support.

Taking into account that the Hilbert scheme $\operatorname{Hilb}(X)$ of zero-dimensional subschemes of X exists also when X is a quasi-projective scheme (see [FGAIV]), the proof of the second statement is equal to that of Lemma 1.23 in [BBH08]: the sheaf $\mathcal{F}_U$ induces a map $\alpha\colon U\to\operatorname{Hilb}^P(X)$ to the Hilbert scheme of zero-dimensional subschemes of X whose Hilbert polynomial P is equal to that of the sheaf $\mathcal{F}_s$. The morphism α is characterized by the property $\mathcal{F}\simeq(\alpha\times\mathrm{id})^*\mathcal{Q}$ where $\mathcal{Q}$ is the universal sheaf on $\mathrm{Hilb}^P(X)\times X$. By the hypothesis, α is injective on closed points and, since the base field k has characteristic zero, the tangent map $T_s\alpha\colon T_sU\to T_{\alpha(s)}\mathrm{Hilb}^P(X)$ is injective at least at one point $s\in S$. Moreover, the Kodaira-Spencer map $KS_s(\mathcal{F})$ at s for $\mathcal{F}_U$ is the composition of the tangent map $T_s\alpha$ with the Kodaira-Specer map for the universal family $\mathcal{Q}$. Since the later is an isomorphism because of the universality of $\mathcal{Q}$, we conclude that $KS_s(\mathcal{F})$ is injective. □

Lemma 14.4. *Let X and Y be quasi-projective k-schemes and let $\mathcal{F}$ be a coherent sheaf on $X\times Y$ flat over X. Let x be a closed point of X and set $\mathcal{F}_x:=\mathcal{F}_{|\{x\}\times Y}$. Then the associated integral functor $\Phi=\Phi^{\mathcal{F}}_{X\to Y}$ induces a morphism*

$$\Phi\colon \mathrm{Ext}^1_{\mathrm{X}}(\mathbf{k}(\mathrm{x}),\mathbf{k}(\mathrm{x}))\to \mathrm{Ext}^1(\Phi(\mathbf{k}(\mathrm{x})),\Phi(\mathbf{k}(\mathrm{x})))=\mathrm{Ext}^1_{\mathrm{X}}(\mathcal{F}_{\mathrm{x}},\mathcal{F}_{\mathrm{x}})$$

which coincides with the Kodaira-Specer morphism for the family $\mathcal{F}$.

Proof. The proof is the same as in Lemma 4.4. of [Bri99]. □

The main technical result of this appendix is the following criterion (valid in characteristic zero) for an integral functor to be fully faithful. Its proof follows the line of Bondal-Orlov's criterion (see Theorem 1.1 of [BO95]) and its subsequent generalizations: Theorem 1.22 in [HLS07], Theorems 3.6 and 3.8 [HLS09] and Theorem 2.3 in [San09].

Theorem 14.5. *Let X and Y be two schemes quasi-projective over an algebraically closed field k such that* $\mathrm{char}(k)=0$. *Suppose that X is smooth and connected and Y is projective connected and Gorenstein. Let $\mathcal{P}$ be a line bundle on $X\times Y$. Suppose that the following conditions are satisfied:*

(1) $\mathrm{Hom}^{\mathrm{i}}_{\mathrm{D^b(Y)}}(\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x})),\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{y}))=0$ *for any $x,y\in X$, unless $x=y$ and $0\le i\le\dim X$.*

(2) $\mathrm{Hom}^{0}_{\mathrm{D^b(Y)}}(\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x})),\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x}))=\mathrm{k}$ *for any $x\in X$.*

Then, the integral functor $\Phi^{\mathcal{P}}_{X\to Y}\colon D^b(X)\to D^b(Y)$ is fully faithful.

Proof. By Lemma 14.1, $\Phi=\Phi^{\mathcal{P}}_{X\to Y}$ has a left adjoint $G=\Phi^{\mathcal{P}^{-1}\otimes p_2^*\omega_Y[n]}_{Y\to X}$, where n is the dimension of Y. By Proposition 1.18 in [HLS07], to prove that Φ is fully faithful it is sufficient (and necessary) to prove that the composition $G\circ\Phi$ is fully faithful. This composition is still an integral functor; let us denote by $\mathcal{M}^\bullet\in D^b(X\times X)$ its kernel.

The strategy of the proof is as follows: first we are going to prove that condition (1) in the statement implies that the complex $\mathcal{M}^\bullet$ is isomorphic to a single sheaf $\mathcal{M}$ topologically supported on the image Δ of the diagonal immersion $\delta\colon X\hookrightarrow X\times X$. Next, using condition (2) in the statement, we show that $\mathcal{M}$ is the push-foward of a line bundle $\mathcal{N}$ on a closed scheme Z of $X\times X$. Then we prove that Z coincides with Δ. So, the functor $\Phi^{\mathcal{M}^\bullet}:=\Phi^{\mathcal{M}^\bullet}_{X\to X}$ is given by tensoring by $\mathcal{N}$ and then it is fully faithful.

(a) *$\mathcal{M}^\bullet$ is a single sheaf $\mathcal{M}$ topologically supported on the image Δ of the diagonal morphism $\delta\colon X\hookrightarrow X\times X$.*

Indeed, by the convolution formula, one has that

$$\mathcal{M}^\bullet=\mathbf{R}\pi_{13,*}(\pi_{12}^*\mathcal{P}\otimes\pi_{23}^*(\mathcal{P}^{-1}\otimes p_2^*\omega_Y[n]))$$

where π_{12},π_{23} and π_{13} are the natural projections of $X\times Y\times X$ onto the corresponding factors. Being π_{13} a base change of $q\colon Y\to\operatorname{Spec}\mathrm{k}$, it is a projective morphism so that $\mathbf{R}\pi_{13,*}$ sends coherent sheaves to coherent sheaves, and then the complex $\mathcal{M}^\bullet$ belongs to $D^b_c(X\times X)$ and we can make use of Lemma 14.2.

For any closed point $x\in X$, denote by $j_x\colon\{x\}\hookrightarrow X$ and $i_x\colon\{x\}\times X\hookrightarrow X\times X$ the natural inclusions. Fix a closed point $(x_1,x_2)\in X\times X$. Then, we have

$$\begin{aligned}\mathcal{H}^i(\mathbf{L}j_{x_2}^*\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1))^\vee)&\simeq\mathrm{Hom}^{\mathrm{i}}_{\mathrm{D(\{x_2\})}}(\mathbf{L}\mathrm{j}^*_{\mathrm{x_2}}\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(\mathrm{x_1})),\mathbf{k}(\mathrm{x_2}))\simeq\\&\simeq\mathrm{Hom}^{\mathrm{i}}_{\mathrm{D(X)}}(\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(\mathrm{x_1})),\mathbf{k}(\mathrm{x_2}))\simeq\\&\simeq\mathrm{Hom}^{\mathrm{i}}_{\mathrm{D(Y)}}(\Phi(\mathbf{k}(\mathrm{x_1})),\Phi(\mathbf{k}(\mathrm{x_2})))\,.\end{aligned}$$

The first condition on the hypothesis implies that this is zero unless $x_1=x_2$ and $0\le i\le\dim X$. Since X is a smooth scheme, the skyscraper sheaf $\mathbf{k}(x_2)$ at x_2 is a l.c.i. zero cycle, so that the complex $\mathbf{L}j^*_{x_2}\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1))$ is bounded with coherent cohomology because so is $\mathcal{M}^\bullet$. Being x_2 a zero-dimensional Gorenstein scheme, $\omega_{x_2}\simeq\mathbf{k}(x_2)$ is an injective object and then, using (1.7) in [HLS07], we have

$$\begin{aligned}\mathcal{H}^i(\mathbf{L}j_{x_2}^*\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1))^\vee)&\simeq\mathcal{H}^i(\mathbf{L}j_{x_2}^*\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1))^*)\simeq\\&\simeq\mathcal{H}^{-i}(\mathbf{L}j_{x_2}^*\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1))\simeq\mathbf{L}^ij_{x_2}^*(\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1)))\,.\end{aligned}$$

By applying Lemma 14.2 to the closed immersion $j_{x_1}\colon \{x_1\} \hookrightarrow X$, one has that the complex $\Phi^{\mathcal{M}^\bullet}(\mathbf{k}(x_1)) \simeq \mathbf{L}i_{x_1}^*\mathcal{M}^\bullet$ is a sheaf topologically supported at $\{x_1\}$. Then, using Lemma 4.3 in [Bri99], we get that $\mathcal{M}^\bullet \simeq \mathcal{M}$ where $\mathcal{M}$ is a sheaf on $X \times X$ flat over X by the first projection $\pi_1\colon X \times X \to X$. Moreover, since $\mathbf{L}i_x^*\mathcal{M} \cong i_x^*\mathcal{M}$ is topologically supported on $\{x\}$ for any closed point x of X, the sheaf $\mathcal{M}$ is topologically supported on the diagonal Δ (a closed subscheme of $X \times X$ because X is a separated scheme).

(b) Denote by $\bar{\delta}\colon Z \hookrightarrow X \times X$ the schematic support of $\mathcal{M}$, so that $\mathcal{M} \simeq \bar{\delta}_*\mathcal{N}$ for a coherent sheaf $\mathcal{N}$ on Z. Let us see that $\mathcal{N}$ *is a line bundle.* For every closed point $x \in X$, we know that $i_x^*\mathcal{M} \simeq \Phi^{\mathcal{M}}(\mathbf{k}(x))$ is topologically supported at $\{x\}$ and by the hypothesis

$$\mathrm{Hom_X}(\Phi^{\mathcal{M}}(\mathbf{k}(\mathrm{x})), \mathbf{k}(\mathrm{x})) \simeq \mathrm{Hom}^0_{\mathrm{D^b(Y)}}(\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x})), \Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x}))) = \mathrm{k}\,.$$

By Lemma 14.3, $\mathcal{N}$ is a line bundle.

(c) *The scheme Z coincides with Δ.*

The diagonal embedding $\delta\colon X \hookrightarrow X \times X$ factors through a closed immersion $\tau\colon X \hookrightarrow Z$ which topologically is a homeomorphism and we know that $\mathcal{M} \simeq \bar{\delta}_*\mathcal{N}$ is flat over X by $\pi_1\colon X \times X \to X$. Then $\mathcal{N}$ is flat over X by the composition $\bar{\pi_1} = \pi_1 \circ \bar{\delta}$ and since $\bar{\pi_1}$ is a finite morphism, this implies that $\bar{\pi}_{1*}\mathcal{N} \simeq \pi_{1*}\mathcal{M}$ is a locally free sheaf. In order to show that $Z = \Delta$, it is enough to show that $\pi_{1*}\mathcal{M}$ is a line bundle. Since we know that $\pi_{1*}\mathcal{M}$ is a locally free sheaf and since X is connected, it is enough to show that $\Phi^{\mathcal{M}}(\mathbf{k}(x)) \simeq \mathbf{k}(x)$ for some closed point $x \in X$. Given a closed point $x \in X$, consider the morphism $\Phi^{\mathcal{M}}(\mathbf{k}(x)) \to \mathbf{k}(x) \to 0$ given by adjuntion and let us denote by $\mathcal{K}_x$ is kernel. It suffices to find a closed point $x \in X$ such that $\mathrm{Hom_x}(\mathcal{K}_\mathrm{x}, \mathbf{k}(\mathrm{x})) = 0$. Take homomorphisms in $\mathbf{k}(x)$ from the exact sequence

$$0 \to \mathcal{K}_x \to \Phi^{\mathcal{M}}(\mathbf{k}(x)) \to \mathbf{k}(x) \to 0\,.$$

Since, by Lemma 14.4, the morphism

$$\Phi^{\mathcal{M}} : \mathrm{Hom}^1_{\mathrm{D(X)}}(\mathbf{k}(\mathrm{x}), \mathbf{k}(\mathrm{x})) \to \mathrm{Hom}^1_{\mathrm{D(X)}}(\Phi^{\mathcal{M}}(\mathbf{k}(\mathrm{x})), \Phi^{\mathcal{M}}(\mathbf{k}(\mathrm{x})))$$

is the Kodaira-Spencer map for the family $\mathcal{M}$, it is injective at some point $x \in X$ by Lemma 14.3 (using $\mathrm{char}(k) = 0$) and the result follows. ☐

The hypothesis that $\mathrm{char}(k) = 0$ cannot be removed in the above Theorem 14.5, as it is observed in [HLS07, Rmk. 1.25]. However, if we add one more assumption, then we get the following variant of Theorem 14.5 which is valid in arbitrary characteristic.

Theorem 14.6. *Let X and Y be two schemes quasi-projective over an algebraically closed field k (of arbitrary characteristic). Suppose that X is smooth and connected and that Y is projective connected and Gorenstein of dimension n. Let $\mathcal{P}$ be a line bundle on $X \times Y$ and denote by $\mathcal{P}^{-1}$ its inverse. Suppose that the following conditions are satisfied:*

(1) $\mathrm{Hom}^{\mathrm{i}}_{\mathrm{D^b(Y)}}(\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x})), \Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{y}))) = 0$ *for any $x, y \in X$, unless $x = y$ and $0 \le i \le \dim X$.*

(2) $\mathrm{Hom}^0_{\mathrm{D^b(Y)}}(\Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x})), \Phi^{\mathcal{P}}_{\mathrm{X\to Y}}(\mathbf{k}(\mathrm{x}))) = \mathrm{k}$ *for any $x \in X$.*

(3) There exists a closed point $x \in X$ such that $\left(\Phi^{\mathcal{P}^{-1}\otimes p_2^*\omega_Y[n]}_{Y\to X} \circ \Phi^{\mathcal{P}}_{X\to Y}\right)(\mathbf{k}(x)) \cong \mathbf{k}(x)$.

Then, the integral functor $\Phi^{\mathcal{P}}_{X\to Y}\colon D^b(X) \to D^b(Y)$ is fully faithful.

Proof. In the proof of Theorem 14.5, the only place where the assumption $\mathrm{char}(k) = 0$ is used, is in finding a closed point $x \in X$ such that $\Phi^{\mathcal{M}}(\mathbf{k}(x)) \cong \mathbf{k}(x)$. Since $\Phi^{\mathcal{M}} = \Phi^{\mathcal{P}^{-1}\otimes p_2^*\omega_Y[n]}_{Y\to X} \circ \Phi^{\mathcal{P}}_{X\to Y}$ by definition, the existence of such a closed point x is now guaranteed by the condition (3). ☐

Acknowledgments. We are extremely grateful to E. Esteves for several useful conversations on fine compactified Jacobians and for generously answering several mathematical questions. This project started while the first author was visiting the Mathematics Department of the University of Roma "Tor Vergata" funded by the "Michele Cuozzo" 2009 award. She wishes to express her gratitude to Michele Cuozzo's family and to the Department for this great opportunity. M. Melo was supported by the FCT project *Espaços de Moduli em Geometria Algébrica* (PTDC/MAT/ 111332/2009), by the FCT project *Geometria Algébrica em Portugal* (PTDC/MAT/099275/2008) and by the Fundação Calouste Gulbenkian program "Estímulo à investigação 2010". F. Viviani was supported by the FCT project *Espaços de Moduli em Geometria Algébrica* (PTDC/ MAT/111332/2009).

References

[Ale02] V. Alexeev: *Complete moduli in the presence of semiabelian group action.* Ann. of Math. (2) 155 (2002), no. 3, 611-708.

[Ale04] V. Alexeev: *Compactified Jacobians and Torelli map.* Publ. Res. Inst. Math. Sci. 40 (2004), no. 4, 1241-1265.

[AN99] V. Alexeev, I. Nakamura: *On Mumford's construction of degenerating abelian varieties.* Tohoku Math. J. (2) 51 (1999), no. 3, 399–420.

[AIK76] A. B. Altman, A. Iarrobino, S. L. Kleiman: *Irreducibility of the compactified Jacobian.* Real and complex singularities (Proc. Ninth Nordic Summer School/NAVF Sympos. Math., Oslo, 1976), pp. 1–12. Sijthoff and Noordhoff, Alphen aan den Rijn, 1977.

[AK79] A. B. Altman, S. L. Kleiman: *Bertini theorems for hypersurface sections containing a subscheme.* Comm. Algebra 7 (1979), no. 8, 775–790.

[AK80] A. B. Altman, S. L. Kleiman: *Compactifying the Picard scheme.* Adv. Math. 35 (1980), 50–112.

[AK79] A. B. Altman, S. L. Kleiman: *Compactifying the Picard scheme II.* American Journal of Mathematics 101 (1979), 10–41.

[Ari11] D. Arinkin: *Cohomology of line bundles on compactified Jacobians.* Math. Res. Lett. 18 (2011), no. 06, 1215–1226.

[Ari] D. Arinkin: *Autoduality of compactified Jacobians for curves with plane singularities.* Preprint arXiv:1001.3868v2.

[Art69] M. Artin: *Algebraic approximation of structures over complete local rings.* Inst. Hautes Études Sci. Publ. Math. 36 (1969), 23–58.

[BPV84] W. Barth, C. Peters, A. Van de Ven: *Compact complex surfaces.* Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 4. Springer-Verlag, Berlin, 1984.

[BBH08] C. Bartocci, U. Bruzzo, D. Hernández Ruipérez: *Fourier-Mukai and Nahm transforms in geometry and mathematical physics.* Progress in Mathematics, 276, Birkhäuser Boston Inc., (2009).

[BNR89] A. Beauville, M. S. Narasimhan, S. Ramanan: *Spectral curves and the generalised theta divisor.* J. Reine Angew. Math. 398 (1989), 169–179.

[Ber70] J. E. Bertin: *Generalites sur les preschemas en groupes.* Exposé VIB in SGA3.I "Schémas en groupes I", Lecture Notes in Mathematics 151 (1970), 316–408.

[BO95] A. I. Bondal, D. O. Orlov: *Semi orthogonal decomposition for algebraic varieties.* MPIM Preprint 95/15 (1995), Preprint math.AG/9506012.

[BLR90] S. Bosch, W. Lütkebohmert, M. Raynaud: *Néron models.* Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 21. Springer-Verlag, Berlin, 1990.

[BGS81] J. Briançon, M. Granger, J.P. Speder: *Sur le schéma de Hilbert d'une courbe plane.* Ann. Sci. École Norm. Sup. (4) 14 (1981), no. 1, 1–25.

[Bri99] T. Bridgeland, *Equivalences of triangulated categories and Fourier-Mukai transforms.* Bull. London Math. Soc. 31 (1999), 25–34.

[BH98] W. Bruns, J. Herzog: *Cohen-Macaulay rings.* Cambridge Studies in Advanced Mathematics. 39. Cambridge: Cambridge University Press, 1998.

[Cap94] L. Caporaso: *A compactification of the universal Picard variety over the moduli space of stable curves.* J. Amer. Math. Soc. 7 (1994), 589–660.

[CCE08] L. Caporaso, J. Coelho, E. Esteves: *Abel maps for Gorenstein curves.* Rend. Circ. Mat. Palermo (2) 57 (2008), 33-59.

[CE07] L. Caporaso, E. Esteves: *On Abel maps of stable curves.* Michigan Math. J. 55 (2007), no. 3, 575–607.

[Cat82] F. Catanese: *Pluricanonical-Gorenstein-curves.* Enumerative geometry and classical algebraic geometry (Nice, 1981), pp. 51–95, Progr. Math., 24, Birkhäuser Boston, Boston, MA, 1982.

[CL10] P. H. Chaudouard, G. Laumon: *Le lemme fondamental pondéré. I. Constructions géométriques.* Compos. Math. 146 (2010), no. 6, 1416–1506.

[CL12] P. H. Chaudouard, G. Laumon: *Le lemme fondamental pondéré. II. Énoncés cohomologiques.* Preprint arXiv:0912.4512.

[CP10] J. Coelho, M. Pacini: *Abel maps for curves of compact type.* J. Pure Appl. Algebra 214 (2010), no. 8, 1319–1333.

[DH88] S. Diaz, J. Harris: *Ideals associated to deformations of singular plane curves.* Trans. Amer. Math. Soc. 309 (1988), no. 2, 433–468.

[DP12] R. Donagi, T. Pantev: *Langlands duality for Hitchin systems.* To appear in Invent. Math. (available at arXiv:math/0604617).

[EGAII] A. Grothendieck, J. Dieudonné: *Eléments de Géométrie Algébrique II.* Publ. Math. Inst. Hautes Études Sci. 8 (1961).

[EGAIII1] A. Grothendieck, J. Dieudonné: *Eléments de Géométrie Algébrique III-Part 1.* Publ. Math. Inst. Hautes Études Sci. 11 (1961).

[EGAIV2] A. Grothendieck, J. Dieudoné: *Eléments de Géométrie Algébrique IV-Part 2.* Publ. Math. Inst. Hautes Études Sci. 24 (1965).

[EGAIV3] A. Grothendieck, J. Dieudoné: *Eléments de Géométrie Algébrique IV-Part 3.* Publ. Math. Inst. Hautes Études Sci. 28 (1966).

[EGAIV4] A. Grothendieck, J. Dieudoné: *Eléments de Géométrie Algébrique IV-Part 4.* Publ. Math. Inst. Hautes Études Sci. 32 (1967).

[Est01] E. Esteves: *Compactifying the relative Jacobian over families of reduced curves.* Trans. AMS 353 (2001), 3045–3095.

[Est09] E. Esteves: *Compactified Jacobians of curves with spine decompositions.* Geom. Dedicata 139 (2009), 167–181.

[EGK00] E. Esteves, M. Gagné, S. Kleiman:*Abel maps and presentation schemes.* Special issue in honor of Robin Hartshorne. Comm. Algebra 28 (2000), no. 12, 5961–5992.

[EGK02] E. Esteves, M. Gagné, S. Kleiman: *Autoduality of the compactified Jacobian.* J. Lond. Math. Soc. 65 (2002), No. 3, 591–610.

[EK05] E. Esteves, S. Kleiman: *The compactified Picard scheme of the compactified Jacobian.* Adv. Math. 198 (2005), no. 2, 484–503.

[ER] E. Esteves, F. Rocha: *Autoduality for curves of compact type.* Preprint July 2012.

[FGvS99] B. Fantechi, L. Göttsche, D. van Straten: *Euler number of the compactified Jacobian and multiplicity of rational curves.* J. Algebraic Geom. 8 (1999), no. 1, 115–133.

[FGA05] B. Fantechi, L. Göttsche, L. Illusie, S. L. Kleiman, N. Nitsure, A. Vistoli: *Fundamental algebraic geometry. Grothendieck's FGA explained.* Mathematical Surveys and Monographs, 123. American Mathematical Society, Providence, RI, 2005.

[FGAIV] A. Grothendieck: *Techniques de construction et théorèmes d'existence en géométrie algébrique IV: les schémes de Hilbert.* Séminaire N. Bourbaki, exp. 221 (1960–1961), 249–276.

[FGAVI] A. Grothendieck: *Tecnhique de descente e théorèmes d'existence em géométrie algébriques VI. Le schéma de Picard: propriété générales.* Séminarie N. Bourbaki, exp. 236 (1961–1962), 221–243.

[Har66] R. Hartshorne: *Residues and duality.* Lecture Notes in Mathematics. 20. Berlin-Heidelberg-New York: Springer-Verlag, 1966.

[Har77] R. Hatshorne: *Algebraic Geometry.* Graduate text in Mathematics 52, Springer-Verlag, 1977.

[HLS07] D. Hernández Ruipérez, A. C. López Martín, F. Sancho de Salas: *Fourier-Mukai transform for Gorenstein schemes.* Adv. Math. 211 (2007), 594–620.

[HLS09] D. Hernández Ruipérez, A. C. López Martín, and F. Sancho de Salas: *Relative integral functors for singular fibrations and singular partners.* J. Eur. Math. Soc. (JEMS) 11 (2009), 597–625.

[Hit86] N. Hitchin: *Stable bundles and integrable systems.* Duke Math. J. 54 (1987), no. 1, 91–114.

[Huy06] D. Huybrechts: *Fourier-Mukai transforms in algebraic geometry.* Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006.

[HL97] D. Huybrechts, M. Lehn: *The geometry of moduli spaces of sheaves.* Aspects of Mathematics. E 31. Braunschweig: Vieweg (1997).

[Kas09] J. L. Kass: *Good Completions of Néron Models.* PhD thesis, Harvard, 2009.

[Kas12(a)] J. L. Kass: *Degenerating the Jacobian: the Néron Model versus Stable Sheaves.* To appear in Algebra and Number Theory (available at arXiv:1012.2576).

[Kas12(b)] J. L. Kass: *An Explicit Non-smoothable Component of the Compactified Jacobian.* Preprint available at arXiv:1108.2706.

[Kle71] S. Kleiman: *Les théorémes de finitude pour le foncteur de Picard*, Exposé XIII in SGA 6 "Théorie des intersections et théoréme de Riemann–Roch", Lecture Notes in Math. 225 (1971), 616–666.

[KK81] H. Kleppe, S.L. Kleiman: *Reducibility of the compactified Jacobian.* Compositio Math. 43 (1981), no. 2, 277–280.

[Kle81] H. Kleppe: *The Picard Scheme of a Curve and its Compactification.* PhD thesis, Massachusetts Institute of Technology, Cambridge, Massachusetts, 1981.

[KM76] F. Knudsen, D. Mumford: *The projectivity of the moduli space of stable curves. I: Preliminaries on "det" and "Div".* Math. Scand. 39 (1976), 19–55.

[LM05] A. C. López-Martín: *Simpson Jacobians of reducible curves.* J. Reine Angew. Math. 582 (2005), 1–39.

[Mat89] H. Matsumura: *Commutative ring theory.* Translated from the Japanese by M. Reid. Second edition. Cambridge Studies in Advanced Mathematics, Vol. 8. Cambridge University Press, Cambridge, 1989.

[MRV2] M. Melo, A. Rapagnetta, F. Viviani: *Fourier-Mukai and autoduality for compactified Jacobians. II.* In preparation.

[MV12] M. Melo, F. Viviani: *Fine compactified Jacobians.* Math. Nach. 285 (2012), no. 8-9, 997–1031.

[Muk81] S. Mukai: *Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves.* Nagoya Math. J. 81 (1981), 153–175.

[Mum70] D. Mumford: *Abelian varieties.* Tata Institute of Fundamental Research Studies in Mathematics. 5. London: Oxford University Press. VIII (1970).

[Nit91] N. Nitsure: *Moduli space of semistable pairs on a curve.* Proc. London Math. Soc. (3) 62 (1991), no. 2, 275–300.

[Ngo06] B. C. Ngô: *Fibration de Hitchin et endoscopie.* Invent. Math. 164 (2006), no. 2, 399–453.

[Ngo10] B. C. Ngô: *Le lemme fondamental pour les algèbres de Lie.* Publ. Math. Inst. Hautes Études Sci. No. 111 (2010), 1–169.

[GIT94] J. Fogarty, D. Mumford, F. Kirwan: *Geometric invariant theory.* Third edition. Ergebnisse der Mathematik und ihrer Grenzgebiete (2), 34. Springer-Verlag, Berlin, 1994.

[Ray70] M. Raynaud: *Spécialisation du foncteur de Picard.* Inst. Hautes Études Sci. Publ. Math. 38 (1970), 27–76.

[Reg80] C. J. Rego: *The compactified Jacobian.* Ann. Sci. Ècole Norm. Sup. (4) 13 (1980), no. 2, 211–223.

[San09] F. Sancho de Salas: *Koszul complexes and fully faithful integral functors.* Bull. London Math. Soc. 41 (2009), 1085–1094.

[Ser06] E. Sernesi: *Deformations of algebraic schemes.* Grundlehren der Mathematischen Wissenschaften 334. Springer-Verlag, Berlin, 2006.

[Sch98] D. Schaub: *Courbes spectrales et compactifications de jacobiennes.* Math. Z. 227 (1998), no. 2, 295–312.

[Tei80] B. Teissier: *Resolution simultanee. I.* Exposé n. 8 in Seminaire sur les Singularites des Surfaces (Palaiseau, France 1976-1977, M. Demazure et. al., eds.). Lecture Notes in Math., vol. 777, Springer- Verlag, Berlin and New York, 1980.